


No. A22: 6. 22 18
GLASGOW
UNIVERSITY

LIBRARY.

WALKER-ARNOTT COLLECTION.





LECTURES

ON

HISTOLOGY,

DELIVERED AT THE

ROYAL COLLEGE OF SURGEONS OF ENGLAND,

IN THE SESSION 1850—51.

ELEMENTARY TISSUES OF PLANTS AND ANIMALS.

BY JOHN QUEKETT,

ASSISTANT CONSERVATOR OF THE MUSUEM OF THE ROYAL COLLEGE OF SURGEONS OF ENGLAND.

ILLUSTRATED BY ONE HUNDRED AND FIFTY-NINE WOOD-CUTS.

LONDON:

HIPPOLYTE BAILLIERE, 219, REGENT STREET,

AND 290, BROADWAY, NEW YORK, U.S.

PARIS: J. B. BAILLIERE, RUE HAUTEFEUILLE.

MADRID: BAILLY BAILLIERE, CALLE DEL PRINCIPE.

1852.

LONDON:
Printed by Schulze and Co, 13, Poland Street.

PREFACE.

ALTHOUGH the Leetures contained in the following pages have already appeared in the "Medical Times" for 1851, it has been thought advisable to publish them in a separate form, and for this purpose they have been carefully revised, and a considerable amount of new matter has been added.

The last two Leetures have been omitted in this volume, but will be included in a subsequent one, as they relate to the Structure of the Skeleton of Invertebrate Animals, which formed the subject of the Course of Leetures for 1851—52.

The preparations exhibited, were, with few exceptions, those described in the first volume of the Histological Catalogue, to which work the Author would refer those of his readers, who may be desirous of having a more detailed account of the individual specimens.

The Author begs to tender his best thanks to his friend, Dr. P. B. Ayres, for much valuable assistance during the progress of the work.

^{32,} Blandford square, APRIL 27, 1852.



HISTOLOGY OF VEGETABLES.

		· LECTURE I.								
		,						PAGE		
INTRODUCTION	٠.							1		
MEMBRANE								6		
FIBRE .		•		.•				13		
•		LE	CTURI	E II.						
FORMS OF CELLS								17		
CONTENTS OF CI	ELLS	* *		•	. •			26		
		. TE	CTURE	III.	•					
STARCH .			**					28		
CHLOROPHYLLE			• .					34		
MILKY JUICES							•	35		
FIXED OILS						·		37		
VOLATILE OILS							•			
RAPHIDES						•		40		
					•			4.7.		

LECTURE IV.		
SILICA		PAGE 49
LECTURE V.		
SCLEROGEN		
PHYTOZOA	•	70
LECTURE VI.		
FIBRO-CELLULAR TISSUE		
DOTTED OR POROUS TISSUE		~~
T TAMILITE TITE		
LECTURE VII.		
VASCULAR TISSUE.—SPIRAL VESSELS		
POROUS OR DOTTED DUCTS		
LECTURE IX.		
CONCLUSION OF VEGETABLE TISSUES		109
HISTOLOGY OF ANIMALS.		
LECTURE X.		
ANIMAL TISSUES		115
MEMBRANE		117 120

	LEC	TURE	XI.			
						PAGE
YELLOW FIBROUS TISSUE						124
AREOLAR TISSUE						127
	LECT	FURE	XII.			
CARTILAGE .					•	135
	LECT	URE	XIII.			
ARTICULAR CARTILAGE		٠.		•		143
	LECT	TURE	XIV.			
ARTICULAR CARTILAGE.	-synov	IAL ME	MBRANE		•	151
	LEC'	TURE	XV.			
CARTILAGE.—ENCHONDRO	OMA			•		163
OSSIFYING CARTILAGE						170
MOLLITIES OSSIUM						175
			•			177
	LEC	FURE	XVI.			
ADIPOSE TISSUE .				•		180
	LECT	URE	XVII.			
FATTY DEGENERATIONS						193

LECTURE XVIII.

	LEO.	E O 16 12	A V 1.1	1.	
PIGMENT					PAGE . 202
		·····	·····		
INDEX					911

HISTOLOGY OF VEGETABLES.

LECTURE I.

HISTOLOGY,* the science of the minute structure of the organs of animals and plants, may be truly said to be the creation of the present century; some glimpses, however, of organic structure had been obtained by the earlier observers, Leeuwenhoek, Malpighi, Hooke, Grew, and others; but these, for the most part, consisted of unconnected observations, from which it was impossible to educe any of the general laws of formation and development. The microscope, except as a mere toy, fell into disuse during the greater part of the eighteenth century; nor was it until within the last twenty years that the instrument was rendered capable of yielding such a magnifying power, together with such clearness of definition, as is necessary for the investigation of this most interesting and important field of research.

^{*} From ιστος, a tissue or web; and λογος, a discourse.

One of the primary and most invaluable results brought out by elaborate examination of the minute structures of organized beings is, that even a closer unity of organization exists among these structures than is found among the larger organs of animals, visible to the unassisted eye. While little, if any, anatomical analogy is discoverable between the larger parts of animals and plants, the strongest ties of organization are demonstrated by the microscope; and it is found, as we shall hereafter see, that an unity of plan in structure and development, may be traced throughout the organic world.

In organized beings, nature works out her most secret processes by structures far too minute for observation, unless with the assistance of the microscope. Hence we find that our best modern works on human and comparative physiology, arc filled with descriptions and illustrations of minute structure. The processes of sccretion, of nutrition, of generation, nay, even the mysterious actions of the brain and nervous system, unintelligible, except in their results, by the gross means of investigation heretofore employed, are now being gradually evolved by the labours of microscopic physiologists. Nay, more, what is of greater importance, having discovered the healthy structure of the organs, the microscope is brought to bear on the changes of structure of these organs, in their abnormal conditions, so that we now have not only a microscopic physiology, but also a microscopic pathology.

greatest possible aid is thus afforded to seientifie medieine in the diagnosis of diseases, that in many eases would be undistinguishable without the assistance of this instrument. It would be easy to quote numerous examples of the utility of the microscope, in establishing a sound pathology, but it will suffice for my present purpose to eite urinary deposits and tumours, the nature of which is most readily determined by microscopie examination.

Among those who have employed the microscope not only with the greatest assiduity, but with the utmost benefit both to science and their fellow-creatures, have been the members of the medical profession. It would be foreign to my purpose to point out the many advantages to be derived from a knowledge of the intimate structure of organized bodies, in health and disease, as this is amply demonstrated, not only by the publications of the day, but also by the increasing demand in all parts of the globe for this kind of information.

However striking the difference between an animal and a plant may seem at first sight in the higher groups, a more extended examination shows that animals and plants gradually approach each other as we descend in the seale until they meet in a common centre—the simple or individual cell. At this point, all means of distinction between the vegetable and animal organism end, and no feature exists which, in the present state of science, can enable even the most distinguished microscopist to determine to which of

the two kingdoms the individual cell belongs, since it possesses characters common to both. Zoologists, botanists, and chemists, have, each and all, bestowed the labour of years on the solution of this question, and have endeavoured, but without success, to establish some decisive character by which animals and plants may be distinguished.

In the earlier periods of natural history, the power of spontaneous motility, and the presence of a stomach, were considered as the distinguishing characteristics of an animal. It is now well known that certain confervæ possess the power of locomotion, while the sponges and some allied families of animals, are destitute of both these attributes. Chemists in former times, considered the presence of nitrogen as a distinct proof of the animal character of the tissues containing this element; but more modern researches have demonstrated that nitrogenized matter is a necessary constituent of the growing parts of plants. Histological inquiry, as I have already stated, has proved equally incapable of solving this question; the appeal to the higher powers of the microscope having rendered the problem more complex by the discovery of a common in place of a distinctive character; namely, the primary cell, or starting point for all organic beings.

It being demonstrated by histological examination that the cell is the primordial condition of the animal and the plant, it is essentially necessary that the student of histology should be himself familiar with the primary elements discoverable in both these grand divisions of the organic world.

The vegetable kingdom is divided by the philosophical botanist into two great classes, the cellulares and the vasculares; the former containing the lowest, and therefore the least complicated forms. In this class the fungi, algæ and lichens, are composed of simple cells alone, occasionally elongated or otherwise modified in shape in the higher types of each order. The lowest form in each of these orders is a simple globular or ovoid cell. As we proceed, two or more cells are united in a definite Still further development occurs in higher groups until we find distinct organs, which, in the highest of the algae and lichens bear a striking similitude to the leaves and seed-vessels of the vascular or flowering plants. Some orders of alga, the Desmidea and Diatomaceæ for example, are equally claimed by the botanist and the zoologist, so uncertain is it to which department of science they truly belong.

In the vascular class of plants, the structure is more complex, the organs are numerous and serve distinct purposes, and their elementary tissues have been divided into cells, fibres, and vessels: this subdivision has been proved by recent investigation to be illusory; vessels being merely modified or elongated cells; and fibres, elongated cells, the walls of which are thickened, or the entire cavity solidified by subsequent deposit on the internal surface of the cell wall.

The lowest types of animal life consist, like the analo-


gous vegetable forms, of simple cells; but the progress of development in animals is much more rapid than in plants, producing a greater variety of external form and eomplexity of structure among groups which would otherwise hold a parallel position in the two kingdoms.

One great distinction between the elementary tissues of plants and animals, which should be always kept in view by the student is, that while in the plant the cell, however modified in form, still possesses all the characters of a eell; in the animal it usually undergoes a development into tissues, in which the cellular form completely disappears, and the cellular origin of the tissue can only be discovered by studying it in its embryonic condition.

MEMBRANE.

It is now generally considered by vegetable physiologists that, with the addition of an intercellular mucus or cambium, all plants are made up of a membrane existing in the form of cells or utricles, having as an organic basis a tough insoluble material called cellulose. Membrane, therefore, is regarded as the sole element of plants. Some years since, fibre was also enumerated as an element, but more recent observation has shown that this is always subsequently deposited on the inner surface of membrane.

Membrane in its earliest stages is thin, transparent, and structureless, as shown in the outer membrane of the seed of a *Gourd*, Fig. 1. B, or at A, which represents eells obtained from the young flower-stem of a *Leek*. It is generally eolourless, or has a greenish-white hue; in some cases, however, as in ferns, it is brown. The beautiful and varied eolours of the eorollæ of flowers do not depend upon the membrane eom-




A, cells from the flower-stem of a Leek; a, cell-wall; b, nucleus. в, cpidermis of the secd of the gourd, composed of cells.

posing the eells, but on the eolouring matters which, chiefly in a fluid state, are contained in their interior. Membrane is also, at the stage at which I am describing it, entirely free from visible pores, although fluids readily pass through it by endosmose and exosmose; and, notwithstanding many cells and vessels, hereafter to be described, have their walls studded with dots and apparent foramina, the membrane is always present, but, in most instances, so excessively thin, that it

cannot be demonstrated until charred, or stained of a brown colour by the application of tincture of iodine.

Examples are not unfrequently met with in which cells or vessels having become old and dry, the membrane has disappeared, and holes have been left, and such is the ease in the vessel from a *Balsam*, Fig. 2, A.



A, old cell from the *Balsam* exhibiting perforations. B, fibrous cells of *Sphagnum* with perforations.


Truc foramina, however, are found in the walls of newly developed eells in certain mosses of the genus *Sphagnum*. These cells (Fig. 2, B), are of a fusiform shape, have a spiral fibre developed within them, and exhibit a series of perforations on each side, which, were first described by Mr. W. Valentine. I have satisfied

myself that they are true perforations by observing infusoria pass in and out of the eells in a young plant growing in water.

In process of growth, the thin membranous cell-walls become thickened by the deposition of new matter, either on their external or internal surface—more frequently the latter. The deposit is termed secondary, and, according to Schleiden, usually assumes a spiral direction. It is generally considered, that if the deposit take place before the eell has completed its growth, its form will be that of a spiral fibre or band, in eonsequenee of the cell-wall elongating or growing more rapidly than the deposition takes place within the eell; but that, if the deposit take place after the complete development of the cell, it may then so happen that certain parts being left uncovered by the deposited material, pores, or pits having the appearance of pores, are produced. We have striking examples of eells eontaining spiral fibres, or bands, in many plants, especially the Orchidacea, as in the Pleurothallis, two cells from the leaf of which are shown in Fig. 4, E F, and in some elongated cells from the stem of a Balsam in Fig. 4, B.

The other form of deposit is more eommon, and is well exhibited in its earliest stage in the two specimens of the root of *Marchantia polymorpha*, in Fig. 3, B, C; or in A, which is a representation of the cells forming the pith of the *Elder*; the same fact is strikingly shown in the plant furnishing the Rice paper, and in the

Vine, as at Fig. 3, D; or it may be seen to still greater advantage in the elongated cells known as porous or



A, pitted or dotted tissue of the pith of the Elder. Bc, pitted ducts from the root of Marchantia polymorpha. D, pitted cells of the Vine (Vitis vinifera).

dotted ducts, specimens of which are readily obtained from the *Alder*.


In all these examples there are no apertures in the membrane or cell-wall; but in some old cells from the stem of a *Balsam*, Fig. 2, A, the membrane has been destroyed, and holes are left. This destruction of the cell-membrane is particularly evident, both in the cells and vessels of many specimens of fossil wood, which no doubt had undergone some slight

decomposition before being silicified. Two preparations are described in the Histological Catalogue of the Museum of the College of Surgeons, H 58-59, in both of which the membrane has been destroyed. In the one a series of oval holes, and in the other only an irregular hexagonal network, indicate the original porous type of the vessels.

Membrane may also be thickened by a deposit so dense as nearly to resemble bone. We have examples of this in the stones of the plum, peach, cherry, &c., and more especially in a nut known as vegetable ivory. This hard material called sclerogen by Turpin may occur in a homogeneous form, or in concentric strata, but the deposit being rarely if ever sufficient to fill the cell entirely, leaves a central cavity with radiating canals or pores, which, upon section, resembles one of the lacunæ of bone; but in these sclerous vegetable tissues the radiating canals never pass beyond the walls of their proper cells, and never anastomose with those of neighbouring cells. The thickening of the walls of cells is well shown in the vertical and horizontal sections of one of the scales of the cone of Pinus Webbiana, represented in Fig. 4, G H, in both of which the pores radiating from the central cavity, and proceeding as far as the cell-wall, are distinctly seen; in Fig. 4, D, is shown a large ligneous cell from the Snake-wood, in which the deposit has taken place in concentric strata.

The specimens already described are instances of

deposit on the internal surface of membrane; examples, however, are not wanting in which the deposit has



A, hair from the fornix of Anchusa italica. B, cells from the stem of a Balsam. c, tubercles from the hairs of Anchusa. D, ligneous cell of the Snake-wood. E F, cells from leaf of Pleurothallis. G H, vertical and horizontal sections of cells of a scale from the cone of Pinus Webbiana.

taken place on the external surface, but these are rare in comparison with those illustrating the former mode of deposition. In the hairs of the fornix of *Anchusa italica*, a portion of one of which is represented by Fig. 4, A; the outer surface of the thin violet-eoloured membrane is studded with oval tubercles, some of which are shown in Fig. 4, c; these were first pointed out by Schleiden.

FIBRE. 13

In a portion of cutiele taken from an *Aloe*, the tubercles are so large as to give the membrane a rough appearance even to the naked eye.

FIBRE.


Fibre, although at one time regarded as an element of vegetable tissues, is now generally looked upon as a secondary formation deposited within the walls of cells, or, in other words, on the internal surface of membrane. It is solid, sometimes transparent, and generally speaking of a greenish-white colour, though in some few plants, as in the elaters of Jungermannia, it is more or less red. Its direction, as shown in Figs. 5 and 6, is most frequently spiral, the spires running from right to left; in some plants, however, the direction is from left to right. According to Lindley, it is straight in the lining of the anther of Campanula and Digitalis purpurea. Fibre varies considerably in size; in some plants it exceeds $\frac{1}{3000}$ th of an inch in diameter, in others, it surpasses in delicaey the finest hair; it generally adheres very firmly to the sides of the ecll-wall, especially in the young state, when the turns of the spiral are very elose together, but as the cell elongates, the fibre sometimes keeps paee with it, and the turns are more widely separated. Two or more contiguous turns are very prone to anastomose, and it is in this way that all the peculiar markings found in eells are formed, such as the rings, dots, and reticulations hereafter to be described. When the spiral cells are fully developed, as shown in Fig. 4, E, the fibre will separate from the membrane of the eell-wall, and its elasticity is such, that it ever tends to unroll itself, and in old elongated eells or vessels, it frequently ruptures the membrane, and becomes more or less straight. In this way the fibre may be obtained in an isolated condition.

The bulbs of many of the lily tribe, and one species of squill from the Mediterranean, contain such immense numbers of spiral vessels, that the entire series of laminæ of which they consist, appear to be composed of spiral fibres. These fibres are obtained in such abundance from some plants, that they are collected, bound into bundles, and used as a kind of slow match for lighting pipes and eigars. The elasticity of spiral fibre is very bcautifully shown on the testa of the seeds of many plants, as the Collomia grandiflora, Salvia pratensis, &c. When a minute portion of the testa is moistcned with water, eells that were so elosely adherent as not to be distinguished become soft, the spiral fibre released, as it were, from its elosc captivity, uneoils itself and pushes the cell-wall before it, and what was at first a brown shapeless object, becomes a mass of beautiful spiral cells; a few of these cells from the Collomia are represented in Fig. 5.

Fibre, like membrane, is frequently increased in

FIBRE. 15

thickness by successive deposits, so that the spiral thread becomes a flattened band or even a ring, like a



Cells from the testa of the seed of Collomia grandiflora.

quoit, as in Fig. 6, which represents cells of *Opuntia* vulgaris, with fibres in various stages of development.



Cells with thickened fibres from Opuntia vulgaris.

Some of these cells exhibit thickened fibres, others have the fibre converted into a flattened band; whilst in a few may be seen the ring-shaped bodies affording evidence of concentric deposit.

LECTURE II.


CELLS.

In the first Lecture it was stated that the most simple plants are composed of cells or utricles, and that all others, however complicated in structure, are made up of a series of such cells, variously modified by pressure, and arranged according to the conditions under which they are developed and the functions they have to perform. Within the transparent membrane of which the young eell alone eonsists, there is in the growing eell a thin delicate lining, termed the primordial utricle of Mohl; within this is a nucleus or eytoblast, and this again contains smaller cells or nucleoli. The typical form of the cell is either spherical or oval; but by pressure in growth, eells assume almost every variety of shape, and their walls become thickened either uniformly or partially. In the most lowly organized plants among the Alga and Fungi, the eells maintain their original form throughout life; but in the majority of instances,

as has been already stated, they undergo much alteration in shape. The following are the chief forms which have been specially named by botanists; the oblong, lobed, square, prismatic, cylindrical, fusiform, muriform, sinuous, stellate, filamentous, &c.

In pulpy fruits, such as the ripc strawberry, the cells may be readily separated from each other, and obtained in an isolated condition; many of them, in the lemon, will be found to be upwards of half an inch in length; in the shaddock they are much larger, but in most fruits and other parts of plants, the agency of maceration, or boiling water, must be resorted to for their separation.

One of the most striking, and at the same time most interesting examples of the typical form of cell, is the *Torula cerevisiæ*, or yeast-plant, shown in Fig. 7,



a, three cells of Torula cerevisiæ; b, cells of Torula diabetica; c, cells of Torula cerevisiæ developing new cells.

a. This, like the Torula diabetica so constantly present during the fermentation of diabetic urine, is the carliest condition of a fungus, very nearly allied to the common Mucor or mouldiness. When yeast is examined by a high power of the microscope, it is seen to consist of myriads of minute simple ovoid cells. Shortly after,

being placed in contact with a solution of sugar or infusion of malt, these cells develop smaller cells from one or two points of their external surface, Fig. 7, c, which,

when full grown, give the plant the appearance of a necklace. The further development, which cannot be fully described here, proceeds to the formation of jointed tubes and filaments, when these rise to the surface, and are exposed to the air, the complete development into the Penicillium glaucum occurs. Another species of Torula is found in the urine of diabetic patients as soon as fermentation commences; hence it has been named T. diabetica. For the sake of comparison with T. cerevisiæ, a few of these cells are represented in Fig. 7, b; they are much smaller, and more oval in shape, and are beginning to be arranged in a linear series. This is always found to be the case both in new beer and in urine, if kept for a few days.

There is a disease of the stomach characterised by a very constant train of symptoms: dilatation with flatulent distension, a burning or scalding sensation referred to the epigastrium and upwards in the course of the œsophagus, and the frequent vomiting of a clear fluid, with or without the admixture of a small quantity of brown, frothy matter, like yeast. The patient complains that the vomited fluid is excessively acid to the taste, and its acidity is readily demonstrated by litmus-paper. Within an hour or two after ejection, it begins to ferment as distinctly and rapidly as new beer wort, and a head is formed which cannot be distinguished, in appearance or odour, from yeast. When placed under the microscope, it is seen to be mainly composed of minute cubical masses of a greenish


eolour, intersected by erueial lines, and these are again divided and subdivided by similar markings. bodies were first noticed by Professor John Goodsir, and by him described in the "Edinburgh Medical and Surgical Journal," in 1842, under the name of "Sarcina ventriculi," and he eonjectured that they were of the nature of parasitie vegetables of low organization. They much resemble the infusorial animaleules depicted by Ehrenberg, under the name of Gonium pectorale, so named from an ornament worn on the breast of the Jewish high-priest. Mr. Busk has since noticed them, in the "Microscopical Journal," as occurring in three cases under his care in the 'Dreadnought' hospital-ship, one being a patient suffering from rupture of the diaphragm; and my late brother met with and described a ease at one of the meetings of the Microseopieal Society; but in this instance they evidently belonged to a distinct species of the same genus.

The specimen in my possession was obtained by Mr. Monekton from a patient in King's College Hospital, under the eare of Dr. Budd, from whom we may expect an elucidation of the most effective plan of treatment of this serious disease. I am inclined to think that the administration of remedies best calculated to destroy vegetable life, or fermentation, will prove most effectual in removing this affection; although, hitherto, the disease has generally been found in eases which have terminated fatally.

Two masses of the Sarcina are represented in Fig. 8,

21

A, and B, the former magnified eight hundred diameters,



A, cubical masses of Sarcina magnified 800 diameters. B, cubical masses of Sarcina magnified 250 diameters.

a, Torula occurring with the Sarcina; bb, cells of Torula diabetica and cerevisiæ; cd, cells of the Gonium pectorale.


is shown at A. In the same fluid (Fig. 9, a) we have a great number of minute oval eells, resembling the Torula of the Yeast-plant, which are probably the spores or sporangia of the Sarcina. The Gonium pectorale is represented in Fig. 9, cd, for eomparison with the Sarcina. The eolour of the Gonium is always a bright green, and in the fresh state individuals may be seen moving rapidly across the field of the microscope. As they form the nearest approach to the Sarcina, a representation has been given. In the vomited matters many other substances are detected under the microscope, such as fat, starch, museular fibre, &c.; but with these we have nothing to do at present.

The probability of the development of the Sarcina ventriculi from the minute eells, observed in the vomited

fluid, leads me to repeat the remark that the latter bears a strong analogy to what has been ealled the T. diabetica, represented in Fig. 7, b, and Fig. 9, b, in form and in eapability of exciting fermentation. The T. diabetica and T. cerevisiæ, are neither of them perfect plants, but an intermediate and subaquatie stage of the development of fungi, belonging to the order Mucorinæ. Turpin has demonstrated that the Yeast-plant, misnamed T. cerevisiæ, is an early condition of the Penicillium glaucum; and my friend, Dr. P. B. Ayres, has informed me that he has traced the development of the T. diabetica to a species of Mucor, probably Mucor ramosus. Each oval cell or spore, first developes one or two smaller cells, by germination on opposite points of its surface, these increase in size, and each again produces other cells, so as to form a moniliform string of eells, or torulæ. The next stage of development is into jointed eonfervoid, tubes more or less irregular in figure, and these increasing in number form eloud-like masses, which ultimately rise to the surface of the fluid by the aid of bubbles of carbonie acid, and the upper surface being exposed to the air partially dries, and in a short time the vesical fruetification of the fungus is produced. The genus Torula ought elearly to be abolished, its so-called species being simply stages of development of a more perfect plant.


The ripe *Strawberry* affords an example of larger cells of an oval shape (Fig. 10) containing a brownish nucleus, *a*, *b*; and in those from the young flowering

stem of the Leek, represented by Fig. 11, the nucleus a, and nucleolus b, are easily recognized.



Cells from a ripe Strawberry: ab, brown central nucleus.


FIG. 11.



Cells of a *Leek*: a, nucleus; b, nucleolus.

It would, I fear, occupy too much time to exhibit to you each of the varieties of cells before enumerated; but, in this section of the leaf of a *Balsam* (Fig. 12), and


FIG. 12.



Vertical section of the leaf of a Balsam; a, cells from the stem of the same plant.


in the mass a, from the stem of the same plant, most of the forms may be distinguished, more especially the

square, the oblong, and the lobed; in the section of the rind of a *Gourd*, represented by Fig. 13, you will notice, that eells more or less eubical may become prismatic, or even be so much compressed towards one edge, as to assume the appearance of fibres. There are, however, some of the varieties which demand a more



Vertical section of the rind of a Gourd.






Stellate cells from the young petiole of Sparganium ramosum.

particular description; one of the most remarkable of these is the stellate eell, which is found in the pith of some rushes, the petiole of the sweet Burrreed (Sparganium ramosum), and in the stems of many aquatic plants. In the young petiole of the Sparganium there are strata of hexagonal eells which form septa; these, as shown in Fig. 14, have around their margins certain notehes, which, with those in adjacent cells, from triangular apertures, known to botanists as lacunæ; in process of growth these notehes become deeper and deeper, the lacunæ, aa, enlarging at the expense of the area of the eell-wall, which gradually


assumes more and more the appearance of a star, as shown in Fig. 15. In the Rush we have a very beautiful example of this form of tissue, the eell being

FIG. 15.



Stellate cells from fully developed petiole of *Sparganium ramosum*; aaa, lacunæ.


FIG. 16.



Stellate cells from the stem of a Rush; aaa, black lines indicating the cell-walls.


reduced to a six-rayed star, as shown in Fig. 16, but the remains of the eell-wall, indicated by the black lines, aa, are always present. By this alteration in the shape of the cells, the stem becomes lighter, and in this way the large receptacles for air, eommon in most water plants, are formed.

FIG. 17.



Elongated cells of the Mushroom.

Very striking specimens of the filamentous form of tissue are obtained from many of the fungi, as the common edible *Mushroom*, in which the cells are of a long oval figure, and joined end to end, as shown in Fig. 17; but in the *Boletus igniarius* FIG. 18.



Elongated cells of Boletus igniarius.

they are so greatly compressed and elongated as to assume the appearance of fibres; many of them, as shown in Fig. 18, are seen to bifurcate; but a transverse line, indicating the cellwall, is occasionally visible.

Another variety of cell arrangement is known as the muriform, from its resemblance

to that of bricks in a wall; it forms the medullary rays of most woods, and gives to them the peculiar appearance known as the silver grain; such cells are readily seen in the *Coniferæ*.

Other forms, especially the sinuous, occur in the cuticle of most plants; the cylindrical, however, is best seen in the *Characeæ*.

From the form and structure of cells, we proceed to consider the nature of their contents; these are colouring matter, starch, oil, raphides, silica, &c. I have already stated, that the beautiful variety of colours seen in the corollæ of flowers is dependent, not upon the membrane composing the cells, but upon colouring matters, usually of a fluid nature, contained within them. The cells of the cuticle of the common garden Rhubarb afford an excellent illustration of this fact; some of them filled with a red coloured fluid occurring singly among others destitute of colouring matter. In the corolla of a Pelargonium, most of the cells are full

CELLS. 27


of crimson fluid, which imparts to this favourite flower its rich hue. In the case of the ferns before alluded to, the brown colour depends on that of the cell-wall, and not on its contents. In old cells of *Palms* and certain *Coniferæ*, resinuous matter of rich colour is often met with; this, however, is solid, and is very remarkable in the palm yielding the substance called dragon's blood. The green colour, so universally present in plants, is due to a more or less solid material, contained in cells and termed chlorophylle, or green vegetable wax; it consists of minute spherical or oval particles, but as these have certain properties in common with starch, they will be more particularly described with that substance.

LECTURE III.

STARCH.

Starch, which performs a similar office in the economy of plants to that of fat in animals, is a most important ingredient in the aliment of the whole human race, twothirds of mankind subsisting almost exclusively upon it; its existence was known to the ancients. Leeuwenhoek first examined it microscopically in wheat and beans, and has given tolerably accurate representations of the granules in the second volume of his works. Starch is found in the cells of plants in the form of granules, as in Fig. 19, q, from the Rhubarb, the cells of which have been separated by maceration; also in a section of a Potato. It is usually obtained by rupturing the cellwalls and washing out their contents with cold water; when dry it is hard, and if rubbed between the fingers, a peculiar crackling sound is produced. Starch is insoluble in cold water, alcohol, ether, volatile and fixed oils; boiling water, however, changes its properties, causing

rupture of the granules and the liberation of starch, gum or dextrinc. For this reason the starch granules



a, Granules of Wheat-starch; b, Sago-meal; c, Rice-starch; d, Potato-starch; e, Tous les mois; f, bursting of starch grains on the application of heat; g, isolated cells of the Rhubarb containing starch granules.

almost wholly disappear in boiled potatoes. When examined microscopically, starch is found to consist of more or less oval granules of various sizes, having definite and peculiar characters in different plants; thus in the *Tous les mois*, Fig. 19, e, they are of very large size; in the *Potato*, Fig. 19, d, they are much smaller; and in the *Rice*, Fig. 19, c, are so very minute as to require a power of two hundred and fifty diameters to discover them.

Each granulc has at one extremity a circular spot, termed the hilum, around which are a large number of eurved lines, passing at first in circles eoncentrically, and subsequently in eurves around the hilum, these markings are not, as has been imagined, the result of consecutive deposits, nor indications of increments of growth of the granule itself; they are confined to the cell-wall, and are most probably merc tranverse puckerings or rugæ in the membrane, of which, together with its amylaceous contents, the starch granule consists. If sulphurie acid, and heat be gradually applied to starch, the granules swell to three or four times their ordinary size, and the distended cell-walls lose all trace of the markings which previously existed. When starch is boiled in water, the cell-membrane swells and bursts; the first stage of which is shown in Fig. 19, f, and the amylaecous matter becomes intimately mixed or dissolved in the fluid.

The presence of starch, however minute in quantity, can be immediately recognized by the addition of free iodine, which forms with it a compound of a beautiful blue colour, the iodide of starch—this test was discovered by Jurine. A little tineture of iodine added to the grains of starch in a section of the potato will make each granule blue, but will not so colour the vegetable cells in which these granules are stored up. If a small quantity of starch be boiled and poured into a vessel of cold water, the addition of a few drops of tineture of iodine will give the water a rich blue colour; if a

little of this water be examined under the microscope, the colour will be seen only in the remains of the granule-membrane; and if the water be allowed to stand, all the colouring matter will fall to the bottom of the vessel as a precipitate, showing that all parts of the starch grain are of greater specific gravity than water.


I have stated that the stareh grains have definite and peculiar characters in different plants; and as starch, under eertain names, such as Tous les mois, tapioea, sago, arrow-root, &e., is largely used by invalids and ehildren, it becomes of importance for us, as medical men, to be able to deteet frauds which are frequently praetised in these articles. In Fig. 19, a, b, c, d, e, &e., you have representations of the principal varieties of starch, all of which are drawn to a fixed seale; they, therefore, exhibit their peeuliar form and proportionate size; and it will be readily seen that by means of the microseope, any mixture of two or more of them, or of other varieties with them could be easily detected, as was strikingly exemplified, some years since, in a case which our distinguished ehemist, Dr. Ure, was employed to investigate. This was an attempt to import the stareh of the Cassava, which bears a duty, as arrow-root, upon which there is no duty; thereby an attempt was made to defraud the revenue to a eonsiderable amount. The difference between the two starches was not sufficiently obvious to the naked eye; but by means of the microscope, and a knowledge

of the nature of the grains, the Doctor was enabled to detect the imposition.

Could it be proved that a direct relation exists between the size of the starch granules and their nutritive quality, we should, of course, possess a ready method of determining such quality in any given specimen; and, on this principle, *Tous les mois* would be considered as much more nutritious than the potato, and the potato than the rice starch.

Starch possesses the property of polarizing light; each grain shows a black cross surrounded by coloured rings, changing to a white cross, with the complimental colours of lower intensity when the analysing prism is revolved.

In the euphorbiaeeous plants, the starch granules




a, elub-shaped stareh grains of *Euphorbia splendens*; b, three of the laticiferous vessels with stareh grains in situ.

have a peculiar form and situation; they are found in the interior of the laticiferous vessels, floating in the latex or milky juice, Fig. 20 b, and their shape is that of an elongated dumbbell, or two-headed club, as shown at a. They are easily exhibited by wounding the plant, and placing a drop of the milky juice under the microscope, and

these singular granules are proved to be really stareli, by the addition of tineture of iodine, which gives them a deep blue tinge.

Before leaving the subject of starch, allusion may be

made to the recently prevalent and destructive epidemic among the *Potatoes*, which I believe to have been a disease of the tuber, not of the haulm or leaves. Examined in an early stage, such *Potatoes* are found to be composed of cells of the usual size, but they contain little or no starch; and hence it may be inferred, that the natural nutriment of the plant being deficient, the haulm dies, the cells of the tuber soon turn black and decompose, and *fungi* are developed as on most other decaying vegetable substances. A portion of healthy *Potato*, shown in Fig. 21, consists of a series of more



Portion of healthy *Potato* with stareh grains *in situ*.





FIG. 22.

Portion of diseased *Potato* showing absence of stareh.

or less hexagonal cells, full of granules of starch of different sizes; the granules, when highly magnified, exhibiting the peculiar markings represented in Fig. 19, d.

In the diseased tuber, illustrated by Fig. 22, it will be noticed that the starch-grains are absent from nearly all the cells, their place being occupied by numerous minute

granules of a brown colour. In many of the cells, a large circular nucleus or cytoblast is present. In other



Portion of diseased Potato with fungus.


parts of the same *Potato*, in which the disease is further advanced, not only is the starch absent, but a moniliform *fungus*, as shown in Fig. 23, occupies many of the cells, and threads its way between others. In this section, the parts containing numerous brown granules exhibit the greatest number of *fungi*.

The next, and one of the most common of all the cell-contents, is known as *chlorophylle*. It has been before alluded to as the cause of the green colour of plants, and exists in the leaves and young stems of almost all the flowering plants, when not deprived of the action of light.

Chlorophylle is soluble in alcohol and ether, but is not acted on by water. It forms a thin coating to the granules found in the interior of eells, some of which are said to be composed of mucilage, and others of starch, as is proved by their being rendered blue by iodine. That most remarkable phenomenon, the circulation of the contents of vegetable cells, or, as it is technically termed, cyclosis, may be examined at the same time as the chlorophylle granules; the specimens selected for this purpose being either transparent water-

plants, such as the various species of *Chara*, *Nitella*, and *Vallisneria*, or the hairs of the *Groundsel* and *Tradescantia*.

The stem of the Nitella consists of cylindrical cells, in most of which a movement of granules may be observed; they pass up on one side of the cell and down on



Portion of leaf of Vallisneria spiralis.

the other, but never escape into adjoining cells. The circulation may, however, be more plainly seen in the Vallisneria; a thin layer of one of the flattened leaves of this plant will exhibit a series of oblong cells, as shown in Fig. 24, in each of which an active circulatory movement

of the green granules may be noticed; among the granules so moving may be discerned one larger and more transparent than the rest—the nucleus or cytoblast of the cell, having a nucleolus within it.

I shall now proceed to describe very briefly some of the other contents of cells, such as milky-juices, oils, resins, &c, all of which are important to mankind, some being universally employed as food, others extensively used in the arts.

Milky-juices are true secretions found in many plants, and contained in distinct vessels called milk or laticiferous ducts, formed by the union of elongated cells, which,


unlike other vessels in plants, are eommonly subdivided or branched. The *Dandelion* contains a large quantity of this juice; Indian-rubber, or eaoutehoue, is the concrete milky juice of the *Ficus elastica*, and many species of this genus secrete a similar milky fluid.

Gutta-pereha resembles eaoutehoue in its origin, but the milky-juiee that yields it, eontains a more eonsiderable amount of stareh. As this material is now eoming into such general use, it may not be out of place to give you the derivation of its name, as I have never yet seen it eorreetly stated in books. On referring to a map of Asia, you may observe that the Island of Sumatra presents a ragged outline. The Malay name for this island is Pulo-Pereha, "pulo" signifying "island," and "pereha," "ragged." "Gutta" is the Malay term for gum, hence this substance received the name of "gutta pulo percha," or gum from the ragged island. It has, however, become the eustom of late years to omit the word "pulo," as in the ease of Penang, which was formerly ealled Pulo Penang, and thus this substance has received the name of gutta-pereha. The above information was communicated to me by a friend residing in Penang.

Our countryman, Nehemiah Grew, was not only aware of the presence of milks in plants, but, in his anatomy, speaks of many which "bleed a lympha." He even gives drawings of the vessels which contain it, but does not appear to have been cognizant of their

branched character. These vessels, which have been particularly investigated by Schultz, were supposed by

FIG. 25.



Milk-vessels from the stipule of Ficus elastica.

him to exhibit a circulation; but, it has been found, by subsequent examination, that the circulatory movement is due to the contraction of the vessels themselves, which have always a tendency to expel their contents when torn or divided.


In the stipules of Ficus elastica, the Indian-rubber plant, a layer of branching laticiferous vessels exists, of which Fig. 25 is a drawing; these vessels contain granular matter. In another specimen taken from the Dandelion, the milk-vessels are

readily recognised by their frequent bifurcations, and by their contents having been changed to a brown colour by the liquid used to preserve them.

Fixed Oils and other matters are found as contents of cells; they are highly nutritious, and are chiefly met with in the seeds of plants, in which they serve the purpose of nourishing the embryo, until the development of the radicle and plumule has provided organs capable of deriving sustenance from other sources. In

the Cocoa-nut, the cells contain a concrete oil, which when extracted by pressure, is an article of considerable commercial importance, being much employed for combustion in lamps, and for the manufacture of candles. In the process of germination the oil is absorbed by the embryo, so that the albumen of the seed is gradually

FIG. 26.



Portion of a *Cocoa-nut* with cells full of globules of concrete oil.

diminished and shrivels, a fact well exhibited by a Cocoa-nut after germination. It is somewhat difficult to demonstrate the oils, in situ, in the cellular tissue, since the section almost necessarily allows the escape of the oil; but, in some parts of the Cocoa-nut this is concrete, and we have a better chance of displaying it in the cells.

Some of these are represented in Fig. 26. The oily matter consists of minute spherical globules, which are so closely packed as to render the section partially opaque.

The presence of oily matters in vegetable cells was known, and even depicted by Leeuwenhoek, in the second volume of his works.

Palm Oil resembles fat rather than oil, and is used by the natives of the western coast of Africa as a substitute for butter. It is now extensively employed in the manufacture of candles, and likewise on the railroads, for greasing the wheels of the carriages, sulphur being added to prevent it from melting too rapidly. This oil is contained in the cotyledons of the seeds, from which it may be extracted by heat and pressure; when sulphurie acid is added to it, copious fumes of gas are evolved, and the oil turns black, it is then subjected to distillation, stearic acid comes over, and the residue is a material like black sealing-wax. This is the principle of Price's patent process. Human fat is capable of being separated by pressure into a solid and fluid portion; the solid principle is termed margaric acid, the liquid glycerine. In animals as well as in vegetables, the oil or fat is produced by and stored up in the interior of closed cells.

Castor, croton, linseed, rape, poppy, and numerous other oils, are examples of these vegetable fats. An oil expressed from the seeds of a species of Jatropha, burns exceedingly well in lamps, and gives a steady white light. It is manufactured in Bristol by Messrs. Visger and Miller, and sells at four shillings per gallon; but it is of a poisonous nature, and cannot be even tasted with impunity. A bag of the nuts having accidentally burst on the quay at Bristol, many children picked them up and eat them, and in a short time numbers of the children were taken into the infirmary with symptoms of irritant poisoning. Some of them suffered severely, others less so; but I believe no fatal result ensued in any ease. A gentleman, well skilled in the properties of oils, imprudently dipped his finger into a specimen of this oil, and carried it to his mouth, in order

to judge of its quality; he was rendered uncomfortably ill for two or three days.

Nor is it in the animal and vegetable kingdoms alone that we meet with oil, since it has also been obtained from the mineral kingdom. Mineral oil derived from bitumen is of a dark colour, somewhat resembling brandy, and burns exceedingly well in lamps. The paraffine extracted from peat by destructive distillation, by Mr. Owen's process, is employed for making candles. Peat appears to be capable of conversion into a great variety of useful products, many of which are at this time gaining much attention from practical men.

Volatile Oils are also secreted in cells, which, from their peculiarities of shape and situation, have received the names of vesicles, vittæ, glands, &c. They occur on the surface or in the parenchyma of leaves, and in the pericarp of fruits. They are not nutritious, are soluble in alcohol, but not in water. The volatile oil of the rind of the lemon and orange, is contained in such vesicles, and connoisseurs in punch are well aware, that by rubbing a lump of sugar over the outer surface of the lemon, the sugar absorbs the essential oil, and thus imparts a fine flavour to the liquor; the sugar being rough acts the part of a grater, rupturing the cells in which the oil is contained, and allowing it to escape.

Many of these essential oils, when evaporated, yield a resinous substance. The resin may be sometimes seen on the surface of plants yielding a perfume. There is a striking example of this in the seed of the common

lavender, which, when examined as an opaque object, presents rows of small globular bodies of highly refractive powers situated in the furrows of the seed. If the seed be dipped in alcohol, the resinous matter will be immediately dissolved.

Camphor is a substance nearly allied to the volatile oils, differing from them in being solid at ordinary temperatures. It may be found in larger or smaller masses, in the cracks and fissures of the wood of Laurus camphora, as in a specimen belonging to John Hunter, which is still preserved in the Museum.

Gum is another substance, formed originally in cells, but subsequently separated from the plant as an excretion. The purest form of gum is that known as the gum arabic, produced by the Acacia vera, which is very soluble in water, forming a transparent solution; the gum tragacanth, from the Astragalus tragacantha, cannot be said to dissolve, but forms a soft opaque mucilage, resembling paste.

Wax and Tallow are also found in plants, the former in large quantities on the surface of the Wax-palm, Ceroxylon andicola, whilst the latter occurs in the seeds of Pentadesma and Bassia, of Sierra Leone, both having the specific name of Butyracea. The bloom of the plum and the grape are also examples of wax.


Resins abound in some plants, especially in the coniferous species; they are not unfrequently contained in large reservoirs, termed turpentine vessels. When that from the common Pine is distilled, oil of turpentine

comes over, and the residue is known as the "black" or "fiddler's resin." If, however, water be added prior to the distillation, the residue is the white resin of commerce.

Canada balsam now so largely used for mounting microscopic objects, is an example of a liquid resin.

RAPHIDES.—Besides the various contents of cells which have already occupied our attention, there are certain mineral substances commonly found in them in the form of crystals. These were first noticed by Malpighi, in the Opuntia; and were subsequently described by Jurine and Raspail. They occur principally in two forms, either in stellate masses, as shown in Fig. 27, B, C, or in bundles of sharp-pointed crystals, very much resembling needles, as in Fig 27, A. It is these last which have obtained for themselves and their fellows the general name of raphides, from the Greck Paφις, a needle. They are also not unfrequently found as single crystals, in the form of octohedra, rectangular, and oblique prisms. According to Raspail, the needleshaped, or acicular, are composed of phosphate, and the stellate of oxalate of lime. There are others having lime as a base combined with tartaric, malic, or citric acid. These are easily destroyed by acetic acid, and are also very soluble in many of the fluids employed in the conscrvation of objects. Some of them are as large as $\frac{1}{40}$ th of an inch, others as small as $\frac{1}{1000}$ th; they occur in all parts of the plant, in the stem, bark, leaves, stipules, sepals, petals, fruit, root, spiral vessels, and even in the pollen. They are always situated in cells, and not as


has been stated by Raspail and others, in the intercellular passages. Some of the containing cells become much elongated, but still the cell wall can readily be traced.



A, portion of outer layer of bulb of Scilla maritima, having acicular raphides in some of its cells. B, conglomerate raphides in the cuticle of a Cactus. c, stellate raphides in root of Rhubarb. D, raphides from the bark of the Lime-tree.

In a thin section of the leaf of an aloe, the *Aloe verrucosa*, small silky filaments may be discerned by the naked eye, which, when magnified, are found to be bundles of acicluar raphides. In the bulb of the medicinal squill, *Scilla maritima* (Fig. 27, A), certain large

cells are detected immediately beneath the euticular layer, filled with bundles of acieular crystals.



A, octohedral crystals of oxalate of lime in the cells of an *Onion*. B, prismatic crystals of oxalate of lime in the cells of another part of the same *Onion*.

In the eutiele of the Onion, every eell is oeeupied by an octohedral or
prismatic erystal of oxalate
of lime. In one specimen,
Fig. 28, A, the octohedral
form predominates; whereas, in others, Fig. 28, B,
the erystals are ehiefly prismatie, and are beginning
to form stellate groups.


All who are in the habit of examining the deposits from urine must be familiar with the appearance of the erystals of oxalate of lime, and will readily reeognize their close resemblance to these in the eells

of the *Onion*. I have also met with them in the urine, contained in the interior of eells, so that, both in the animal and vegetable kingdoms, we have erystals of oxalate of lime as a product of eell secretion.

Raphides of oxalate of lime are found in very great abundance in the medicinal *Rhubarb*, as shown in Fig. 27, c; the best specimens from Turkey, containing as much as thirty-five per cent. of them, those from the

East Indies twenty-five per eent., and the English or that sold in the streets by men dressed as Turks, ten per cent. Buyers of this drug usually judge of its quality by its grittiness, that is, by the quantity of raphides it contains, and this is the more curious, as the erystalline matter eannot from its nature, contribute to the activity of the drug, since the tincture which contains no raphides, is as efficacious as the powder. Certain plants, of the Cactus tribe, seem to be almost entirely made up of raphides; in some instances every eell of the cutiele, as shown in Fig. 27, B, contains a stellate mass of crystals; while in others, the whole interior is full of them, rendering the plant so exceedingly brittle, that the least touch will oeeasion a fraeture. On this aecount, some specimens of Cactus senilis, said to be one thousand years old, which were a few years since sent to Kew Gardens from South America, were obliged to be packed in cotton, with all the eare of the most delieate jewellery, to preserve them during the transport. Raphides of peculiar figure are eommon in the bark of many trees. In the hiccory, Carya alba, masses of flattened prisms,

FIG. 29.




Prismatic raphides in the bark of the Lime-tree.

having both extremities pointed, may be observed; similar erystals are present in the bark of the *Lime-tree*, and in Fig. 29 they are represented in situ; they occur in rows, their pointed extremities nearly touching each

other, their principal situation being in the cellular tis-

sue, in contact with the medullary rays. Other forms

FIG. 30.




Prismatic raphides in cells of the pith of *Elæagnus angusti*folia.

FIG. 31.



Raphides in the bark of the *Apple-tree*.

FIG. 32.



Raphides in the testa of a seed of the Elm.

of crystals, as the rhombohedron, and a small stellate form, shown in Fig. 27, D, are also found in the bark of the lime. In a vertical section of the stem of *Elæagnus*, represented in Fig. 30, numerous prismatic raphides of large size are visible in the pith. Raphides are also found in the bark of the *Apple-tree*, as shown in Fig. 31, and in the testa of the seeds of the elm, as shown in Fig. 32; each cell contains two or more very minute crystals.


It is not at present known what office raphides perform in the economy of the plant. Some have gone so far as to state that they are deposits to be applied towards the mineral part, or skeleton of the plant, but the fact of their being insoluble in vegetable acids would prove this view of their use to be erro-

neous. The more rational supposition is, that they are accidental deposits, formed by the action of vegetable acids upon lime or other bases, derived by the

plant from the soil. Many of the larger kinds of raphides, as those from the Cactus, Fig. 27, B, do not consist entirely of carthy material, if they be acted on by an acid just strong enough to dissolve the lime, it will be found that an organic basis is left behind retaining, to a certain extent, the shape of the mass of crystals. Some of the masses even exhibit a laminated centre with crystals on their exterior, and when mounted in Canada balsam, they very much resemble small calculi, leading one to suppose that this was their true nature. They may, however, be formed artificially, and my late brother succeeded in doing so in the following manner: If oxalic or phosphoric acid be added to lime-water, the precipitate will be pulverulent and opaque. If, however, a vessel containing oxalate of ammonia in solution be connected by means of a few filaments of cotton, with another vessel containing lime-water; crystals will be formed at the end of the fibres in contact with the lime-water. This led him to attempt to form them in the interior of cells. He selected for the purpose a portion of ricc-paper; this substance was placed in limewater under an air-pump, in order to fill the cells with the fluid. The paper was then dried, and the process again and again repeated, until many of the cells were charged with lime-water. Portions of the paper were then placed in weak solutions of oxalic and phosphoric acid, and at the end of three days, crystals were found in the cells in both instances, those in the oxalic acid being

of the stellate form, while those in the phosphoric acid were rhombohedral. No acicular crystals however

FIG. 33.



Portion of *Rice-paper*, in which are two masses of raphides produced artificially.

No acicular crystals, however, were formed although the process was continued for ten days. One of these pieces of *Rice-paper* exhibits very clearly a stellate mass of crystals in some of the cells. Each of these, as shown in Fig. 33, precisely resembles the raphides found in *Rhubarb*.

Raphides are sometimes found in the fossil state. In a section of a fossil palm, from Saugur, in the Deccan, a stellate mass, evidently of a crystalline nature, occupies nearly the whole of the interior of some of the larger cells.

LECTURE IV.

SILICA.

Besides the raphides, another and more insoluble inorganic material, silica, is met with abundantly in certain orders of plants, not in a crystalline form, nor contained in the interior of cells, like the raphides, but diffused generally throughout the structures in which it occurs, and this connexion is so intimate and equable, that it forms a complete skeleton of the tissues after the soft vegetable matters have been destroyed; in fact, the part it plays with reference to the organized tissues in which it is deposited is precisely analogous to that existing between the animal and earthy elements of shell. Silica exists in such great abundance in the cuticle of a plant known as Equisetum hyemale, or Dutch rush, that on this account the stems are employed by carvers in wood and modellers in clay as a substitute for sand-paper.

When the stems are rubbed together, a grating noise is heard, as if they were composed of glass. In the Graminaceæ, especially the canes, silica is also very abundant, but is by no means limited to this order of plants. It is contained principally in the cuticle and in the various structures that are developed from it, such as hairs, spines, &c.; but in some instances layers of cells lying much deeper than those of the cuticle also abound in silica, and it may be met with in woody fibres and in spiral vessels.

In certain of the canes, as the bamboo, silica sometimes collects in large solid masses in the interior of the joints, forming the substance called "tabasheer." It would seem that in this case, the silica should be viewed in the light of a secretion, as it is poured out in a liquid state into the cavity of the bamboo, and I possess a specimen in which a spiral vessel, lying in the cavity of the joint, has been entirely surrounded by the tabasheer.

It is generally known that after the burning of haystacks, masses of irregularly-shaped, but perfectly formed glass, are always to be found among the ashes; these result from the fusion of the silica contained in the cuticle of the hay, in combination with the potash of the vegetable tissue, by which a silicate of that base (or glass) is formed.


In a portion of the bark of a tree from Ccylon, the name of which I have never been able to ascertain, the

amount of silica is so great as to cause it to be employed in the manufacture of pottery.

In order to display effectually the silieeous matter in plants, it is necessary to expose the tissue under examination to the flame of a blowpipe, or better still to boil it for some days in nitrie acid. By these means the organic portion is entirely destroyed, and the siliea withstanding these destructive agents, remains as a perfect model or cast of the original tissue.


The stem of Equisetum hyemale, after having been

FIG. 34.



A portion of the cuticle of *Equisetum hyemale*, after long boiling in nitric acid.

FIG. 35.



A portion of the euticle of a leaf of the Bamboo.


boiled and macerated in nitrie aeid for a eonsiderable period, is a mass of pure siliea, and, as represented in Fig. 34, not only do the forms of the eells of the euticle remain, but even a eonsiderable amount of the detail of the stomata, as shown in the rows of small oval bodies with serrated markings. In the leaf of the bamboo, Fig. 35, of which one part A has been imperfeetly, and another B perfectly deearbonised, an exact model, or east in silica, of the original specimen still remains; the more complete the decarbonisation the whiter is the siliea. On the surface of that part of the leaf, shown at B, even

FIG. 36.



A portion of the husk of a grain of Wheat.

FIG. 37.



A portion of the cuticular and subcuticular layer of cells of the *Manilla cane*.

the hairs are plainly exhibited. In the husk of a grain of wheat, Fig. 36, not only the cells of the eutiele and layers of cells beneath, but also the fibres of the spiral vessels are silicified.

The smooth, glassy outer surface which many of the varieties of eane present, is entirely due to silica; for if a thin slice of the cutiele be removed and exposed to the flame of a blowpipe, or to the action of nitric acid as in Fig. 37, from the Manilla cane, not only every cell of the cutiele remains beautifully defined, but in this instance, the layer of cells immediately beneath; these have no nuclei, and are seen at the

bottom of the figure, the cutieular layer having been removed from this part.

Of all the grasses used as food by man, rice contains the largest proportion of silica. This is demonstrated by boiling the husk of the rice grain in nitric acid to remove the organic matter, after which the microscope gives evidence of the large quantity of silica contained in the husk. This circumstance accounts for the worn and flattened surfaces of the teeth of those races of men FIG. 38.




Portions of woody fibres from the husk of the *Rice*.

that live almost entirely on raw rice; examples are found in the skulls of Malays in the Museum of the College of Surgeons. This condition of the teeth frequently enables us to distinguish the animal or vegetable nature of the food by which the individual has been supported.

In the husk of the rice, the woody fibres are also coated with silica. In most specimens the woody fibres are abundant, some of them, as seen in Fig. 38, presenting peculiar serrated margins somewhat like those of the fibres of the crystalline lens of the cod-fish; others, as shown at b, are smooth.

of the Rice. As I have already stated, the modifications of the epidermic cells of plants partake largely of the siliceous deposit, and this is particularly the case with the hairs or setæ which stud the surface of the husk of the oat, wheat and other grains. I may here refer to a very interesting pathological circumstance in connexion with these minute and seemingly insignificant hairs. The occurrence of intestinal concretions of anomalous character, and of no ordinary size, was at one time far from uncommon in this country, especially in the north of England and in Scotland, and many such specimens are preserved in the Museum of the College of Surgeons. Some of them are of considerable bulk. One of these was


brought to the College many years since, for the pur-FIG. 39. pose of ascertaining its nature and the ra-



Hairs from the palea of an Oat. bread is extensively used as an article of diet.

The Pharus cristatus, an exotic grass, is remarkable

FIG. 40.




Portion of the cuticle of a leaf of Pharus cristatus.

for the arrangement of its silicified cells. In this grass, two rows or masses of silica, somewhat resembling an hour-glass in shape are arranged in two sets; between these, stellate masses like raphides, arc interspersed. Both forms are illustrated in Fig. 40, but never having seen any specimen of the

grass in its natural condition, I am unable to form a

conclusive opinion as to their nature. One of the com-




Portion of palea of Festuca pratensis.

mon meadow grasses, Festuca pratensis, here also requires notice. In the paleæ, cups of silica as they have been termed by the Rev. J. B. Reade, are very abundant. In the specimen represented, by Fig. 41, which is a portion of one of the paleæ in the natural state, longitudinal rows

of these cups may be observed; the hairs also on their edges are exceedingly sharp, and abound in silica.

The leaves of most species of *Deutzia* arc remarkable for having stellate hairs developed from the cuticle



Siliceous cutiele from the under surface of the leaf of Deutzia scabra, exhibiting sinuous cells and stellate hairs.

of both their upper and under surfaces; in Fig. 42, some of these are represented. This cuticle, in common

with the other examples already mentioned, exhibits a beautiful series of colours under polarized light.

In the cuticle of the petals of a Mallow Malva sylvestris, nearly every cell contains a stellate mass of calcarcous matter, but as the masses are all crystallinc, they are probably only raphides.

While noticing silica in plants it will be well to make mention of the great abundance of this substance in the beds of our ponds, streams, and rivers, and in the waters of the ocean, formed by the accumulated remains of countless myriads of organized beings; whether these are animal or vegetable is still a matter of contention between botanists and zoologists; the former classing them with the Algæ, the latter styling them Infusoria. They may, however, be regarded as cells, having their walls strengthened by a coating of silica, which like that of the cuticles before mentioned, is probably a secretion, and indestructible both by fire and acids.

According to botanists, the Algæ arc divided into five natural orders, viz.: Diatomaceæ, Confervaceæ, Fucaceæ, Ceramiaceæ, and Characeæ. Of these the Diatomaceæ are characterised as "angular fragmentary bodies, brittle, and multiplying by spontaneous separation." This order is divided into three genera, viz: Cymbelleæ, Hydrolineæ and Desmidieæ; the two first have a siliecous, and the last a horny skeleton. All are remarkable for their beauty of form, as well as for the markings on their surfaces, which are so delicate in many cases, as not only to be employed as a test of the


defining qualities of a microscope, but, in some instances, their minuteness is such as even to defy the highest magnifying powers we at present possess, to render them visible.

The Diatomaceæ inhabit both salt and fresh water; the Desmidieæ fresh water only; these last have lately been classified and arranged by a member of our own profession, Mr. John Ralfs, and in his excellent work you will find all that relates to their structure and mode of development. If the Diatomaceæ be animals, the Desmidieæ are certainly vegetables, Mr. Ralfs having detected in them the presence of starch.

The Diatomaceæ possessing siliceous skeletons, require a brief notice; a good example is the Isthmia obliquata found on our coasts, attached to fuci, and remarkable for the elegance of its form and markings. A single shell is represented by c in Fig. 43. The Isthmia, although having a siliceous shell or covering, is claimed by the botanists, and the first figures of it were given in Sowerby's "British Botany." Many of the Diatomaceæ, on account of their silica, are not only employed largely in the arts, as a material for polishing metal and other hard structures, but in the form of a light white powder, resembling magnesia, are said to be used as an article of food, being known in Germany as the berg-mehl, or mountain-meal. The berg-mehl consists principally of the remains of navicula, one of which is represented by B, in Fig. 43.

In certain parts of the world, as in Germany, and

especially in America, there are strata of infusorial remains, many miles in extent. The city of Richmond, in Virginia, is built upon a stratum eighteen feet deep,



A, Synedra. B, Navicula grandis. c, Isthmia obliquata. D, Campylodiscus clypeus. E, Eunotia. F G, Gallionella sulcata. H, Zygoceros rhombus. I, Dictyocha fibula. K, Cocconeis.

and upwards of thirty miles in length, which is shown by the microscope to consist of little else than well-marked infusorial remains, some of which are represented by E, F, G, H, I, K, in Fig. 43.

These minute ereatures once lived in the water, from which they had the power of separating the silica, in the same manner as their representatives of the present day—as one generation died off, it was replaced by another; and so, in process of time, was a stratum formed.

In Bohemia there is an infusorial earth termed Kiesel-guhr, which is remarkable for containing peculiar shield-shaped discs, termed by Ehrenberg, Campylodiscus clypeus; one of these is represented by D, in Fig. 43.


On the shores of South America, and in many islands of the Pacific, sea-birds, principally penguins, resort and deposit their excrement, which, in the lapse of ages, has accumulated so as to form masses of enormous extent and depth; and thousands of tons have been brought to this country as a manure, under the name of guano. Guano consists of a mixture of flesh, bones, and sand, together with various salts of potash, lime and ammonia. When examined microscopically, a great abundance of beautiful siliceous skeletons of Diatomaceæ are found amongst it; and curiously enough, the best samples of guano contain the greatest number of these remains, which were first detected by my late brother in 1845. The infusorial skeletons are chiefly circular, and many of them of a blue colour; they are the siliceous shells of animalcules once inhabiting the depths of the ocean, and which had been swallowed by the fishes, on which the penguins feed, but had not been digested; they belong principally to three genera: Actinocyclus, Gallionella and Coscinodiscus, and are no doubt of the same species with others still living in the sea in the neighbourhood of the Guano Islands. Now, when we

consider the vast amount of silica that must be removed from the soil with the straw of wheat, barley, oats, and other grasses, it must be evident that a supply of this substance ought to be restored to the soil in order to insure good crops; hence, it follows, that the value of good guano as a manure may depend, not entirely upon its ammonia, lime, and potash, but in a certain degree also upon the silica it contains.

LECTURE V.

SCLEROGEN.


I now proceed to speak of another kind of deposit in cells, which still retains the name of sclerogen, originally given it by Turpin; this deposition always takes place in the interior of the cell, but rarely if ever completely fills it, a central cavity, and certain radiating tubes or canaliculi being left at intervals in the deposit. In the Histological Catalogue of the Museum of the College of Surgeons, all the principal varieties of the deposit of sclerogen, are classified and described under the name of Hard tissues, and contrasted with bone and teeth, which form the hard tissues of animals. Those who are in the habit of eating Pears will remember, that near the centre of the fruit, and in the neighbourhood of the seeds, there is something which grates between the teeth: it is known to botanists as the Gritty tissue,



Thin section of *Pear*, showing a mass of gritty tissue amongst the cells.

and consists of a number of cells aggregated together in small bundles, somewhat in the shape of a star, or a mass of conglomerate raphides. When a thin section of *Pear* is examined with a power of ninety diameters, as shown in Fig. 44, these stellate masses are readily seen amongst the cellular tissue; but when a

higher power is employed, as of two hundred and fifty diameters, and the cells have been slightly separated



Isolated cells from the sclerogenous, or gritty tissue of the *Pear*.


from each other by maceration or boiling (Fig. 45), each exhibits a distinct central cavity with tubes radiating from it, and the solid deposit contained within the cell, is almost as clear and transparent as quartz, and refracts light very strongly. This is the sclerogen of Turpin, and it is the same mate-

rial that gives hardness to the stone of the *Plum*, *Apricot*, *Peach*, &c., to the shell of the *Cocoa* and *Coquilla nut*, and which is so dense and white in the fruit of a *Palm*, *Phytelephas macrocarpa*, as to have obtained the name of vegetable ivory.

Some of these structures are so hard as to be employed in the arts for various purposes; the albumen


of the *Ivory-nut*, and shell of the *Coquilla-nut*, being turned into handles for walking-sticks, umbrellas, and other articles as a substitute for true ivory. The *Coquilla* is a species of *Cocoa-nut*, and full two centuries ago received the name of *Cocos lapidea* from its stony hardness. The *Ivory-nut*, when first imported into this country is so soft as to be readily cut with a knife, and is of a milk-white colour, but it soon hardens, and although so very dense, yet, when cut sufficiently thin, still exhibits its cellular character. It is composed of long oval cells, having large elongated central cavities, from which, numcrous canals radiate towards the margin

FIG. 46.



Vertical section of the Ivory-nut

of the cell, at this point, as shown in Fig. 46, they are considerably dilated. It is curious to observe that the pores or canals in adjoining cells almost invariably correspond at their circumferential extremities, and vet they in no case meet or otherwise communicate, the intervening cell wall being always present, as seen in Figs. 46 and 47. In a transverse section of the same nut, Fig. 47, the size of the central cavity, and the pores coming off from it as



Transverse section of Ivory-nut.

so many radii, are seen; their dilated ends arc well shown, also the fact of their non-communication with adjoining pores, and the presence of the cell wall.

In specimens mounted in preservative liquids, the canaliculi and central cavities become filled with the fluid;

but those mounted in a dry state contain air, and the points above mentioned arc more clearly shown, as both cavities and canaliculi are perfectly black, and they have precisely the appearance of bone cells, the surrounding parts being transparent.

The outer coating or husk of the Ivory-nut is of a

FIG. 48.

Vertical section of the testa, or husk of the *Ivory-nut*.

brown colour, and exhibits a structure quite different from that of the white interior. Some of the cells, as shown at a, in Fig. 48, are, when transversely divided, of hexagonal figure, and consist of a small central cavity, with a very great number of minute equal sized pores or caniculi radiating from it towards the wall of the cell. In imme-

diate connexion with these cells, as shown at b, are others divided longitudinally, in which the system of

pores is well shown. A curious point in this section is, that cells appearing so truly different in structure, should occur so close to each other, and that there should be no gradual transition from one to the other.


In the cells of some plants the deposit takes place in concentric layers. In a section of the Snake-wood, Brosimum guianensis, a wood not unfrequently used in the manufacture of bows for the archer, there are many large ligneous cells, in which the deposit has taken place in concentric laminæ, the pores running through them being small and few in number; and the central spot, although of very minute size, is occupied by a mass of rich brown resinous matter. These cells have been already described, and one of them is represented by D, Fig. 4; the same kind of tissue is found in the eones of the Common Fir, which consist of clongated cells, nearly full of deposit. Four of these, divided vertically, are represented by G, in Fig. 4, and a mass of the same in transverse section by H, in the same figure. In the old ligneous cells of the bark of the Beech, the concentric deposit is remarkably well seen with scarcely any trace of pores, as illustrated by Fig. 49. A singular form of laminated deposit, combined with a branched eondition of the pores or canaliculi, is found in the seed of the Star-anise represented in Fig. 50. It is best seen in specimens mounted in Canada balsam, and when the central cavity and pores of the cell are full of air. The laminæ are well shown in sections of the cells, but the branching of the pores, which are very

FIG. 49.



Old ligneous cells of the bark of the Beech, exhibiting concentric deposit.

FIG. 50.



Cells in the testa of the seed of the Star-anise (Illicium anisatum).

numerous, ean be better studied when the outer surface of the eell is in focus, or when a thin layer of the eells is uninjured in the preparation of the specimen.

A knowledge of these hard structures is often of eonsiderable importance, much more so, indeed, than many are apt to imagine. The following is an example of the praetical utility of such an aequaintance with minute structural anatomy: About two years since, I received from a medical gentleman in the eountry, some specimens mounted as microscopic objects, that had been passed from the bowels by a female. One of them I found to be the cutiele of a plant, and this turned out subsequently to be the cutiele of a Gooseberry; the other puzzled me, but I made up my mind that it also was of vegetable origin, and that it was, in all probability, the testa of some seed. I wrote to my correspondent to this effect, but the patient denied having eaten any dried fruit for the space of twelve years, and the physician, believing the statement of his patient, considered that the

microseopist was in error. I, however, still maintained my point, and, when preparing the series of specimens known as hard tissues, for the Histological Catalogue of the College of Surgeons, I examined, among other things, the *Tamarind*, and in the testa of the seed found the disputed structure. I subsequently learned that the patient was the daughter of a groeer, and might have had free access to the tamarind jar. This is another instance of the value of the microscope to our profession.



Portion of the testa of the seed of the Tamarind.

FIG. 52.




Portion of the testa of the seed of Croton tigtium.

A portion of this specimen is represented in Fig. 51. It consists of a series of elongated cells or woody fibres, which exhibit pores on their margins, and oecur in several laminæ; the direction of the cells in one part of a lamina is often nearly at right angles to that in close proximity to it; the entire testa is exceedingly tough, and no doubt very indigestible. A somewhat similar tissue to that above described exists in the core of the *Apple*, but the cells are much coarser than those of the *Tamarind*.

In the *Croton-oil* seed, the eells are remarkable for their minute size, being much smaller than in any other plant I have yet examined; they, however, exhibit very well the central cavity and radiating pores. A portion of one of

the seeds is represented in Fig. 52.

FIG. 53.




Section of the testa of the seed of a Euphorbiaceous nut.

In certain Euphorbiaceous nuts, the sclcrogen is of a brilliant crimson colour, and in the specimen, represented by Fig. 53, the cells are of a peculiar shape, having indented or wavy margins, and when light is transmitted through such objects, the red colour is so bright as to interfere with distinct definition.


Each cell is nearly filled with the crimson deposit.

In the Walnut, Cocoa, and Hazel-nuts, the same kind of tissue is found; in these, the pores with their central cavities, resemble in a most striking manner the cells of bone; but, as I have so often stated, in vegetables the cell-wall always remains, forming a strong barrier between the ends of the pores or canaliculi of adjoining cells; whereas in bone, the cell-wall disappears, and the canaliculi anastomose.

In the specimen represented by Fig. 54, the cells



Transverse section of the shell of a Cocoa-nut.




Vertical section of the shell of a Cocoa-nut.

in a transverse section of the shell of a *Cocoa-nut* are seen; and in Fig. 55, a horizontal section of the same shell, not only the thickened cells are shown, but close to them a part of a bundle of peculiar vessels, somewhat like those occurring in *Ferns*.

Spiral vessels are oeeasionally met with both in the shells of the Walnut and Hazel-nut.

In addition to the substances already described as examples of the more common contents of eells, a few of a peculiar nature demand some notice.

Certain plants are provided with offensive weapons, some of which are known as stings, being (as in the Nettle) elongated eells or hairs developed from the eutiele, and usually of a eonical figure. They contain a poisonous or irritating fluid, in which, in some eases, a cyclosis or circulation is visible. When examined with a power of one hundred diameters, as represented by a, b, in Fig. 56, each sting is seen to have either



a, b, stinging hairs of the Nettle (Urtica dioica); c, d, e, ciliated spores of Confervæ.

a minute bulbous or triangular free extremity attached by a narrow neek to the apex of the shaft; the part connected with the cuticle is much expanded, and around its base are certain more or less polygonal cuticular cells. The contents of the hair are always in a state of tension, and when the hand is passed along the part of the leaf on which these stings are situated, the free extremity is rubbed off, the point of the sting perforates the skin, and, by the pressure of the eells at the base of the hair, the poisonous matter is forced into the wound in a similar manner to the venom from the fang of a scrpent.

We now arrive at the most highly-organized products of secretion of the vegetable eell, viz.: those especially engaged in the development of future plants; they are ealled *Phytozoa*, or plant-animals, being the representatives of the *Spermatozoa* in the animal. They appear to have been first discovered by Mr. Cornelius Varley, in 1833, in that part of the fructification of the *Chara vulgaris* known as the globule, and have since been noticed by Meyen and Unger in mosses, and by Nägeli in ferns.

Mr. Varley's account of these bodies is so true to nature, that I may perhaps be permitted to extract a short paragraph from his paper in the second volume of the "Transactions of the Microscopical Society:"

"The ripe globule spontaneously opens. The filaments also expand and separate into elusters. They are so numerous, that I have not been able to decide upon their exact number.

"The tube-like filaments are divided into numerous eompartments, in which are produced the most extraordinary objects I have ever observed of vegetable origin. At first they are seen agitated and moving in their cells, where they are coiled up in their confined spaces, every cell having one. They gradually escape from their cells, by what means or through what opening I have not been able to ascertain, and the whole field soon appears filled with life. They are generally spirals of two or three coils, and never become straight, though their agitated motion alters their shape in some degree. They have at their foremost end a filament so fine as only to be seen by its motion, which is very rapid and vibratory, running along it in waves.

"These objects, although they have every appearance of life, swim about with no apparent consciousness of each other, or any choice of direction. Their filamentary end goes foremost; when they eome into contact with each other they become entangled, and their motion is hindered. This does not eause any retreating action or attempt at separation except by accident.


"They appear to be an example of life without selfwill or choice. Their motions gradually slacken, and in about an hour they become perfectly still.

"If a globule be forcibly opened before it is ripe, the filaments will give no indication of life."

Unger has described them in Sphagnum as Infusoria, under the name of Spirillum; they have been the subject of great controversy, and Schleiden, denying their animal nature, has stated that they are nothing more than fibre in an early stage of development. From what I have been able to observe, I think that Schleiden's notion is decidedly erroneous, and that they are strictly analogous to the Spermatozoa.

In some of the lowest plants, as the *Confervæ*, an aet, called by botanists *conjugation*, takes place. Two filaments lie side by side, and very soon a nipple-shaped growth is protruded from opposite points of the filaments, these touch and eventually communicate; one cell becomes empty, and in the other a spore is developed. The spore makes its escape from the parent

FIG. 57.



Conjugation of Conferva; a, spore in process of development.


eell, and swims about very actively by means of eilia attached to some part of its outer surface. In some species there is a pink spot in the centre of the spore, and such spores are no doubt not unfrequently described as *Infusoria*. The conjugation of a *Conferva* is shown in Fig. 57, and at a, a spore in process of development. Spores provided with eilia are represented at

cc and d, in Fig. 56.

If we examine mosses of the genus Sphagnum and Polytrichum, in the spring, we find among the parts eoneerned in reproduction, eertain elongated bodies termed Antheridia. If one of these be ruptured, a number of eells make their escape, and, if this be done in water, in a few moments we notice, with a power of two hundred and fifty diameters, the rapid movement of a spiral filament or phytozoon, in each eell. The filament has a globular head and a long tail, exactly like that of a Spermatozoon. When the phytozoa escape from the


cell, their true form may be made out, and they will be seen to be precisely of the shape exhibited in Fig. 58; a representing one of the antheridia, b a portion of the

FIG. 58.



Reproductive parts of *Polytrichum* commune: a, antheridium; b, portion of an antheridium more highly magnified; c, cells escaping from ruptured antheridium; d, cell containing a phytozoon; e, f, phytozoa detached from the cell.

FIG. 59.



a, a, spores and elaters of an Equisetum; b, spores surrounded by the elaters; c, portion of filament of Chara vulgaris dividing phytozoa in the cells; d, d, phytozoa of Chara vulgaris taken out of the cells.

same ruptured, with numerous eells escaping. Each (as shown at c and d) contains a spiral filament, which, when detached from the cell, presents the appearance represented by e or f. The spores of the common Equiseta or horse-tails also possess a power of spontaneous motion, depending, however, on hygrometric conditions of the atmosphere. Each spore, as shown at a a, in Fig. 59, consists of a central portion with three or four clavate filaments attached to it which are denominated elaters. In some states of the spore, as represented by b, the claters are seen surrounding it in a spiral manner; in others, as shown at a a,

they will be found standing out in straight lines from the spore. But the most curious point connected with these elaters is the rapidity with which they embrace the spore when slightly moistened by the breath. If the moisture of the breath fall upon them, they will immediately eoil themselves around the spore, and almost as quiekly resume their original position, and this may be repeated an indefinite number of times. It has been considered by some persons that the phytozoa resemble the elaters, but that their motion is continuous. the sake of comparison, the phytozoa of the Chara vulgaris are shown in Fig. 59; at c, a portion of one of the filaments is seen, it is eomposed of a series of cells of a somewhat square figure, arranged one above the other; in each cell is a phytozoon. Two of these, detached from the cells, and more highly magnified, are represented by dd.

LECTURE VI.

FIBRO-CELLULAR TISSUE.

WE now arrive at a class of eells which, for the sake of distinction, may be termed compound, from the circumstance of their being composed of membrane and fibre; they occupy definite positions in plants, and form the tissue known as the fibro-eellular. The cells are of a more or less oval figure, and the fibre often attains a very large size, even as much as $\frac{1}{3000}$ th of an inch.

Fibro-cellular tissue is eommon in the leaves of many orchidaceous plants, and oeeurs abundantly in the testa of seeds. I have already mentioned the existence of this tissue on the testa of the seed of Collomia grandiflora, in which the cells containing a spiral fibre are protruded from the surface after the seed has been moistened with water. The leaf of an Orchis, Pleurothallis ruscifolia, exhibits, by vertical section, a large

number of angular eells (Fig. 60) containing spiral fibres. Immediately beneath the cuticle, on the upper edge of many of the sections, a row of small oval cells may be seen, in which the fibre is of very large size,

FIG. 60.




Vertical section of the leaf of *Pleurothallis ruscifolia*, showing fibro-cellular tissue *in situ*.

FIG. 61.



Cells of fibro-cellular tissue of *Pleurothallis ruscifolia* more highly magnified.

and in some eases unrolled. If this layer of cuticle



A, single cell of the leaf of Oncidium stramineum. B, cell from testa of the seed of Cobwa scandens.

be peeled off after some slight degree of maceration, and the attached surface examined, a beautiful view of the fibro-eel-lular tissue, as shown by Fig. 61, is afforded; the terminations of the fibres being seen at the free extremity of the cell. These cells may be isolated from the leaves of most *Orchidaceous* plants, as in *Oncidium stramineum*, Fig. 62, A, one of the finest

FIG. 63.



Elongated cell from the leaf of Saccolabium guttatum with spiral fibres crossing.

examples of the size of the eells and spiral fibre. Detached eells of *Pleu-rothallis ruscifolia* have already been represented by E and F, Fig. 4.

In the leaves of another Orchis, Saccolabium guttatum, there are very long cells, in which the spiral fibres run obliquely, and cross each other at right angles, whereby a series of diamondshaped markings are produced. One of these eells, obtained by maceration, is represented in Fig. 63, as seen under a magnifying power of two hundred and fifty diameters.

The testa of the seed of $Cob\alpha a$ scandens yields, by maceration, the isolated fibrous eells represented in

Fig. 62, B; they are of much smaller size, and contain a fibre remarkable for its extreme tenuity and the number of its coils.

Another form of fibro-cellular tissue occurs in the testa of the almond, in which the fibre is interrupted, and the eells approach in character to those termed porous cells. Similar cells exist in great abundance in certain seeds, whose testæ project laterally, forming thin membranes, or wings as they are called by botanists. In a specimen from the *Sphenogyne speciosa*, represented by Fig. 64, a delicate layer exists, composed of clongated cells of fibro-eellular tissue; in another specional

cimen (Fig. 65), from the seed of a Bignonia, the cells are longer and the fibre coarser than in the pre-



Portion of testa of the seed of Sphenogyne speciosa.

FIG. 65.


Portion of testa of the seed of a Bignonia.

ceding. In the seed of Lophospermum erubescens, in which the thin membranous wing surrounds the entire circumference of the seed, the cells with their spiral fibres are well shown. The most remarkable specimen of wing, however, and one in which this tissue is largely developed, occurs in a plant, from the East Indies, Calosanthes Indica, the wing being more than an inch in length on each side of the seed.

DOTTED OR POROUS TISSUE.


The variety of tissue known as the *porous*, is formed by an unequal deposit of secondary matter on the interior of the cell wall; the spaces pre-occupied by the deposit, being thinner than the other parts, produce the appearance of pits or pores. It is one of the most common vegetable tissues, and is generally met with in the pith and other light parts of plants. The material known as Chinese rice-paper, is not really paper, but a thin shaving of a species of the genus **Eschynomene*. I have often heard ladies complain of the difficulty of getting the colour to lie smoothly on this material, which is readily accounted for by the fact that it is composed of a series of large cells (Fig. 66), the walls of which are minutely porous. A spurious kind of rice-paper, procured from a species of **Desmanthus*, (D. natans*,) is still more difficult to draw on than the last; the walls of many of the cells being dotted with large pores.

FIG. 66.



Thin slice of Chinese rice-paper (Eschynomene).

FIG. 67.



Porous tissue of the *Dragon-* palm.

The most striking example of porous tissue is that from the oldest living inhabitant on the surface of our globe, the *Dracæna draco*, or *Dragon-palm* of Tencriffe. In a section I possess of this *Palm*, the wall of every cell (Fig. 67), however small, is covered with

large pores. In all the examples above-mentioned, the eell wall is invariably present, but there are instances when the eells have become old, of the eell-wall disappearing, and true pores being formed.


WOODY TISSUE.—This tissue, which makes up almost the entire bulk of certain plants, is by far the most useful and important to man of all those occurring in the vegetable kingdom, since it supplies material to every branch of art, and to it we are indebted for our linen, ropes, cordage, &c., to say nothing of paper.

It eonsists of clongated transparent tubes of great strength, but varying much in size in different plants; each tube is more or less pointed at both extremities, and in some eases the tubular part has become almost solid by the internal deposition of new matter, so that the trace of the central cavity is only indicated by a line. Such tubes are remarkable for their toughness, and have received the name of fibres. The outer membrane of the fibre is generally structureless; but in the Flax and other plants, in which the fibres are of great length, there are here and there traces of transverse markings.

The fibres of Flax are aggregated together in bundles, and are separated from the plant for use by a process termed hackling, which may be described as an alternation of beating and combing. In the rough condition, in which it is imported from Russia, the fibres have been, to a certain extent, separated by hackling. On its arrival here, it is again subjected to maceration, to a repetition of the hackling, and to the bleaching process in order to reduce it to the white silky condition

required by the spinner and weaver. Flax, after its final preparation, consists of bundles of structureless tubes (Fig. 68, A) which are seen separated in Fig. 69, each fibre exhibiting a central line.

Similar bundles of woody tissue occur in the "Chinagrass" of commerce, which is said to be obtained from a plant of the Nettle tribe, and is brought to this country in the form of coarse fibres, two of these are represented in Fig. 68, c. After being hackled,



A, bundle of woody fibres of the Flax plant. B, two of the ultimate fibres of the leaf of the Pine-apple. c, two woody fibres of the China-grass plant.

macerated, and bleached, it is made into very delicate shining linen handkerchiefs, the material of which is known as Grass-cloth. The fibres of the Phormium tenax, or New Zealand Flax, are much stronger than those of the common Flax, but are much more easily broken, in consequence of the outer membrane being excessively brittle. Although these fibres will resist a very considerable force applied to them longitudinally, if tied into a knot so that the fibre is twisted, or if the force be applied transversely, they readily break.

Of all the woody fibres I have examined, those obtained from the leaves of the *Pine-apple* plant appear to be the finest. For the sake of comparison, representations are given of the fibre of *Flax* (Fig. 68, A), of *China-grass* (Fig. 68, C), and of the *Pine-apple* (Fig. 68, B), as seen under a power of four hundred diameters, by which it is well shown that the last is much smaller than either of the others.


It is important, under eertain eireumstanees, to determine the true nature and eomposition of some of the textures used as artieles of clothing, &e., and this is readily done if we bear in mind the component threads merely longitudinal, rounded, unmarked fibres, such as have been already figured in Flax, Fig. 69; but where cotton has been employed solely, or by admixture, its fibres, as shown in Fig. 70, are recognised as flattened and more or less twisted bands, bearing other marks of resemblance to hairs, which, in fact, they are, since in the condition of clongated cells, they line the inner surface of the pods. Fig. 70, represents the isolated

elongated cells of cotton, in which the peculiar characters I have described, are delineated, they may be contrasted



Fibres of Flax.

FIG. 70.




Fibres of Cotton.

with the figures of true woody fibre, and also with wool (Fig. 71), and silk (Fig. 72), and it will be



Wool of sheep.

FIG. 72.



Filaments of silk.

seen that these textile materials are readily distinguishable by the microscope.

Mummies have, at various periods, been brought to this country and unrolled, and it has been a matter of euriosity to determine the nature of the clothing material, as giving evidence of that in use in the country and age in which they were embalmed. By microseopie investigation of the texture of mummy eloths, it has been shown that the Egyptians used Flax only in their manufacture, whereas, in the Peruvian mummy cloths, Cotton alone has been discovered. In some fabrics from Otaheite, Wool is mixed with Cotton, and is recognisable, as shown in Fig. 71, by the zigzag transverse markings on the fibre, which are characteristic of hair.

In the foregoing examples of woody fibre, the membranous walls are structureless, having no visible markings; but other kinds exhibit certain peculiarities of structure, which depend on secondary deposits of lignine in their interior. In coniferous woods, such as Deal, the fibres are not filled with secondary deposit, and consequently remain tubular.


In a transverse section of a fossil Palm from Antigua, most of the woody fibres will be found to be occupied by a concentric deposit like that of the Beech, represented in Fig. 49. In the Clematis, the woody fibres are exceedingly short, and more or less pointed at both extremities; the walls, as shown in Fig. 73, are marked with numerous simple pores. These pores are more evident, and of a compound or bordered character, in a section of Deal, Fig. 74, made in the direction of the medullary rays; in every fibre a row of large circular dots or porcs may be seen, which are characteristic of

the coniferæ, each pore having a smaller one in its centre. In some woods, and especially in fossil speci-



Woody fibres of Clematis.

FIG. 74.



Woody fibres of *Deal*, exhibiting bordered pores.

mens of Araucaria, three or more rows of pores occur in each fibre.

Spiral fibres may also co-exist with pores, as in the Yew, (Fig. 75) each woody fibre of which has two spirals interlacing each other, and one or two pores



Section of Yew, showing bordered pores and spiral fibres.

FIG. 76.



Section of Araucaria excelsa exhibiting large bordered pores and spiral fibres.

between the interlacements of the fibres where these are most distinct. The eo-existence of large pores with a well-developed spiral fibre is best seen in a section of the Norfolk Island pine, Araucaria excelsa, (Fig. 76) a tree, which in its own climate, often attains the enormous height of two hundred feet. The nature of the pores, or glands as they have been termed, of the woody fibres of the Conifera has long been a matter of dispute.

The foregoing illustrations have been taken from compact woods, but in certain palms the woody fibres are so hard and stiff, and the bundles so easily separated that they are employed for making brushes. The larger and coarser bundles are made into brushes for sweeping the streets, while the smaller are used for scrubbing and other lighter purposes.

In a section of the stem of the eocoa-nut palm, the bundles of woody tissue are of a brown colour, and form the spots seen upon the surface; the surrounding texture being so compact as to bind them firmly together. A transverse section of one of these bundles affords a striking example of the interior of a woody tube almost entirely filled up by dense secondary deposit, and accounts for the strength of the individual fibres. In many light woods, especially one from China, the name of which I have not been able to obtain, but which from its softness, is employed as a substitute for cork in the lining of insect boxes, the woody fibres are short, of rather large diameter, and, instead of being

filled up with hard matter, as in the cocoa-nut, their interior is occupied with granules of stareh, arranged in single, double, or treble rows; these might very easily be mistaken for pores, but their true nature is readily detected by the addition of iodine.

WOODY TISSUES.


When speaking of the various kinds of woody tissue employed in the arts, I mentioned ordinary Flax, China Grass, and Hemp, and stated that, from the peculiarly hard nature of the outer coating of each fibre of the latter, it was rendered very brittle when twisted. I then remarked that linen is at all times more difficult to work, and to absorb colouring-matter, than cotton, which may perhaps, be accounted for by the density or toughness of the woody fibre; I have since learned that a process has lately been invented by M. Claussen for subdividing the elementary fibres of Flax by boiling and steeping, after they have been hackled and bleached in the usual manner, and specimens prepared in this way are made to imitate not only cotton, but silk and wool. In some samples of Flax which have undergone these processes, the fibre has been reduced to the fineness and suppleness of cotton, in another to the texture of silk, and in a third to that of wool. When, however, these

specimens are examined microscopically, the woody fibres will, in most cases, be found entire; some indeed are ruptured tranversely, and others partially separated into finer fibres. Mr. Donlan has proposed another plan of preparing Flax, which if successful, will in all probability yield a more durable fabric than any now in use. In this process the fibres of the fresh stems are at once reduced to the smallest conceivable size by mechanical means, without previous steeping or bleaching.

I have already stated that woody fibres are elongated cells, generally more or less pointed at both extremities, and having their walls strengthened or solidified by internal deposit. Occasionally, however, as in Flax and Hemp, they are marked with tubercles or tranverse lines at short intervals; in some plants, especially those in which the woody fibres arc short, as the Clematis and Elder, Fig. 73, when marked with pores, or little dots, from deficiency of the internal deposit at certain points, these arc called simple pores; but in the Conifera, represented in Fig. 74, the pores are surrounded by a larger circular ring, giving to the whole the appearance of a circular cell, with a small central nucleus; these are called bordered porcs, and are well seen in all specimens of conifcrous wood. The true nature and mode of formation of these bordered pores was for many years a subject of dispute among botanists; by some they were supposed to be glands; others have considered them to be either thick or thin spots in the membranous cell-wall of the fibre, whilst a few have


asserted that a hole really exists in the eentre of each pore communicating with a similar hole in a neighbouring fibre. It is now, however, ascertained, by improved means of investigation, that these bordered pores are not confined to one fibre, but are formed between two eontiguous fibres, and always exist in greatest number on the sides of the woody fibre parallel to the medullary rays. The bordered pores are hollow, their shape is bi-eonvex, as represented by c in Fig. 77; and a small

FIG. 77.



a, section through two contiguous woody fibres of Salisburia adiantifolia; bc, portions of two contiguous woody fibres of Deal (Pinus Strobus); d, one of the bordered pores of Pinus Strobus magnified 500 diameters.

FIG 78.




A, two woody fibres of a fossil coniferous wood, with spiral fibres and bordered pores; aa, casts of the interior of the bordered porcs. B, fossil coniferous wood with three rows of bordered pores.


circular or oval spot is visible in the eentre of each. In some plants each fibre has one row of pores, in others two and in a few instances, as at B in Fig. 78, which is a representation of a specimen of fossil coniferous wood from Kennawha Ohio, the entire fibre is covered with them. In order to get a satisfactory view of these

pores, a tangental section of the wood must be made; when, as in the specimen of deal (portions of two contiguous fibres of which are represented by b c, in Fig. 77,) it will be seen that there are certain oval spaces between these fibres, which have in their centre a porc extending nearly through the entire thickness of the wall of the fibre, but closed by membrane on its outer margin; a highly magnified representation of onc of these pores, showing the mode in which it projects from the side of the woody fibre of the deal, is shown at d. The correctness of this account of the real significancy of the structure of the *qlandular* woody fibre has been strikingly confirmed by the examination of sections of fossil coniferous wood. Some years since, a specimen of fossil wood from Fredericksburgh, in Virginia, was sent to my late brother by Professor Bailey of the Military Academy of West Point on the Hudson, which, on microscopic examination, was found to be coniferous, the woody tissue not only exhibiting the bordered pores, but, in addition, numerous minute spiral fibres, as represented in Fig. 78, Some of the disc-like bodies, shown at a a in the same figure, and precisely similar to the porcs of the fibres, were lying loose. On carefully examining the surfaces of the fibres, similar discs were occasionally found projecting from the outer surface, as represented in Fig. 78, A. It then turned out that these dies were in reality casts in silica of the bordered pores; in short, the bi-concave cavities shown by b c in Fig, 77, had been filled with silica. Thus the description given of their true nature, after careful observation, was fully confirmed by the fossil specimen.

In some of the orehidaeeous plants, as those of the genus Oncidium, and especially in Aporum anceps, small globular bodies are found projecting from the woody fibres, as represented in Fig. 79, A. Sometimes they oeeur in many rows, as shown at A, or in a single one as at B, or in two rows as at C, each being surrounded by a ring-like marking. These bodies, like those in the fossil specimens, are sometimes found loose, as shown by E, but, the places they occupied, as represented at C, were eavities in the wall of the fibre.



A, woody fibre of Aporum anceps with four rows of globular bodies. B, woody fibre with one row. c, woody fibre in which the cavities occupied by the globular bodies are shown. D, globular bodies projecting from the side of a woody fibre. E, globular bodies detached from the fibre.



A, woody fibre of Salisburia adiantifolia with upper surface in focus. B, woody fibre of the same plant with both upper and under surfaces in focus. c, woody fibre of Cycas revoluta showing at the upper part the porcs on the under, and at the lower those on the upper surface.

In some plants, as in *Cycas revoluta*, the eentral spot or pore is not round, but more or less oval, and is inclined

at an angle of 45° to the length of the fibre, and the pores lying on one surface have the direction of the central spot at right-angles to that on the other surface; the outer or bordering lines eorrespond; but if the thin part of a fibre be examined, it often happens that both the upper and under surface are in foeus at the same time, and then the appearance of a eell with a crucial nucleus, as shown by c in Fig. 80, will be presented. faet is even more strikingly exemplified in Salisburia adiantifolia, the pores being of much larger size. Portions of two of the woody fibres of this plant are represented by A B in Fig. 80. At A the central pore on one surface of the fibre is shown, and at B the two opposite sides of the same fibre, in which the crueial appearance is most strikingly exemplified. At a, in Fig. 77, a section through two contiguous woody fibres of the same plant is represented, by which it will be seen how these peculiar pores are formed.

LECTURE VII.

VASCULAR TISSUE.—SPIRAL VESSELS.

Vascular Tissue.—This tissue, bearing as it does the same relation to the growing portion of the plant as the vascular system of animals does to their general organism, may be considered as the most important part of the plant. It forms no exception to the general rule with which we started, that all the textures originate in cells, for these vessels are but elongated cells joined end to end, and ultimately communicating with each other; their walls are in most cases supported either by a complete or modified spiral fibre. These spiral vessels, were called by the older microscopic botanists, trachea, from their resemblance to the ramifying air tubes of insects; nor is the analogy far fetched, since in either case the tube is kept patulous by an elastic spiral coil of fibre, which has a tendency to unrol itself when

freed from confinement within the vessel. Moreover, in old dry parts of the plant in which their function has ceased, they are found to contain, as in insects, only air, but in the young and growing parts they contain fluid, and floating raphides are found in abundance in the spiral vessels of the young branches or shoots of the common grape vine. Spiral vessels, however, differ from the air-vessels of insects in being generally straight and unbranched.

In order to illustrate what I have been describing, it will be well for you to see the tracheæ of the larva of *Dyticus marginalis*; in any of the large branches, as shown by Fig. 81, the presence of a spiral fibre within a membranous tube, will be remarked. For the sake of comparison, a portion of

FIG. 81.




Portion of a tracheal tube of the larva of Dyticus marginalis.

FIG. 82.



Portion of a spiral vessel of a plant.

a spiral vessel of a plant is represented by Fig. 82, in which it will be seen that the principal point of




Tracheæ ramifying upon the stomach of a *Bee*.

distinction is, the size of the fibre in the vegetable being larger than that in the animal. In both, the fibre is within the membranous tube. In Fig. 83, the tracheæ are displayed ramifying on the coats of the stomach of a *Bee*, I may here remark that the tracheæ serve the purpose of carrying oxygen

to the blood or nutritious fluid, which in insects is distributed through the system generally, and not in vessels.

Spiral vessels are elongated cells, acuminated at each end (Fig. 86, a) with one or more fibres developed within them, and capable of unrolling by the tearing of the delicate membrane of the cell-walls. I have already described the mode of formation of the spiral fibre by a secondary deposit in a spiral direction, on the internal surface of the cell-wall. The spiral vessel is subject to many modifications, in consequence of the variable arrangement of the sccondary deposit, and these modifications have heretofore been classed as distinct orders of vessels—the term spiral vessel having been restricted to such as contain one or more spiral fibres, and are capable of being unrolled. Recent and accurate researches have demonstrated, that all the forms of vascular tissue are referable to the same origin: the simple cell with its secondary interrupted deposit. We have already

seen simple ovoid eells eontaining a spiral fibre; let one of these become elongated and pointed, and we have at once the typical spiral vessel. Other arrangements of the secondary deposit produce the annular, the reticulated, the barred or scalariform, and the dotted duets of botanists. That such is the ease is demonstrated by the occasional presence of the spiral, the annular, the barred, and even the dotted variety in different parts of the same vessel. Nothing is more common in plants, especially among the *Monocotyledons*, than to see the spiral fibre in one part broken up into rings, as in a spiral vessel from the *Opuntia*, Fig. 84, A. Spiral vessels are usually pointed at both



A, short spiral vessels, with thickened fibre, from *Opuntia vulgaris*. B, a spiral vessel becoming an annular one. c, spiral vessel having a hole at one extremity. D, dotted vessel with oblique termination and hole in the eentre. E, reticulated vessel. F, old spiral vessel with perforations. G, H, spiral vessels with fibre running from left to right.


extremities, but occasionally they terminate somewhat abruptly, and a hole is often seen near the extremity,

by which a communication is established with a neighbouring vessel, as in Fig. 84, c, from the petiole of the garden *Rhubarb*. A similar perforation in a dotted duct is represented by Fig. 84, D.

In most spiral vessels the fibre is single, (Fig. 86, a,) it is then called a simple spiral; in others, two or more fibres running in the same direction form a band, which for distinction is termed a compound spiral vessel (Fig. 86, b); sometimes the coils of the fibre separate, and the extremities of each coil uniting form a series of rings; such vessels are called annular (Fig. 84, AB). In other cases, the turns of the spiral may be connected together here and there by the branching of a fibre, or the development of lateral processes, forming what is termed reticular tissue (Fig. 84, E).

Occasionally the spiral fibre bifurcates, or short longitudinal fibres are developed, which connect the spiral coils, and give the vessel a reticulated appearance; such vessels represented by E and F, in Fig. 84, occur in sections from the Balsam. In other cases, the spiral thread is not so evident, but the interior of the vessel (A, Fig. 91) is marked by long slit-like pores, which are placed regularly one above the other, resembling the rounds of a ladder; these are termed scalariform vessels, are almost peculiar to Ferns, and will be alluded to hereafter. That these are really all modifications of the spiral type, is known by the frequent occurrence of several varieties in one and the same plant, and even section, as in the Balsam,

FIG. 85.



A bundle of vessels from the stem of a *Balsam*.

Fig. 85, where in a single bundle, the outermost vessel on one side is dotted, and that on the other annular; between these, spiral vessels occur, with their spires more and more widely separated. This state of things may be also observed in the *Lettuce*, in the root of which plant, bundles of vessels are frequently seen in various

stages of development at one and the same time.


In some plants, and especially in the Canna bicolor, the spiral vessels are remarkable for the occurrence of longitudinal as well as spiral fibres. In Fig. 86, h, is represented what might be taken at first sight for an ordinary spiral vessel of large size, but, if earefully examined, a series of minute longitudinal fibres will also be observed, and when such a vessel is unrolled, the broken ends of these fibres, as represented by c h e, will be found projecting above and below the spirals.

In many eases, both the longitudinal and spiral fibres are of the same size and equi-distant. The vessel represented in Fig. 86, g, is covered with square markings, but retains its capability of unrolling. In another specimen, Fig. 86, f, from the same plant, the longitudinal fibres, although of the same size, are wider apart than the spirals; the vessel in this case presents markings which are true parallelograms, not squares.

These remarkable vessels have amongst them others,

which are unrolled, and occasionally, as represented by Fig. 86, b, you may see very fine examples of the compound spiral vessels before alluded to, as many as






a, simple spiral vessel from petal of Anagallis; b, compound spiral vessel from Canna bicolor; c d e f g h, portions of spiral vessels of Canna bicolor having longitudinal as well as transverse striæ; i, branching spiral vessels from the stem of a Leek.

six coils forming a band; I have, however, in some cases, counted as many as twenty-two. Spiral vessels not unfrequently branch; this fact is well seen in the specimen taken from the flowering stem of the *Leek*, and represented by *i*, in Fig. 86.

In all these examples, it may be observed that the

spiral fibres are uniformly coiled from right to left, but in some instances, this is reversed, and left-handed spirals are found, as in the section of a *Palm* from the East Indies, Fig. 84, G, H; it has been suggested that the direction of the fibre may determine that in which the plant coils round an upright pole. The *Hop*, which exhibits these left-handed spiral vessels is also a left-handed climber, which gives support to the above-mentioned theory.

Whilst speaking of fibres unrolling spirally, I take the opportunity of showing a fact I have lately made out, that even membrane itself may occasionally be met with, that will tear and unroll spirally. Botanists



Hairs from the fruit of Cycas revoluta unrolling spirally.

generally describe membrane as tearing irregularly, but the hairs found on the outside of the fruit of *Cycas* revoluta, which are of some considerable size, will, when separated from the fruit, exhibit the appearance shown in Fig. 87; both extremities are pointed, but it is the one attached to the fruit that unrolls spirally.


I shall next draw your attention to the variety of vessel which, from its peculiar markings somewhat resembling the rounds of a ladder, is called *Scalariform*; but this, like the other varieties, is but a modification and later stage of existence of the true spiral vessel, resulting from an unequal deposit of the secondary matter upon the inner surface of the cell-wall. The parts in which there have been no deposits, or, in other words, the pores, are always more or less elongated, like a slit, and have both extremities rounded. If one of these vessels, when divided vertically, be examined with a power of at least two hundred and fifty diameters, the depth of the pits may be distinctly seen.

These vessels present another peculiarity: they are often of hexagonal figure, and the markings, as well as the sides, are generally of uniform size, as shown in Fig. 88; in other cases, as in Fig. 89, both the sides and the markings are unequal.

The scalariform tissue is best seen in the root of *Ferns*, on tranverse sections of which, the naked eye discerns rows of black dots, with intervals of a lighter colour; the former are bundles of cellular and ligneous

tissue, the latter the vascular tissue of the plant, which is almost entirely composed of sealariform vessels, arranged most frequently in a circular form. An

FIG. 88.



A bundle of scalariform vessels seen in a diagonal section of the root of *Pteris aquilina*.


FIG. 89.



Scalariform vessel of hexagonal figure.

oblique section of the root of *Pteris Aquilina*, or *Common Brakes*, one of the most abundant of our *Ferns* (Fig. 88), exhibits the cut extremity of a bundle of

FIG. 90.



One side of a scalariform vessel from a *Tree Fern* of New Zealand.

these vessels lying among the ligneous tissue; the scalariform, or step-like markings of the walls, are well shown, and also the faet above mentioned, that they are not always eircular in form, but often hexagonal or heptangular.

In Fig. 90, which represents a portion of a vessel found in a vertical section of a *Tree Fern* of New Zealand, these sealariform vessels are ex-

tremely large, being nearly double the size of those of the *Brakes* already described.

So characteristic are these vessels among the Ferns, that I have more than once identified our common Pteris from a small portion of its remains, especially in one instance, in which I found some fragments of a frond in a funercal urn. About two years since an urn, dug up in the island of Anglesey, was, with its contents, brought to mc for examination by one of our most distinguished archæologists, Mr. Albert Way. After having determined the presence of human bones belonging to an adult, and to a child, probably a mother and her offspring, certain filaments were found adhering to the inner surface of the urn; these were of a brown colour, and arranged in definite order like the veins of leaves. Upon microscopically examining sections of these, scalariform vessels were noticed precisely similar to those occuring in the Pteris. This Fern is very abundant in the district in which the urn was discovered, and most probably, portions of fronds were placed in the receptacle before the ashes of the deceased persons were deposited in it.

As I have before remarked, the fact that these several varieties in the vascular tissue are dependent on modifications in the arrangement of the spiral fibre, is proved by the occasional occurrence of several of them in one length of vessel or duct; thus in a specimen from the *Pteris Aquilina* (Fig. 91, B) you will observe one vessel, which at its upper extremity

A B D

FIG. 91. .

A, a portion of a scalariform vessel having nine sides. B, upper portion of a scalariform vessel unrolling in the form of a spiral band. c, middle portion of the same vessel unrolling like an ordinary spiral vessel. D, portion of a scalariform vessel magnified 500 diameters, showing the depth of the pits.

has the scalariform character; lower down, as shown by c, the markings are reticulated; still lower the original spiral form re-appears, and to this again succeeds a portion of ladder-like tissue. When stretched, as represented by the lower extremities of B and c, the scalariform vessels will unroll either in a band, or in single fibres.

I have on a previous occasion spoken of an exception to a general rule which occurs in *Ferns*, namely, that in them the membrane is frequently coloured brown, independently of its contents. When the colour is deep, the vessels collected into bundles, and arranged in a peculiar

form, are easily distinguished by the naked eye.

POROUS OR DOTTED DUCTS.

There are, in addition to the spiral vessels of which I have now spoken, other tubes or canals of large size, termed ducts, which give to woods their different degrees of porosity, and these are visible even to the naked eye. Some of them are marked by an internal deposit of fibre in addition to the pores, but unlike the spiral vessels, they are incapable of unroll-

a b, portions of two porous vessels, or duets, from the stem of a Clematis; c, portion of a porous duet from the Alder; d, pores of the same magnified 500 diameters; e, three bordered porcs from a fossil wood.

ing. The usual markings exhibited are either simple or bordered pores. When these vessels are examined with a low power, the pores appear as dots, and hence the name "dotted duets" has been given to them, the term "duet," being restricted to such vessels as are incapable of being unrolled.

In the Cissampelos Pareira, or Pareira Brava, the duets form bundles, and are remarkable for the minute size of their pores; the extremity of each duct being more or less flattened. In

a vertical section of the root of the *Alder*, the duets as represented in Fig. 92, c, are of large size, and the bordered pores found on their walls, are also remarkable for the extent to which they are developed.

FIG. 93.



Porous vessel in Anthracite coal.

In a section of anthracite eoal, Fig. 93, the duets are of great length, and all the pores are of an oval form. In the eommon *Clematis*, the duets are of small size, but their termination by a septum or septa, situated nearly at an angle of 45° to the long axis of the duet is well shown in Fig. 92, a, b. Now, it often happens that these duets, originally elongated eells, become

eontinuous eanals or vessels, by the absorption of the septa, but, in some cases, in which the process of absorption has either been incomplete, or fibres in the form of parallel bands, have been developed in the septa, these remaining at the edge of the septum, give this part of the duet a very peculiar appearance, somewhat like that of the bars of a gridiron.

In an oblique section of a foreign wood, given me by Dr. Robert Brown, porous duets abound, and the fibres of the septa are well shown; they occur in parallel lines, like the bars of a gridiron, and, as represented in Fig. 94, the fibres join the walls of the pores at the margins of the duets. In a section of a fossil *Palm*, from St. Vincent's, the extremities of the duets are more or less conical, and the fibre is at these points


often developed into broad flattened bands, which are well represented in Fig. 95. One of the most striking examples of the gridiron tissue is to be met with

FIG. 94.



Termination of two porous ducts, the fibres of the septa being transverse and parallel.

FIG. 95.



Termination of two ducts of a fossil Palm.

in vertical sections of the White Birch; in this tree almost all the vessels have septa composed of parallel fibres.

Peculiar forms of porous ducts are met with in the Stigmariæ; they are frequently of great length, and in some specimens there is an appearance of the fibre between the pores of one duct passing across to join those of a neighbouring duct.

All the pores of the ducts I have hitherto mentioned have been either of the simple or bordered kind; but eertain peculiarities occur in some plants which require notice. In the *Clematis*, represented by *a b*, in Fig. 92, an oblique line or marking passes through the centre of each pore; but in a specimen of fossil wood

FIG. 96.



Portion of a duet from the *Elm* showing pores and a spiral band.

from Herne Bay, as described by Mr. Bowerbank,* the markings shown on a large scale by Fig. 92, e, are double, and extend beyond the outer margin of the pore. In all these examples, the markings, as in the case of those of the woody tissue of Cycas revoluta, are produced by the elongation of the central portion of the pore.

In many porous ducts, and more particularly in those of the *Elm*, a spiral

band co-exists with large bordered porcs. A portion of a duct exhibiting this peculiar structure is represented in Fig. 96.

^{* &}quot;Transactions of the Microscopical Society," Vol. 1.

LECTURE IX.

CONCLUSION OF VEGETABLE TISSUES.

HAVING in my last Leeture completed the description of all the important tissues entering into the formation of vegetables, I shall now pass on to eonsider those composing the animal body; but before doing so, I propose to take a rapid review of the subjects which have occupied our attention since the commencement of this eourse.

I stated in my first Lecture, that all plants were made up of an elementary membrane existing in the form of cells, each of which in the young state is provided with a nucleus or cytoblast and nucleoli; the membrane being generally of a greenish colour, but, in *Ferns*, occasionally brown. The membrane in process of growth becomes thickened by secondary deposit; this may be either in a homogeneous form, and occupy the whole

interior of the cell, or be deficient in certain parts, when pits or pores result, or it may occur in a spiral form, the general direction of which is from right to left.

I next called your attention to the various forms under which cells are found in plants, and explained, that in the young state and when unaltered by pressure, they are of a spherical figure, which becomes modified in process of growth into dodecahedrons, cubes, parallelograms, prisms, &c. &c., and even into fibres. I then passed on to the consideration of the contents of these cells, and gave examples of various colouring matters in a fluid state, which produce all the varied hues of the corollæ of flowers. I next noticed one of the most common of cell-contents, and the most useful to mankind, viz.: starch, which possesses definite and peculiar characters in different plants, and consists of granules of a more or less oval figure, composed of an outer membrane exhibiting a central spot or hilum and concentric lines, within which the amylaceous or starchy material is contained. I pointed out the relative sizes of the starch granules in Rice, Potato, and Tous les mois, and drew attention to the peculiar club-shaped bodies found in the milky juice of the Euphorbiaceæ.

Chlorophylle, a substance of a starchy nature, and the material that gives the green colour to all plants growing in the light, was next considered; striking examples of it occur in the *Chara* and *Vallisneria*, in both of which plants the granules may be seen in active circulation within the vegetable cell. Other cell-con-

tents, such as milky juice, oil, resin, gum, were then mentioned, and separately described. Another product of cell-secretion is always found in a crystalline state, and known by the name of raphides. These are of various forms, sometimes occurring as single crystals, but more frequently in stellate masses of oxalate of lime in the *Rhubarb*, or in needle-shaped crystals of phosphate of lime in the *Squill*.

Another inorganic material, found principally combined with the tissue of the walls of cells, is silica, which I described as being so intimately blended with these tissues in certain plants, as to form a perfect cast of their original structure when all the soft vegetable matter has been removed; abundant examples of this being easily obtained from the *Grasses* and *Canes*.

I then exhibited a series of preparations in which sclerogen or hard tissue, approaching bone in many of its characteristics, occurred as a cell-content, but in no one case was a cell entirely occupied by it, a central cavity, with a system of radiating porcs being always present. The last product of cells which I mentioned consisted of *Phytozoa*, or plant animals; these being found in the *Chara vulgaris*, in *Mosses*, *Ferns*, *Confervæ*, &c. Each cell of the antheridia of some *Mosses* is occupied by a spiral filament which exhibits a peculiar gyrating motion, precisely similar to that of the spermatozoa in animals, to which they are no doubt analogous.

The aggregation and modification of the cells previously described, make up the entire structure of all classes of plants, each variety being characterised by the name of tissue; thus we have the woody, the vascular, the porous, and other tissues. The first of these is the most important, since to it we are indebted for our linens, our eordage, and our paper. It eonsists of elongated tubes or fibres of a more or less cylindrical figure, and occurring in bundles; by maceration, and other processes, the fibres are detached from each other, and are then eapable of being worked into fabrics of various kinds.

The vascular tissue consists of eells more or less elongated, joined end to end, or overlapping each other, in which either a spiral fibre or a modification of the same has been deposited; hence, if the spiral be perfect, such a vessel is ealled a true spiral vessel; if interrupted at certain parts, and the fibre breaks up into a series of rings, it is then ealled annular; if the fibre or rings be connected together by branching fibres, in such a manner that a network is produced, the vessel is ealled reticulated; if the fibres be nearly elose together, and the vertical connecting bands short and equi-distant, the vessel is called scalariform, from the markings resembling the rounds of a ladder. Spiral vessels, from their resemblance to the air-tubes of inseets, have been termed Trachea by some authors; the analogy, however, is not far fetched, as in both eases, the tubes are composed of a transparent membrane, kept open by a spiral fibre.

In all the above-described vessels the spiral fibre has

a tendency to unroll, but other eells, which are ealled porous ducts, are of great size, have their walls eovered with large pores, and are not capable of being unrolled. These are the tubes visible to the naked eye in almost every wood. There are other duets found in all plants yielding a milky juice, called laticiferous or milk vessels by Schultz, who, from supposing them to exhibit a eirculation of their contents, likened them to the eapillaries of animals; but by recent examinations, both the existence of the eirculation, and the entire continuity of the vessels, have been proved not to exist.

Such, then, is a brief outline of the subjects that have occupied our attention during the first part of the course. One great object which I have kept in view throughout, has been that of endeavouring to impress on you the fact, that each cell of a plant should be considered as having an independent or individual existence; that in one situation it may secrete colouring matter, in another starch, gum, sugar, oil, &c.; and in another the material for the reproduction of its species.



HISTOLOGY OF ANIMALS.

LECTURE X.

ANIMAL TISSUES.

I now proceed to consider the elementary tissues of animals, and see how far they correspond with those of vegetables. In their earliest condition the cells in both are nearly alike, in some cases the cellular character is maintained throughout life, but in others they rapidly undergo change of form, and all appearance of cell and nucleus is entirely lost. The animal cell, though in some degree possessing an independent existence, cannot perform its functions, as for example, that of secretion, without being in a certain relationship with a series of vessels earrying the nutritive fluid or blood. In the Animal as well as in the Vegetable Kingdom, we have membrane as an element; it may

be met with in the walls of cells, but principally in the form of an investing membrane or sheath, in which case it is perfectly structureless and as transparent as glass. So far as I am aware of, no such extended membrane is found in vegetables unless cellular structure is visible. It is in this coalescence of cell-walls that animal structures principally differ from those of vegetables; in the latter the cell-wall is always present, however old or hard the tissue may be; but in the former, with the exception of those tissues termed cellular, it soon disappears, and in some cases, no trace is left either of nucleus or nucleolus.

It has been found very useful for the purpose of study, to arrange the elementary tissues of animals in a tabular form, and various have been the modes in which this has been carried out by different anatomists. of the best of these tables, and that, which, with some slight alteration, has from the first been adopted in these Lectures, is given by Todd and Bowman in their "Physiological Anatomy." It would, however, be impossible to go through the entire series of tissues in onc course of lectures; I have, therefore, thought it best to follow the order laid down in the above mentioned table, commencing with the simple tissues, and when there is occasion to allude to any one of the compound tissues, a short account of it will be given, in order that its relations to the subject under consideration may be the more readily understood.

TABULAR VIEW OF THE ANIMAL TISSUES.

1. Simple membrane: employed alone or EXAMPLES:—Walls of cells. Posterior layer in the formation of compound mem-
branes colemma of muscle, &c.
2. Fibrous tissues
3. Cellular tissues
4. Sclerous or hard tissues {Rudimentary skeleton of invertebrata. Bone. Teeth, &c.
5. Compound membranes: composed of simple membrane, and a layer of cells of various forms, (epithelium or epidermis,) or of areolar tissue and epithelium.
6. Compound tissues: a, composed of tubes of homogeneous membrane containing a peculiar substance Muscle. Nerve.
b. Composed of white fibrous tissues and cartilage


MEMBRANE.

Membrane is thin, transparent, and structureless, it never presents any visible pores, although fluids pass through it readily; wherever it exists, it is more or less nourished by blood-vessels, which run on its outer surface, but never enter it. The posterior layer of the capsule of the lens of the eye of a Sheep, affords an excellent example of this elementary tissue, which is completely structureless, and as transparent as glass, having no trace of cell in or upon it, so that its very presence can only be discerned by accidental folds, or other irregularities. The capsule of the lens, especially its posterior layer, is in carly life, very vascular, that is to say, vessels ramify freely on its surface, but they do not enter into its forma-

tion or structure. In the adult state, the vessels of the posterior capsule, if they do not altogether disappear, are rarely, if ever, capable of being injected; but in the *Snake*, the *Frog*, and some other reptiles, as will hereafter be shown, they are generally present.


The same simple membrane forms a sheath for the minute elementary particles of muscle and nerve, being called in the former case, sarcolemma, by Bowman, and in the latter neurilemma, by Schwann. The sarcolemma is readily distinguished surrounding a portion of a fasciculus of muscle from an Eel, represented by a in Fig. 97. I have already said that

FIG. 97.



a, muscular fasciculus of an *Eel* showing the sarcolemma; b, blood corpuscles of *Lepidosiren annectens* magnified 500 diameters; c, one of the same seen edgeways.

FIG. 98.




Nucleated cartilage cells from the chorda dorsalis of a *Lamprey*.

simple membrane forms the walls of cells, as in the case of the blood corpuscles, which retain their original cellular character throughout life; those from the Lepidosiren, Lepidosiren annectens, Fig. 97, b, c, are

nucleated, of an oval figure, and with a single exception, are the largest known, being $\frac{1}{570}$ th in the long by $\frac{1}{940}$ th of an inch in the short diameter.

Other examples of elementary membrane occur in the early stages of cartilage, and in adipose tissue; of the former, a specimen (Fig. 98) from the *chorda dorsalis* of the *Lamprey*, bears a remarkable resemblance to the cellular tissue of plants; the nuclei, however, are sometimes wanting. In the cartilage from the ear of the *Mouse* (Fig. 99) the nuclei

FIG. 99.



Cartilage from the ear of a Mouse.

FIG. 100.



Cartilage from the ear of a Bat.


are absent, but in that from the ear of a Bat, *Plecotus* auritus (Fig. 100), they are generally present, each cell having a transparent nucleus in its centre.

FIBROUS TISSUE.

The fibrous, or filamentous form of tissue, enters largely into the construction of tendons, ligaments, fasciæ, and other analogous parts; it is of two kinds, called from their respective tints, white and yellow. Examples of white fibrous tissue are best obtained from tendinous structures, which are almost entirely composed of it. For the yellow, in an isolated condition, we resort to the liquid and subflava of the human spine, or the ligamentum nuchæ of the lower The tissue termed areolar, or the cellular tissue of the old anatomists, is composed of an intricate interlacement of the white and yellow fibres. white fibres, which are most conspicuous in the natural state, may be made to disappear by the addition of acetic acid, so that the yellow fibres, which remain unaltered, can then be separately examined. areolar tissue is more extensively diffused than any other throughout the animal body; it forms the connecting material between organs and tissues, and in some instances is found in such quantities in the Elephant, between the walls of the chest and the costal layer of the plcura, that it may be taken out by handfuls. The arcolar tissue from the thorax of the *Elephant*, affords an excellent object for the exhibition of the nature and appearance of the white fibrous tissue, which occurs

either in white bands, eomposed of minute, equal-sized

FIG. 101.




Tendinous fibres from rectus muscle of the eye of an Ox.

fibres, more or less wavy, or in separate fibres also wavy; these, however, may be rendered straight by extension.

In the tendons of the recti muscles of the eye of an Ox, this tissue is of silvery whiteness, and, as represented in Fig. 101, eonsists of flexuous fibres, all taking the same direction. In some parts, por-

tions of the same tissue are observed erossing the flexuous fibres, these are derived from the areolar tissue with which the tendon is invested.

Having explained and illustrated the nature and eharaeters of white fibrous tissue in tendons, I shall digress somewhat, to describe the eonnexion of muscular fibre with tendons. This connexion is not effected, as has been generally supposed, by union with the investment or sheath of the musele, but, as was first pointed out by Mr. Bowman, by a close connexion of the white fibres with the terminal disc of each fasciculus, as shown in a fibre from one of the recti muscles of the eye of a Cat, represented by A, in Fig. 102. The same mode of connexion is evident in the muscles of invertebrate animals,—as, for instance, in the common Fly. In most works on Entomology you will find that the brown, horny shaft B, with which the muscular fibres are



A, muscular fibre and tendon from rectus muscle of the eye of a *Cat*. B, muscle from the leg of a *Fly*. c, muscular fibre and tendon of a *Fly*.

connected, has been described as the tendon. If, however, the termination of any fasciculus be carefully examined, a perfect tendon of white fibres will be seen to spring from the terminal disc, and by this tendon, represented by c, in Fig. 102, the fasciculus is connected with the horny shaft. The tendinous fibres in some of the lower animals, as the *Mol*-

lusca, are of larger size than those in Mammalia. In the Terebratula they are remarkably large, nearly straight, more or less flattened, $\frac{1}{500}$ of an inch in breadth, and collected in strong bundles, which present to the naked eye a very beautiful silvery aspect. In birds the tendons, especially of the legs, are of large size and great strength, eonsisting of large bundles of white fibrous tissue, connected together by areolar tissue in which the blood-vessels run, as shown in a transverse section of one of the long tendons from the Ostrich. In some birds the tendons are more or less ossified; and all who may have been so unfortunate as to have been helped to the drumstiek of a Turkey or Goose, must be familiar with the long tendons of these birds, which are flattened and tough near their muscular attachments, but bony near their insertions.

When a vertical section of a tendon of any large animal is examined, it always presents a silvery lustre, and the fibres, collected in bundles, proceed in parallel lines, each fasciculus being connected with its neighbour by areolar tissue. If a portion of tendon be dried, the greater part of the silvery lustre disappears, but when viewed by polarised light, not only is the direction of the bundles seen, but each displays a series of brilliant colours.


In the human body we have other structures besides tendons, which are composed almost entirely of white fibrous tissue; such are the periostcum, dura mater, &c. A peculiar form of fibrous tissue occurs in the membrana putaminis or membrane lining the interior of the shell of the egg of most birds and reptiles; it consists of several layers of fibres of nearly uniform diameter, which interlace with each other and produce a series of spaces or cells sufficiently large for gases to pass through. The fibres in an egg of the common Fowl are on an average $\frac{1}{8500}$ of an inch in diameter, and are not acted on by acetic acid. A thin layer of this membrane, having the fibres separated from each other, very much resembles the section of Boletus igniarius, represented in Fig. 18.

LECTURE XI.

YELLOW FIBROUS TISSUE.

The yellow fibrous tissue differs considerably from the white, consisting of large more or less branched fibres, of a yellow colour, with curled extremities. It is found most abundantly, as I have already stated, in the ligamentum nuchæ of quadrupeds, and in the ligamenta subflava of the human spine. In the large Pachydermata, as the Elephant and Rhinoceros, it is frequently employed in the form of a belt or girdle, to support the abdominal parietes. In a specimen, from the ligamentum nuchæ of the Sheep, Fig. 103, A, the fibres are of large size and yellow colour, and much branched, their free extremities being more or less curled. In that from the ligamentum nuchæ of the Giraffe, Fig. 103, B, the same characteristic curled extremities are presented, but each fibre is marked with transverse striæ. When

examined with a power of 500 diameters, the striæ, as shown by c, do not extend across the entire diameter of the fibre, but appear to be principally confined to its centre. The ligament from which this specimen was taken was 6 feet 2 inches in length before it was



A, yellow elastic fibres from the ligamentum nuchæ of the Sheep. B, yellow fibres from the ligamentum nuchæ of the Giraffe. c, one of the same magnified 500 diameters. D, vessels of the ligamentum nuchæ of a young Calf.

separated from its attachments; when detached, however, it immediately contracted to 4 feet, or about onethird of its original length: it even now possesses some degree of elasticity, but the power must be enormous to stretch it even one foot. The weight of the entire ligament was upwards of 8lbs, and the striped fibres are most abundant on the outer surface, the interior being occupied by plain fibres alone. The striped fibres are not confined to the *Giraffe*, I have lately seen them in the *Rhinoceros*, the *Sheep*, and even in arteries.

A modification of the yellow fibrous element constitutes the elastic coat of arteries; which is well shown in a fine specimen taken from the aorta of a Whale upwards of 50 feet in length; the diameter of the vessel is about 12 inches, and the thickness of the elastic coat $1\frac{1}{2}$ inch. When examined by the microscope, the fibres of arteries exhibit all the characters of those of ordinary elastic tissue, except that they are much more minute and less easy of separation.

Another variety of elastic tissue occurs in the ligament supporting the expanded wings of all our larger birds, such as the *Eagle*, *Crane*, *Heron*, &c. The ligament from the *Eagle* is quite as elastic as caoutchouc.

The last variety of this tissue which I shall notice, is that situated between the valves of Conchiferous Mollusca, in which it performs the office of opening the valves whenever the adductor muscle ceases to contract. In some shells, as the Oyster, this elastic substance is placed within the hinge, in others, as the Cockle, it is external to it; in the former, the expansive power of the compressed substance separates the valves, but in the latter, the same object is gained by the contraction of the hinge ligament. The structure of this form of

tissue has been fully described by my late brother, in the 1st vol. of the "Transactions of the Microscopical Society."

AREOLAR TISSUE.


We now pass on to a mixture of the white and yellow fibres forming a tissue most extensively diffused through the animal eeonomy under the various names of eellular, fibro-eellular, and areolar tissue; the latter being the term now usually employed. The older anatomists imagined, that by the erossing and intertexture of the fibres, a series of more or less round or oval eells were formed; these spaces, however, have no distinct eell-walls, but are merely areolæ or meshes, the eontinuity of which is proved in eases of anasarea and emphysema, and by the inflation of this tissue with air by means of a blow-pipe.

The principal use of areolar tissue is that of connecting other tissues, and at the same time allowing greater or less freedom of motion between them. It forms, with few exceptions, the principal means of support to blood-vessels and nerves, accompanying both to their minutest subdivisions. It is largely developed in certain situations, as for instance, under the skin, where it has received the name of subcutaneous arcolar tissue; under mucous and scrous membranes where it is sometimes very abundant, and is then styled submucous or sub-

serous areolar tissue. In the mesentery of small animals, the Rabbit for instance, the subserous layer is represented only by a few branched fibres, these form a delicate framework over which the serous membrane with its epithelium is stretched. In all the situations, where it is normally found, it may be developed to such an extent as to form tumours, which when subcutaneous, often attain an enormous size. Most of those remarkable growths known as fibrous tumours, are composed of this tissue. They often occur in the neighbourhood of glandular organs, but specimens which have been described as fibrous tumours of the breast, are found on minute examination to be chiefly composed of hypertrophied glandular tissue, intermixed with only so much fibrous tissue as is sufficient to give support to the enlarged lobules of the gland.

When areolar tissue is examined with a power of two hundred and fifty diameters, it will be found, as

FIG. 104.




Areolar tissue of the *Elephant*: $a \ a \ a$, white fibres; $b \ b \ b$, yellow elastic fibres.

principally of white fibres, the yellow or elastic element being only occasionally seen. In the areolar tissue from the pleura of the *Elephant* (Fig. 104) a few yellow fibres (bb) are visible, but when acetic acid is added to a portion of the same tissue, a remarkable change takes

place,—the greater part of the white fibres disap-

FIG. 105.



a a b, fibre forming cells of areolar tissue; c c, stellate cells from a tumour in connection with the parotid gland; d, yellow elastic fibres seen in areolar tissue after the action of acetic acid; e, areolar tissue from the vessels at the base of the brain of a Sheep.

pear, and nothing but long imperfect oval cells or nuclei remain; these, however, as shown by d in Fig. 105, indicate the direction previously occupied by the fibres. The yellow element undergoes no change when treated with the acid, and its fibres therefore stand boldly out, being as it were isolated from the surrounding white fibres, and exhibiting all the peculiar characters I have already described as belonging to this form of tissue. The vellow elastic tissue agrees in all points except in greater minuteness with that of the ligamentum nuchæ of the

Sheep. When areolar tissue is growing rapidly, as in tumours, the cells from which it is developed are often seen; some of the principal varieties are represented by a b, in Fig. 105, they are mostly of a fusiform figure, and what was once the cell-wall, in process of growth becomes a mass of fibres. In tumours connected with the parotid gland, I have more than once


noticed that eells of the shape represented by $c\ c$, are very abundant.

A peeuliar form of areolar tissue is found surrounding the vessels at the base of the brain of the human subject and some of the lower animals; when acted on by acetic acid, the white fibres, which are arranged in separate bundles, assume a tubular form, and the yellow element is seen surrounding them, either in the form of rings or spiral bands, a circumstance which has led some anatomists to suppose that these are blood-vessels in an early stage of formation. A specimen of this peculiar arrangement from the vessels at the base of the brain of a *Sheep* is represented by e in Fig. 105.

I now proceed to describe the mode in which simple membrane and fibrous tissues are supplied with blood-vessels, and the first example I shall take is that of the posterior layer of the capsule of the lens, which has already been described as a structureless membrane, nevertheless, it gives support to large blood-vessels derived from the arteria centralis retinæ, which pass forward over the capsule as far as the iris, where they divide into two branches. These vessels can be readily injected in any young animal, but in the adult I have never yet succeeded in filling them, except in the case of certain reptiles, as for example, the Frog, Toad, Newt, Common English Snake, and Tiger Boa. In none of these animals do the vessels pass farther forwards than what may be termed the equator of the lens; they there

form a border composed of the largest vessels of the net-work, which are probably veins, joining those of the iris. Whilst on this subject, I will point out what I consider the true structure of the membrana pupillaris, which, as you are well aware, is generally looked upon as a distinct membrane closing the pupillary aperture, and when injected, differs from every other known membrane in the peculiarity of the arrangement of its blood-vessels. In three specimens of injected capsules of the lens in the Histological Series, the following peculiarities may be observed. In the first, the entire capsule is seen by the naked eye to be covered with

FIG. 106.




Vessels of the capsule of the lens of a *Puppy:* a. anterior layer of the capsule; b, posterior layer.

vessels, those on the posterior surface represented by b, in Fig. 106, being smaller than those on the anterior α ; but when carefully examined with the microscope, the two sets will be found to communicate; those on the anterior surface a, are the so called vessels of the membrana pupillaris, and may readily be known by their forming a series of loops near the centre, but leaving that spot without any vessels. It therefore appears that at one stage of development of

the lens, the whole eapsule is covered with vessels, and if it should so happen in the eourse of the dissection that the anterior layer be detached from the posterior, this anterior layer would be described as the membrana pupillaris; but if the lens come away entirely covered with vessels, no such membrane is found. The vessels from the posterior eapsule, as before stated, when they reach the iris, divide into two sets of branches, one of which joins the vessels of the iris, the other those of These points are readily seen the anterior capsule. in speeimens from the eye of the Kitten, Wolf, and Puppy, as well as in many other animals, which it is unnecessary here to mention.

We now proceed to describe the vessels of white fibrous tissue. The specimen represented by a, in Fig. 107, is a portion of the tendon of an Ostrich; the

FIG. 107.



a, vessels of the tendon of an Ostrich; bc, vessels of muscle and tendon of rectus externus of the Ostrich.

arrangement of its capillaries is very like that of musele, the vessels are straight, and eonneeted by branches more or less oblique; each vessel runs in the areolar tissue between the bundles, but never amongst the white fibres themselves. The number of the vessels in a tendon is very small as compared with those of the muscle to which it is attached, and in all injected specimens as in the

Diaphragm, or in the Rectus externus muscle from the eye of an Ostrich, Fig. 107, the precise point where the muscle b ends and the tendon c begins, is readily perceptible even to the naked eye, not only by the diminished vascularity of the latter, but also by the difference in the arrangement of its vessels.

The vessels of the yellow fibrous tissue are few in number, and their arrangement, as shown by D, in Fig. 103, is somewhat similar to those of tendon; the eonneeting branches, however, are not transverse, but pass off at angles of about forty degrees, so that the spaces enclosed by the vessels have a somewhat diamond-shaped or rhomboidal outline.

The vessels of areolar tissue, on the contrary, are of small size and very numerous; they follow the direction of the principal fibres, and, as a general rule, form a coarse hexagonal network, which is filled up by capillaries or single vessels. This arrangement is repre-

FIG. 108.



Vessels of areolar tissue from the neck of a young Pig.

This arrangement is represented by Fig. 108. As areolar tissue supports both bloodvessels and nerves, it often happens that the bloodvessels of the sheath of the former, or the neurilemma of the latter may be distinguished from those of the areolar tissue itself. In a specimen from the *Pig*, the vessels of

the neurilemma are plainly seen proceeding in a straight

line, and their arrangement is so characteristic to a practised eye, that the presence of a nerve may be asserted with eonfidence, notwithstanding all the nervous matter has been destroyed in the process of drying.

Although areolar tissue is the ordinary nidus in which the eapillaries as well as the vascular trunks run in the various organs and tissues of the body, still there are certain situations in which the eapillaries are wholly destitute of this means of support; to these, the term naked has been applied by Professor Bowman. He first detected naked eapillaries in the Malpighian bodies of the kidneys; as the capillaries of the brain have no investment of areolar tissue, these also may be termed naked. I have discovered such capillaries in the interior of the glands of Peyer; in a specimen from the Dog, the glands have been divided transversely, and capillaries may be noticed crossing and forming loops in the interior of the gland, but with no investment or support of areolar tissue; the secreting eells usually found in these glands, in all probability, perform this office.

LECTURE XII.

CARTILAGE.

Having described the principal fibrous structures, I pass on to consider the cellular tissues; of these, eartilage first claims our attention. This tissue possessing great clasticity and flexibility, enters largely into the formation of the vertebrate skeleton. It is usually white, or of a light grey colour, changing to a yellowish tinge after long immersion in spirit. Some animals, the eartilaginous fishes, for example, have a skeleton composed almost entirely of cartilage. In man, and the higher vertebrata, this tissue is employed more sparingly in the construction of the adult skeleton; but in the fœtal state the form of the skeleton is sketched out in cartilage, which in the process of development is entirely converted into bone.

Cartilage is usually divided into two kinds, the per-

manent and the temporary, The first, or permanent variety is employed, as before stated, to supply the place of bone, in the skeleton of eartilaginous fishes; or in the form of a membrane, either as tubes enclosing eavities, when it is ealled membraniform, or as a coating to the ends of bones, entering into the formation of joints, when it is termed articular. The second, or temporary form, includes the eartilage of the young of vertebrate animals, as well as all other kinds, which, in process of growth, are converted into bone. All these varieties of eartilage, except the articular, are covered with a strong layer of fibrous tissue termed perichondrium, which being analogous to the periosteum of bone, serves as a support to the blood-vessels and nerves.

The simplest form of cartilage, when examined microscopically, is found to resemble the cellular tissue of vegetables. It consists of a series of cells of a spherical or hexagonal shape, capable, in some cases, of being separated from each other, each of which as previously shown in Fig. 98, possesses a nucleus. In this condition, it forms the *Chorda dorsalis* or rudimentary spinal column of the *Lamprey*, and of the tadpoles of the *Toad* and *Frog*.

In the form of a firm membrane, it is employed in the construction of the framework of the ears of small mammalian animals, such as the Bat, Mouse, and Rat, In the Bat, especially the long-eared English variety, $Plecotus\ auritus$, the cartilage consists of a series of hexagonal cells, on an average $\frac{1}{800}$ th of an inch in

diameter. At the edges of the ear, the cells arc in a single layer, but in the thicker parts two or more layers are superimposed. In these situations the cartilage is precisely similar to certain forms of vegetable cellular tissue.

The ear of the common *Mouse* may be taken as a good specimen of simple cartilage; the central portion consists of a series of hexagonal cells arranged in layers one over the other, so that in every other respect, except in the size of the cells, the structure resembles that of a transverse section of the pith of a plant; this, however, has been before alluded to, and represented in Fig. 99.

In the thin cartilage of the ear of the *Bat* above named, the cartilage cells are of smaller size, and have thicker walls than those of the *Mouse*; in all parts, especially on the edges, the cartilage is sufficiently transparent to be seen through; and, in nearly every cell, as central nucleus may be observed. This, like the pre-

FIG. 109.




Transverse section of the *Chorda* dorsalis of a *Lamprey*.

ceding, has been before described, and is represented in Fig. 100.

The transverse section of the *Chorda dorsalis* of a *Lamprey*, a portion of which is represented in Fig. 109, consists entirely of large cells of a more or less hexagonal figure, those near the margins being arranged in the form of radii. Some parts of this Chorda dorsalis, especially those near the centre,

FIG. 110.



Portion of the vertebral column of a *Trout* three weeks old: A, part of a spine composed of transitional cartilage. B, *Chorda dorsalis*.

are soft and pulpy; in these, the eells can be separated from each other, but, nearer the circumference, the eells are more compressed and very firmly adherent.


These large eells are not confined to the Chorda dorsalis of cartilaginous fishes, they exist also in the embryos of osseous fishes, and in the Tadpoles of batrachians. A portion of the vertebral column of a Trout three weeks old is represented in Fig. 110; the larger eells seen at B constitute the true Chorda dorsalis, which never becomes ossified; but above them, as

at A, a rudimentary neural spine is seen, this is composed of eells of much smaller size, and characteristic of eartilage that is afterwards converted into bone. At this early stage, then, we can distinguish between permanent and transitional cartilage.

In the examples I have hitherto given, no other substance than the cells has been alluded to; the majority of eartilages, however, will be found to consist of a matrix

in which the eells are imbedded. The matrix may either be perfectly homogeneous in structure, or granular, and it is not unfrequently fibrous; to this substance I would

FIG. 111.



A, cartilage of the Cuttle-fish, Sepia officinalis. B, portion of an Enchondroma.

give the name of intereellular substance or matrix. The lowest animal in which I have been able to meet with eartilage is the *Cuttle-fish*. In this cephalopod it occurs in the form of a rudimentary skeleton; the cells (Fig. 111, A) are small, more or less irregular in figure, like the cells of bone; and are imbedded in a

structureless intercellular substance. Around some of these eells there is a faint trace of cell-wall, and it becomes a question whether many of them should not be considered as altered nuclei.

Cartilage is sometimes found in large masses, as a product of disease, in the form of tumours, termed *Enchondroma* by Müller. These tumours may be formed in connexion with any of the bones, but are more frequently attached to the phalanges, and the bones of the extremities. They are made up of cells like those of cartilage, but occasionally they also contain cells of a very peculiar form, Fig. 111, B, bearing a striking resemblance to the lacunæ of bones, but more especially to the cells found in the cartilage of the *Cuttle-fish*, above mentioned; to these tumours I shall

have oceasion to revert in a future Lecture In eartilaginous fishes generally, all parts of the skeleton, except the *Chorda dorsalis*, are composed of a firm hyaline intercellular substance, in which, numerous oval nucleated cells are imbedded, they are generally arranged in groups of five or six; distinctly nucleated cells exist in the cartilage of the head of a *Ray*, which also occur in fasciculi or groups imbedded in a slightly granular matrix. In another section of the cartilage from the same *Ray*, the granular appearance is more evident; this portion of the cartilage, which will be alluded to hereafter, is undergoing the process of ossification, and the granules are those of osseous matter.

In Reptiles, the eartilage-eells are larger than in Fishes, and in the Siren they attain their greatest size; they are smallest in the Crocodilia. In Birds there is seareely any eartilage, except that entering into the formation of the joints; in these animals all the eartilage is converted into bone, at a very early period of life.

In the Mammalia we have large eells, and a firm structureless intercellular substance; the largest eells, according to my own observations, being those in the Elephant. In some eases, the intercellular substance is of a fibrous character; if thin sections be taken from the eartilage of the ear of the Rabbit or Dog, the eells can be detached from the meshes formed by the fibres. In a vertical section of the cartilage of the auricle of the human ear, the cells are in some parts

uniform in size, and closely packed; in others, the intercellular substance is fibrous, and oceasionally some of the cells, as I have just mentioned when speaking of the ear of the *Rabbit*, can be separated from the fibrous net-work surrounding them. The reverse is the case in sections of the costal cartilage of the human subject, in which the cells are few, and widely scattered, whilst the matrix is most abundant; the cells are of large size, and frequently arranged in rows, their nuclei are clear and transparent, and the matrix minutely granular.

I now proceed to describe the mode in which this permanent form of cartilage is supplied with blood. Cartilage, as I have already stated, is invested with a fine glistening membrane ealled *perichondrium*, which supports the vessels and is prolonged with them into the interior of all the thicker cartilages; the distribution of the vessels is very like that in arcolar tissue.

In an injected specimen of eartilage from the ear of the *Rabbit*, the vessels are of large size, and each artery is accompanied by two veins, many of which are perceptible to the naked eye, but the vessels are by no means numerous.

On the cartilage of the auricle of the human ear, I have observed that the vessels of the outer surface are more numerous and larger than on the inner surface. On the former, the vessels are exceedingly large and tortuous, (Fig. 112), and their arrangement is manifestly different from those of areolar tissue. On

FIG. 112.



Vessels of auricle of the human Ear.

the latter, the vessels have the same arrangement, but it may be readily seen that they are smaller and much less numerous. This description applies only to the most vascular portions of the cartilage, for in those parts which have a less amount of vascularity, the arrangement

of the vessels is precisely that of areolar tissue.


LECTURE XIII.

ARTICULAR CARTILAGE.

Having now described to you the principal forms of membraniform and permanent cartilage. I pass on to notice that variety of permanent cartilage, which, from its entering into the formation of joints, is called Articular. It differs in structure in the young and in the adult animal, and it is this form of cartilage which, in the young condition is largely supplied with blood-vessels, and undergoes a constant and successive process of ossification, on its attached surface. If a vertical section of fœtal cartilage be examined, the part nearest the articular surface (Fig. 113, A) exhibits numerous small cartilage-cells, arranged without much order, and the nucleus of each cell occupying its whole diameter. As we proceed towards the attached surface, the cartilage-cells begin to be arranged in parallel rows,

and the distance between the nucleus and cell-wall becomes much greater; and last of all, close to the attached surface B, we find spicula or lines, projecting between the parallel rows of cells; these are sections

FIG. 113.



Vertical section of articular cartilage from the head of the humerus of a feetal Wolf: A, articular surface. B, attached surface.

FIG. 114.



Vertical section of articular cartilage from a human metacarpal bone: A, articular surface. B, bone of the shaft. c, articular lamella.

of the walls of tubes of newly-formed bone, and which, when fully developed, become solid, each having enclosed a column of large cartilage-cells. In many sections of articular cartilage, as in Fig. 113, A, taken from the head of the humerus of a young Wolf, there are foramina through which blood-vessels pass to nourish

it, these are indicated by the four oval spots or holes, two of which are shown as being filled with injection.

In adult Mammalian articular cartilage, represented by Fig. 114, the arrangement of the cells is very peculiar; the cartilage is separated from the bone by a white layer (c) of variable thickness, and which contains bone-cells three or four times as large as those of ordinary bone. This is the non-vascular lamella first described by Mr. Toynbee, the lower portion is connected with the osseous tissue of the shaft of the bone (B), while the upper is more or less tuberculated for the attachment of the cartilage. The cells in the lowest layer of cartilage, or those immediately above the lamella, are arranged in columns parallel to the axis of the shaft of the bone, but as we proceed to the articular surface, the columns become smaller, more numerous, and their direction is changed, so that they are now at right angles to the axis of the shaft. Immediately on the articular edge of the section (A), the cells are so much flattened as to present the appearance of epithelium cells, and the intercellular substance or matrix of the cartilage, assumes a fibrous appearance. Adult articular cartilage is not vascular; the capillaries of the osseous tissue of the shaft of the bone terminating in loops as they approach the non-vascular lamella. From this brief outline of the structure of articular cartilage in the young and adult condition, I proceed to notice some peculiarities in that of fishes, reptiles, and birds.

In fishes, as in the vertical section of the jaw of a Conger eel, the eartilage-cells are small in size, very few in number, and arranged without much order in a distinctly fibrous matrix, the fibres of which follow the direction of the shaft of the bone. Upon the articular surface, however, the eartilage-eells are minute, and arranged in parallel rows; but they are so abundant as to obscure all trace of fibres.

In reptiles, as in the *Tortoise*, the articular or free surface of the cartilage has a distinctly fibro-eartilaginous structure; the fibres interlace and produce meshes, in which the cartilage-eells are lodged. Near the bone the fibres disappear, and the eells are of an oval shape, and few in number. The most remarkable circumstance in connection with this specimen is the great thickness of the fibro-eartilage of the articular surface.

In the Batrachia, as may be observed in a section of the head of the femur of a Frog, the eartilaginous matrix is clear and transparent; the cartilage-cells are smallest and most numerous in the part nearest the bone, but as we proceed towards the articular surface they become larger, and the nuclei are more plainly seen. The most interesting fact connected with the cartilage of the Batrachia is, the large size and uniformity of arrangement of the cells upon the articular surface; they are perfectly flat, resemble scales of tessellated cpithelium, and many of them exhibit an appearance of one large cell dividing into four by cruciform fissures at right angles to cach other.

In birds, the articular cartilage is always more or less fibro-cartilaginous. In the *Turkey*, the fibres proceed in straight lines in the direction of the shaft, the cartilage cells being arranged in rows between the fasciculi of fibres. The articular surface of the specimen Fig. 115, exhibits fibres running in various directions among the cartilage-cells; this arrangement is constant in birds, and is precisely similar to that occurring in fishes. It is a remarkable fact, that in birds there is little or no cartilage except that of the

FIG. 115.



Articular surface of cartilage of the femur of a *Turkey*.

FIG. 116.



Vertical section of articular cartilage of the femur of a *Turkey*.


articular surfaces of the joints, and occasionally the rings of the trachea; all the parts of the skeleton of these animals are rapidly converted into bone, and even in the joints, the coating tissue, Fig. 116, often of considerable thickness, is composed of fibro-cartilage.

The moment, however, we examine sections of articular cartilages in the Mammalia, we find a constant arrangement of the cells, and in the adult

the non-vascular lamella makes its appearance. In the part of the eartilage nearest the articular lamella, as secn in a section of the articular surface of the bonc of an Ox, the cells are arranged in columns, but near the articular surface their direction is changed. The same thing is evident in vertical sections of the articular cartilage of the humerus of a Piq, in which also, a welldeveloped non-vaseular lamella is visible, and as in the Ox the cartilage-cells near the articular surface are arranged at right angles to those nearest the bone. In a vertical section of the head of a femur of a female, aged ninetcen, the same arrangement is perceptible; but it will be noticed, that the articular lamella is sparingly developed. In a vertical section of the corresponding bone of a female upwards of seventy years of age, in which the arrangement of the cells in the cartilage precisely resembles that in the preceding case, the non-vascular lamella is nearly twice as broad. If the free surface of the articular cartilage of the young woman be examined, the eells will be seen to be more or less flattened, and to contain nuclei; in the older woman the cells are present, but the nuclei have disappeared. Having demonstrated the principal varieties of articular cartilage, it now becomes necessary that I should speak of the synovial membrane, which is said by some persons to invest the whole articular surface. In the feetal condition of man and of the higher mammalia each joint is eovered with synovial membrane and an epithelium; as soon, however, as the joint is used, the

epithelium, and subsequently the synovial membrane itself disappears from all parts subjected to friction, although, as Mr. Toynbee remarks, it is sometimes persistent on the patella. In all my observations on the synovial membrane, and in the sections made for description in the Histological Catalogue, which include specimens taken from a large number of animals of each of the four great classes, I have never found it very far in advance of the vessels.





Vessels of synovial membrane and articular cartilage: A, glenoid eavity of human fœtus. B, looped vessels of the synovial membrane. C, vertical section of radius of human fœtus. D, vessels of fœtal articular cartilage running in canals.

Articular cartilage in the feetal condition is supplied with blood-vessels, which pass into its substance, and all parts in the neighbourhood of its articular surface are nourished by the vessels of the synovial membrane, which, as will be presently shown, always terminate in looped extremities immediately on the margin of the surface subjected to pressure. The glenoid cavity of a human feetus, Fig. 117, A, is surrounded by a margin of capillaries which project inwards as far as the socket for the reception of the head of the humerus, where each terminates in a looped extremity, the loop itself, Fig. 117, B, being sometimes dilated to two or three times the size of that of the vessel from which it is derived, it would seem that the synovial fluid is poured out from these dilated vascular loops. In a vertical section of the head of a metacarpal bone of an adult, the looped vessels of the synovial membrane are distinctly seen passing upon the articular cartilage as far as the part subjected to friction; these vessels do not enter the cartilage, neither is there any other source from which adult cartilage can derive its nutrition, except these vessels of the synovial membrane.

LECTURE XIV.

ARTICULAR CARTILAGE.—SYNOVIAL MEMBRANE.

Having demonstrated that the vessels of synovial membrane, both in the fœtus and in the adult, pass upon the surface of articular eartilage as far as the part subjected to friction, I will now endeavour to point out in what way the interior of the cartilage is supplied with vessels, and I shall in the first place describe a vertical section of the upper half of the right radius of a fœtus, Fig. 117, c, in which, vessels from the periosteum may be seen passing into the cartilage immediately above its connexion with the shaft, these divide and subdivide into numerous branches. The shaft of the bone also is injected, but its vessels are not continued into the cartilage, the layer of bone upon which the cartilage is situated, termed by Mr. Toynbee the non-

vaseular lamella, and indicated by the dotted line, not admitting of their passage through it. The vessels of the shaft on reaching the lamella are said to terminate in loops. In the opposite half of the same specimen, which is represented in the Histological Catalogue, Plate VIII, Fig. 4, it may be seen, even with the naked eye, that the shaft in the immediate vicinity of the eartilage has a greater number of vessels than any other part, and it is at this spot that the growth of new bone is taking place. Immediately above these vessels the eartilaginous epiphysis is seen, and upon all parts of its surface, except those subjected to friction, the eapillaries of the synovial membrane are distributed, all of them terminating in loops. In a vertical section of the head

FIG. 118.



Vertical section of the head of a tibia of a human fœtus.

of a fœtal tibia with its epiphysis attached, a rich network of vessels belonging to the synovial membrane ramifies upon its upper margin; there are numerous vessels also in the epiphysis, some of which enter the eartilage near the upper margin, but others, as shown in Fig. 118, may

be seen winding round between the upper part of the shaft and the epiphysis; these last are derived from the articular vessels of the periosteum.

In the shaft, Haversian eanals may be observed, some of these have vessels in them. The portion of bone

immediately beneath the epiphysis is whiter than the rest; this is the non-vascular lamella before alluded to, and no vessels pass through it to supply the cartilage.

All the vessels of fœtal cartilage run in eanals. In the early stage of embryonic existence, as Mr. Toynbee has demonstrated, the cartilage has deep notehes or indentations for their reception, and at a later period of growth the canals are still visible in the interior. In a section from the head of the humerus of a human fœtus, the vessels, Fig. 117, D, are very numerous and irregular in their outline. If any one of them be accurately focussed, it will be seen enclosed in a tube or eanal. The vessels themselves are small in comparison with the canal in which they run, but other vessels may be noticed in the neighbourhood, which are so distended as to fill the entire canal.


In healthy adult articular cartilage, there are no vessels; but in morbid conditions of the same tissue, vessels oceasionally become more or less numerous. The latest period of life at which I have discovered vessels in healthy eartilage is twelve years.

In the specimen before represented in Fig. 114, A, which is a vertical section of the head of a metacarpal bone of an adult, the vessels of the shaft of the bone could be traced until they reached the yellowish band, or the non-vascular, or articular amella, and here they ended in loops. The eartilage in this specimen is distinguished from the lamella by being whiter and much more transparent. The lamella is even present

in the fœtus, it is indicated by the dotted line in Fig. 117, c; the vessels of the eartilage, which in this instance are very numerous, being derived from those of the articulation, and not from those of the shaft.

The non-vascular character of healthy articular cartilage, is maintained from the period of youth to old age; but, if disease occur, vessels may soon be found in it. In a specimen from a diseased joint, which after removal was carefully injected, numerous vessels may be observed passing through the eartilage; they are derived from the vessels of the shaft, as the articular lamella being involved in the disease, permits the vessels to pass through it; they proceed in straight lines through the eartilage to the free articular surface upon which they form a network, and anastomose with others probably derived from the synovial membrane. The subject from whom this specimen of eartilage was obtained was fifty years of age, and the disease had existed for nearly twelve months. A preparation which belonged

FIG. 119.




Section of diseased articular cartilage largely supplied with vessels.

to the late Mr. Liston, and which he was in the habit of exhibiting in his Leetures, consists of the head of the tibia, with diseased eartilage attached. Not only can vessels be seen by the naked eye, passing from the bony shaft into the cartilage, in the form of loops, as in Fig. 119, but a

rich net-work may in some cases be observed upon a large portion of the articular surface. As far as I have been able to learn from examinations of diseased articular cartilages, especially those affected with ulceration, I conclude that the change first takes place in the cartilage cells, as is made evident by their becoming rounder and much larger in size, and by their contents assuming a different character, the nuclei disappearing, and globules of oil taking their place. In some cases these oil-globules are of very minute size, and the cells then appear granular; as the disease goes on, the cell-walls are absorbed, a series of cavities are formed, all the hyaline substance in the neighbourhood becomes more or less fibrous, and ultimately blood-vessels are developed in the fibrous tissue. The diseases of cartilage have been studied by very many able observers, especially Sir Benjamin Brodie, the late Mr. Key, Mr. Liston, and more recently by Dr. Redfern, of Aberdeen, whose papers will be found in the "Monthly Journal of Medical Science," for 1849.

The nutriment of articular cartilage is derived from two sources, namely, the articular vessels and those of the synovial membrane, more especially the latter; but neither set ever enters its substance, the one set ending in loops immediately without the parts subjected to friction in the movement of the joint, the other set confined to the cancelli of the shaft of the bone, and separated from the cartilage by the non-vascular lamella. The vessels of the synovial membrane of feetal articular

eartilages have been already noticed. In the adult condition, as shown in a portion of the head of a metacarpal bone, Fig. 120, a, the vessels are equally numerous, and




Vessels of synovial membrane: a, portion of the head of a metacarpal bonc of an adult with the synovial vessels injected; b, vessels from a part of the same magnified 20 diameters.

when magnified at least twenty diameters, their looped terminations b are well displayed; they not only are eontinued upon the eartilage as far as the parts eoneerned in loeomotion, but in some eases processes of the membrane, like fringes or large villi, riehly supplied with vessels, project into every nook and eorner where they

are not liable to injury. These processes, which in the knee-joint were described by Clopton Havers, and stated by him to possess a glandular office, have upon more recent investigation been presumed to be entitled to such a character as Havers ascribed to them, although most anatomists since his time have regarded them as masses of fat. If one of the joints of a finger be laid open, the vessels of the synovial membrane forming the capsule of the joint will be found thrown into a series of processes like villi, which are largely supplied with vessels remarkable for their tortuosity. These villi project into all parts of the cavity of the joint, and the vessels are supposed to pour out the synovia; they may, therefore, be viewed as synovial glands. The

arrangement of the eapillaries of some of the largest specimens is represented in Fig. 122, a; similar processes, which no doubt secrete a fluid somewhat like synovia, are found within the sheaths of tendons, and upon those tendons which perforate or are perforated by others, as in the ease of the flexor tendons of the fingers. Every part of the surface, both of the tendon and of the sheath in which it is contained, has a rich capillary network, except in such parts as are subjected to friction. The vessels of the synovial sheath (Fig. 121) taken from the middle finger of an adult human subject

FIG. 121.




Tortuous vessels from sheath of flexor tendon of the middle finger.

are remarkable for the manner in which they are convoluted into a somewhat heliaeal form, but the helices do not project far beyond the general surface of the synovial membrane. In another specimen, of larger size than the preceding, and which has been dried after injec-

tion, the eapillary net-work and the heliees are beautifully shown, and still nearer the joint the vessels are more numerous. If a portion of this vascular membrane be removed and earefully examined with a power of two hundred and fifty diameters, the eapillaries will be generally found to terminate in loops, but the synovial membrane itself extends some little distance beyond the vessels, in a few eases as far as $\frac{1}{25}$ th of an inch; this non-

vascular portion being covered with epithelium. One of the villi, with its looped vessels, is represented in Fig. 122, c, and a few of the loops more highly magnified in Fig. 122, e; the scales of epithelium upon the basement membrane extending beyond the vessels are shown at d.





Vessels of synovial membrane: a, villi from joint of middle finger; b, looped vessels from flexor tendon of leg of an Ostrich; c, villus from head of metaearpal bone; d, epithelium of synovial membrane in advance of the vessels; e, loops of synovial vessels magnified 500 diameters.

Whilst on the subject of synovial vessels, I will take the opportunity of demonstrating those accompanying the ligamentum teres from its origin to its insertion; and it may not be out of place here to mention the fact, that some few animals, as the

Elephant, have no ligamentum teres; in the Frog there is no ligamentum teres in the hips, but onc in the shoulder-joint; the use of the ligament in this situation being explicable by the habits of the Frog, that of hopping and pitching principally on its fore-feet, whereby a considerable tendency to dislocation must necessarily occur in the shoulder-joint. If the ligament be examined in a well injected fœtus, a large supply of blood-vessels will be found in the synovial membrane of the acctabulum which proceed in straight lines along the ligament, and terminate in loops close to its insertion into the head of the femur. A preparation illustrative of this fact is accurately represented in Plate VIII., Fig. 9, of the "Histological Catalogue," but the looped termination of the capillaries is best seen in Fig. 8, this being a more highly magnified view of that part of the femur into which the ligament is inscrted.

The vessels of synovial membrane, and those of the sheaths of tendons, have been well described by Mr. Toynbee and Mr. Rainey, especially by the former gentleman, who has figured those of the ligamentum teres in a paper published in the "Philosophieal Transactions" for 1841. The beautiful looped vessels are not eonfined to the human subject, but exist in most of the lower animals. In the long flexor tendons of the leg of an Ostrich, eireular patches of eapillaries may be seen, each of which, as shown in Fig. 122, b, terminates in a dilated loop. The vessels of tendon which were exhibited when describing the vascularity of fibrous structures, were

those of the Ostrich, and it was then mentioned that the rich superficial network belonged to the sheath of the tendon, and not to the tendinous fibres themselves.

Whilst treating of articular eartilage, I may mention a peculiar deposit, as it is termed, which takes place in many joints that have been deprived of their eartilage by disease. The parts subjected to friction, in the movements of the joint, present a highly polished appearance, which is known as the ivory-like or porcellaneous deposit. If these specimens be examined, it will be found that all parts of the bone in the neighbourhood of the joint have an additional quantity of bony matter thrown out probably as the result of rheumatic inflammation. The same thing, no doubt, would have taken place upon the polished surfaces, had not the exuberant growth been kept down by the frietion; and, as no bone oeeupying this situation, unless of preternatural hardness, could receive such a polish, I was led to speculate on the eause of the appearance. I therefore removed sliees, and rendered them sufficiently thin for microscopie observation, by grinding away the eut surface; and, having examined them, I found that there was an almost total absence of the Haversian canals, whereby the bone was rendered exceedingly dense, and I concluded that the new osseous matter, having been prevented by frietion from being thrown out upon the surface, was employed in filling up the canals, and by this means a substance, originally porous,

was eonverted into a solid mass, susceptible of taking the highest polish. This view of the subject eorresponds precisely with a practice adopted by French polishers, who are oeeasionally obliged to fill up the pores, or eanals, in many of the hard woods, such as rosewood and mahogany, before they ean bring the polish to any perfection. I must now allude to a point which, for many years, has been the subject of dispute among anatomists, namely, whether the synovial membrane is continued over the surface of articular eartilage. If a joint from a very young animal be examined before it has been used, the surface of the eartilage will be found eovered with synovial membrane, having an epithelium of that variety known as the tessellated; within a short period, however, after the joint has been used, the epithelium disappears, and, subsequently, the synovial membrane itself. This view of the subject is the one generally entertained; but Mr. Toynbee is of opinion that the membrane is not unfrequently present upon the cartilage, although its epithelial eoating may have disappeared. In making the preparations described in the Histological Catalogue, many hundred sections were taken from the eartilages of the four great elasses of animals; all were examined with reference to this point, and I have satisfied myself that the membrane is rarely, if ever, continued over the cartilages of the long bones of the adult. In the patella, and other sesamoid bones, the synovial membrane, as shown by Mr. Toynbee, may oceasionally be

stripped off, but not from the surfaces of the more perfect joints. All my examinations tend to prove that the synovial membrane is continued a short distance beyond the vascular network, and that it ends at the point where friction commences. The cells on the surface of the cartilage, which have so frequently been mistaken for those of epithelium, are nothing more than ordinary cartilage-cells very much flattened; the resemblance of these cells to epithelium is most strikingly shown, as I have already stated, in the *Batrachian* reptiles.

LECTURE XV.

CARTILAGE. — ENCHONDROMA.

We now arrive at the examination of a structure presenting all the characters of cartilage, which occurs in the form of tumours, and named by Müller Enchondroma. Tumours of this character may occur in many situations, and sometimes attain a considerable size. One removed from the neck of a man, by John Hunter, weighing one hundred and forty-four ounces, and consisting of nodular masses of cartilaginous texture, bound together by areolar tissue, is preserved in the Museum of the College. Another specimen, in the same collection, is that of a hand amputated by Sir Astley Cooper, which has enchondromatous tumours attached to almost every bone. A tumour of this kind removed from the cheek by Sir W. Blizard, forming Prep. 202, in the Pathological Series, contains numerous spicula of bone.


The lcg of a young woman having a tumour also of this character, twenty inches in circumference, was lately removed by Mr. Lloyd at St. Bartholomew's Hospital. If these tumours are carefully examined, they will be found to be composed of a tissue resembling either cartilage or fibro-cartilage. The tumour, removed by Hunter, already alluded to, consists of a series of large oval or spherical nucleated cells, imbedded in a firm intercellular substance, which is in some parts structureless, in others more or less fibrous; the cells, Fig. 123, contain nuclei which are very granular, and of a brown colour.

FIG. 123.



Section of an *Enchondroma* from the neck.

FIG. 124.




Section of an *Enchondroma* from one of the ribs.

Fig. 124 represents the structure of an enchondromatous tumour, removed after death from the rib of a man; it more nearly resembles fibro-cartilage than true cartilage, many cells occurring among a dense mass of fibres. In an *Enchondroma* from the cheek, as shown by B, in Fig. 125, some of the cells have

nuclei of large size, others, as at A, are dark and granular, the latter being in the first stage of ossification. In the specimen from which this preparation was taken, there were many points or spicula of bone, but the fibres were least numerous in these parts.

FIG. 125.

FIG. 126.



Portion of an *Enchondroma* from the cheek: A, cells and matrix becoming ossified. B, cells with nuclei.




Portion of an *Enchondroma* from the hand: A, cells imbedded in a fibrous matrix. B, cells detached from the matrix.

The next specimens I shall describe are of great interest, as they throw considerable light on the formation of the lacunæ of bone. The first, Fig. 126, is a thin section of one of the tumours developed from the phalanges, &c., of a hand, removed some years since by the late Sir Astley Cooper; it is composed of a series of small cells, somewhat resembling the lacunæ of bone, each cell having branching tubes or canaliculi projecting from it in all directions. Some of these cells, as shown at A, are firmly imbedded

in a fibrous matrix; others, in the same tumour, as at B, can be separated from the matrix, proving that they are true cells, and not mere vacuities in the hyaline substance. In order to demonstrate that these cells are, in all probability, altered nuclei, the original cell-walls of which have disappeared, I shall describe the structure of a section of the enchondromatous tumour of the leg, before referred to in page 164, as having been lately removed by Mr. Lloyd at St. Bartholomew's Hospital; it was a large mass, measuring twenty-two inches in circumference, and very thin sections exhibit cells of an oval or circular figure, having a central nucleus and a concentric laminated deposit. Some of these, represented in Fig. 127, b, exhibit a minutely granular structure, as though ossification had commenced in them; other parts of the section exhibit cells of similar size, in which the nucleus is not only well developed, but occasionally small projecting points, like commencing canaliculi, may be observed, as in Fig. 127, a. As soon, however, as the canaliculi become somewhat more evident, the cell-wall disappears, and we then find cells of similar shape to those in the first specimen, but of larger size, Fig. 127, c. It sometimes happens that these cells are met with either above or below a cartilage cell, and it then appears as though the altered nucleus were still within a cell wall; but careful focussing will prove that this view of their relative situation is incorrect. In one part of the specimen, there is a thin

layer of bone, in which the lacunæ are large, and have but few canaliculi; these, as shown in Fig. 127, d, very much resemble the nuclei of the cells, and, in all probability, were formed from them.





Cells and bone from an Enchondroma of the tibia: a, cells each having a stellate nucleus; b, cells showing laminated deposit and ossific granules; c, stellate nuclei; d, thin portion of bone in connection with the cartilage.

With regard to the formation of the lacunæ of bone, two views are now entertained by different histologists. The first is that given in the "Physiological Anatomy" of Professors Todd and Bowman, in which it is stated that the lacunæ are developed from the nuclei of the cartilage cells; the other that of Mr. Tomes, published in "Todd's Cyclopædia," article "Osscous Tissuc," in which it is asserted that the lacunæ are not developed

from the nuclei of the eartilage cells, but are cavities left in the newly formed bone, from which canaliculi are subsequently developed. The last described specimen of enchondroma, however, tends to prove that the view entertained by Todd and Bowman is the more correct.

In order to understand what I am now about to describe, it will be necessary for me to give a brief account of the structure of bone. If we take, for example, one of the long bones of the body, we shall find that it consists of a shaft and two extremities; if the same bone be divided transversely, its centre will be found to be hollow, or occupied by a spongy kind of bone, which has received the name of cancellated structure. Within the hollow, the marrow is contained, and the cavity, as you are well aware, is called from this circumstance the medullary cavity. If we examine the outer surface, we shall find a series of minute holes or foramina through which blood-vessels pass towards the

FIG. 128.



Portion of a transverse section of a human feature magnified 40 diameters.

interior of the bonc. If now, a sliec sufficiently thin to be transparent, be made and examined with a power of forty diameters, a series of holes will be noticed, around which innumerable black spots are deposited in concentric circles, as shown in Fig. 128, these are the Haversian canals, and through them the

blood-vessels for the nutrition of the bone are transmitted. These canals are larger on the inner or medullary surface of the bone than they are on the

FIG. 129.



Portion of a transverse section of a femur magnified 200 diameters.

outer. Let the same section be next examined under a power of two hundred diameters, a concentric arrangement of laminæ, as represented in Fig. 129, will be seen around cach canal, and the black dots before alluded to will now present a spider-like appearance; they are known as the lacunæ, or bone-cells, and each con-

sists of a central part or body, from which a number of minute tubes termed eanalieuli proceed. The eanaliculi nearest to each Haversian canal open into it, whilst those more distant from the same eanal, anastomose with the canalieuli of the next lamina; those of the outer row of bone-eells do not anastomose with the canalieuli of neighbouring laminæ, but nearly all bend back and join those of the preceding lamina. By this arrangement, a white line may be observed to surround the outer part of each of the concentrie eireles; so that the bone may be said to be built up of a series of Haversian systems. If the same bone be divided in a vertical direction, as shown by Fig. 130, the course of the Haversian eanals may be readily seen; they run in parallel lines, and are connected together by oblique,

FIG. 130.



Vertical section of human femur magnified 40 diameters.

or more or less transverse branches. Within them, the blood-vessels, from which the nutritious matter is poured out, are contained; this being taken up by the canaliculi opening into the canals, is by them conveyed to the whole concentric arrangement of bone-cells; each bone-cell,

therefore, may be considered as a reservoir of nutriment for the bony matter surrounding it.

It may be asked, since all the structures I have described are of a tubular character, where is the osseous matter? To observe this, we must have recourse to much higher powers, and it will then be found that the bone consists of a congeries of more or less minute angular particles deposited in an organized

FIG. 131.




Portion of a transverse section of a femur magnified 500 diameters, showing the granules of osseous matter.

matrix. Fig. 131 is a representation of a portion of the same section of bone as Fig. 129 was copied from, but the power employed has been one of 500 diameters; the bone-cells and their canaliculi are well shown, and the minute dots occurring in all the spaces between the canaliculi are the granules, or

particles of osseous matter. If such a section be placed in dilute acid, these particles alone are removed, and the matrix in which they were imbedded presents a granular appearance. I have found, that in the crania of very small birds, as the *Canary*, where the bone is too thin to admit of bone-cells, the earthy particles are not only of large size, but each, as shown by B, in Fig. 134, is of a rhombohedral form. In the cartilage of the cranium of *Rana paradoxa*, minute needle-shaped crystals somewhat resembling the raphides in the *Squill* and *Hyacinth*, are occasionally found in the interior of each cell, thus proving that ossific matter may be deposited either in a granular or crystalline form.

In the cartilage of many fishes, as the Shark, Skate, and Saw-fish, the ossific matter is in the form of granules, and occurs principally in the neighbourhood of

FIG. 132.



Cartilage from the eranium of a Skate: A, cells imbedded in a structureless matrix. B, granules of ossific matter around the cells.

the cells; in the Skate, the deposit sometimes takes place within the cell wall. To the latter I give the name cellular, and to the former intercellular ossification. In Fig. 132, are shown two portions of the cranial cartilage of a Skate; the one represented by A, exhibits a mass of cells imbedded in a more or less structureless matrix, whilst in

the other, B, the granules of ossifie matter around the

FIG. 133.




Vertical section of ossifying cartilage from the humerus of a feetal *Wolf*.

cells are well shown. A good example of intercellular ossification has already been given in Fig. 127, b. When a cartilaginous epiphysis is undergoing the process of ossification, the cartilage-cells first arrange themselves in linear series, and at the period of ossification, the cellwalls, as in Fig. 133, become more widely separated from the

nucleus, whilst the cells themselves are studded with minute ossific granules; if a section be made at right angles to the shaft, the walls of the eartilage-cells, as

FIG. 134.




A, transverse section of ossifying cartilage from humerus of feetal *Wolf*. B, osseous granules from eranium of a *Canary*.

represented by A, in Fig. 134, will be found thickened by the granular deposit; subsequently, tubes of bone shoot up between the cells, and so enclose them. The part in which this process is going on may be readily distinguished by the arrangement of the cartilage-eells. In a vertical section of one of the short

phalanges from the foot of a young Pig, Fig. 135, a thin layer of cartilage will be seen between two pieces of bone; if this be more closely examined, it

FIG. 135.



Vertical section of one of the phalanges from the fore-foot of a Pig: A, distal extremity ossification going on slowly. B, proximal extremity, ossification proceeding rapidly.

will be found, that at the proximal extremity B, in which the formation of bone is going on, the eartilage-cells are arranged in linear series, but that at the distal one, A, only a trace of such arrangement is visible.


The ultimate osseous granules were first pointed out by Mr. Tomes, they cannot be well seen in very compact bone until it has been boiled, but may be readily distinguished in imperfectly developed bones. In a vertical section of a portion of the os frontis, belonging to the Histological Collection, a

spot of imperfectly formed bone occurs near the centre, in which the granules are very visible. In certain forms of ossific deposit, as in the early stage of ossification of the coats of arteries, nothing but granules can be seen, and the same holds good in ossified fibrous tumours and cysts found occasionally in various parts of the body.

The ossific granules are generally discoverable in the pus which escapes from a necrosed bone. If a specimen of this pus be examined with a power of five hundred diameters, a quantity of minute granular matter, readily soluble in dilute muriatic acid, may be distinguished among the pus corpuscles. This fact I noticed many years ago, but Mr. Bransby Cooper has ascertained by chemical examination, that a large amount of phosphate of lime is present in such pus, and announced his discovery in 1843, when Professor of Surgery to this College; so that it would appear that the pus has a solvent power, eroding or decomposing the animal matter, while the mineral constituent escapes with the pus in its granular condition.

The bone-cells or lacunæ vary in size and shape in the four great classes of animals, and I have ascertained that they bear a direct relative proportion to that of the blood corpuscles.* They are largest in reptiles, especially those of the *Perennibranchiate* order. In fishes their

FIG. 136.



Portion of the scale of a fish, Lepidosteus osseus.

shape is very peculiar; and, what is remarkable, they are in some cases similar in form to the stellate cells found in the specimen of enchondroma from the hand, before described and represented in Fig. 126, as is illustrated by a thin section


^{* &}quot;Transactions of the Microscopical Society," Vol. 11.

of a scale of the Bony Pike, of North America, Lepidosteus osseus, Fig. 136.

In Mollities ossium, there is a deficiency of the earthy constituent of the bone. The change first begins in the lucanæ, which become larger and larger, and the bone around them more and more transparent; finally, several lacunæ unite to form one cavity, which, however, does not long remain empty, but is occupied by a soft kind of adipose tissue, so that such bones are always extremely thin and full of fat. For this reason Mollities ossium may be considered as an example of the fatty degeneration of bone.

The last form of cartilage which I shall mention is of a very peculiar character, being always more or less tubular. It is found in the vertebræ of the Vaagmaer or Riband-fish,—a specimen of which, about three years since, was captured off St. Andrews. For our knowledge of this interesting structure we are indebted to the late Dr. John Reid; and the account of the anatomy of the fish was the last contribution to science made by that distinguished anatomist. A transverse section of one of the vertebræ, is illustrated at b, in Fig. 137. It exhibits four different structures: the first is an outer coating of dense areolar tissue; the second occupies the centre, and consists of a gelatinous substance somewhat resembling simple cartilage; the third composes the radii, which extend from the centre to the circumference; whilst the fourth consists of a firm fibrous structure, occupying all* the

FIG. 137.



Sections from a vertebra of the Vaagmaer: a, three rows of cells from one of the radii; b, transverse section of a vertebra; c, one row of cells showing the tube passing through them; d, vertical section of one of the radii.

spaces between the radii, and forming the great mass of the body of the vertebra. is to the radii that I would particularly direct attention. as these are composed of a material very like cartilage. In a thin section of the same vertebra, in which five rays may be distinctly seen, each is found to be composed of a series of thick-walled cartilaginous cells, arranged in linear series; some of the rays, as at c, being composed of one row; others, as at a, of three rows of cells. On carefully examining any one of these rows, an aperture of com-

munication between the cells may be observed, and through each of these apertures a delicate tube passes from the centre to the circumference. Whether this tube be a blood-vessel or an absorbent cannot be easily ascertained; its tubular character, however, is very evident. In a longitudinal section of a vertebra of the same fish, the radii, represented by d, are divided in the direction of their long diameter: they generally occur in bundles of three or four, and the apertures through which the tubes pass may be observed at

nearly equal distances from each other, and occasionally, portions of the tubes may be seen in the apertures. Dr. Reid described very accurately the structure of the vertebræ, and the arrangement of these thick-walled cells, but he appears to have overlooked the tubes passing through them, as no mention is made of their presence in his paper on this subject, published in the "Annals of Natural History" for 1849. They may not be present in every vertebra, but they are so in all the specimens in my possession, which were transmitted to me from Dr. Reid himself, through the hands of Dr. Macdonald.

FIBRO - CARTILAGE.

Before concluding the subject of cartilage, there is another tissue which, from being composed of two simple elements, fibrous tissue and cartilage, has been placed by Messrs. Todd and Bowman, as the last of the tissues in their table, namely, fibro-cartilage; but, as this is so intimately connected with some of the forms of articular cartilage and enchondroma, I think it best to consider it in this place.


Fibro-cartilage exists principally in articulations, occurring in the form of discs between the vertebræ in the human subject; these are highly elastic, and serve to diminish the shocks to which this portion of

of immense size; and the articular surfaces of the vertebræ in many of these animals not being anchylosed to the bodies, may also be found detached as discs. These last are smooth and a little concave on their articular surface, convex and rough on their epiphysal; I remember when I first became connected with this College, that the late Mr. Clift told me he once met with a specimen about three or four inches in diameter, in the shop of a dealer in curiosities in the Strand, who gravely informed him more than once that he valued it above everything in his shop, as he considered it a most rare and beautiful example of a fossil crumpet, which it much resembles in appearance.

These discs are very abundant on the sea-shore in northern climates; and when Her Majesty's ship Hecla was wrecked, I was informed by one of the officers, that they served the erew of that unfortunate vessel as plates. The discs of fibro-cartilage between these vertebræ are often more than two inches thick. Masses of fibro-cartilage exist in certain joints, under the name of interarticular fibro-cartilages. We have examples in the temporo-maxillary and sterno-clavicular articulations of the human subject. When either of these is divided transversely, and a section made sufficiently thin for examination by the microscope, it will be found that it is composed ehiefly of a net-work of fibres, within which a few cartilage cells are enclosed.

The eells, Fig. 138, A, are large and tolerably

numerous, in sections of the outer portion of an intervertebral disc; in the centre, which is more pulpy than the outer part, the fibres are fewer in number, and the cells, as shown at a, b, larger and more abundant. Many of the examples of articular cartilage, especially those from fishes and birds, as well as specimens of



A, fibro-cartilage from outer part of an intervertebral disc; ab, cells from the centre of the same disc. B, three dorsal vertebræ showing ossification of the anterior common ligament, with absence of the intervertebral disc. c, fibro-cartilage of epiglottis.

enchondroma, are composed of fibro-cartilage, characterised by the presence of white fibres and cartilage-cells.

Modified forms of fibro-cartilage exist in the epi-


glottis of the human subject, as shown at c, and in the auricles of the ear of the larger mammalia. In these situations, the cells are readily separated from the matrix; indeed, they often drop out, and leave the fibrous framework entire.

Fibro-cartilage is not so prone to ossification as the simple fibrous structures; for in two examples of anchylosis of the vertebræ, selected from a large number in the Museum, the discs have, in each case, disappeared, and the ossific matter is confined to the fibrous tissue, or anterior common ligament, which binds them firmly together; in some instances, the space originally filled by the disc is still present, but remains unoccupied by bone. A striking specimen, illustrative of this fact, is represented in Fig. 138, B. In the two upper vertebræ, the ligament has become ossified, and a considerable quantity of bone has been formed, so as to cover the anterior part of the inter-vertebral space; there is, however, no junction between the two vertebræ, but between the second and third the union is complete, and in both cases, the space originally occupied by the inter-vertebral disc is empty. This fact is quite as evident in the spine of the lower orders of mammalia as in man, and is very common in the Horse and Sheep.

LECTURE XVI.

ADIPOSE TISSUE.

Next in our order of arrangement, among the true cellular tissues, is Adipose tissue, which consists of cells having walls composed of structureless membrane, containing within them the substance known as fat. This exists, in ordinary animals, in three states, either as oil, lard, or tallow; but in one species of Whale, the Physeter macrocephalus, it is in the form of spermaceti. The three first being distinguished by their relative firmness at ordinary temperatures, while the last differs in toto in its chemical properties. In the majority of works on anatomy, and in common parlance, the term fat is given to masses of these cells, but in modern science adipose tissue is the name applied to the mass of cells, and fat to their contents. The cells of adipose tissue, when occurring singly are globular, but when subjected to



Cells of adipose tissue.

pressure as in Fig. 139, they are, like vegetable cellular tissue, of a more or less dode-cahedral shape. In the young subject, they vary in diameter from $\frac{1}{300}$ th to $\frac{1}{800}$ th, and in the adult, they are rarely smaller than $\frac{1}{700}$ th of an inch. When first developed in the embryo,

they have a nucleus, but this very soon disappears. Adipose tissue exists in the form of lobules, which may occur singly or in masses of many pounds' weight. A certain amount of it is considered a sign of health, but an excess, especially if it be in one mass, as in certain tumours, constitutes disease.


In all works on anatomy and physiology, even of so late a period as last year, it is distinctly stated, that adipose tissuc exists in invertebrate animals; this, however, I find to be incorrect, and it cost me no small amount of labour to prove it. Fat certainly does exist in insects, crustacea, and mollusea, but no true adipose cell is ever present; it could not be nourished without its accompanying blood-vessels, and these are not found in invertebrata. The tissue resembling adipose tissue usually belongs to the liver or other glandular organ, and the fat exists in its cells in the form of oil.

In the liver of the larva of a Goat Moth, Cossus ligniperda, which consists of a series of cells or vesicles, containing a large number of globules of oil, and again,

in another specimen, taken from a *Cockroach*, there are tubes, also full of oil globules, but in neither case, and not even in the *Cephalopoda*, is the oil contained in adipose cells.

As soon, however, as we pass the barrier between the invertebrate and vertebrate sub-kingdoms, we find that even in the lowest members of the class of fishes, true adipose cells occur, and all are doubtless aware, that in the liver of the Cod, and of many cartilaginous fishes, fat exists in the form of oil without any adipose tissue;—in this particular, the liver resembles that of an invertebrate animal. If a portion of the liver of a Cod be examined, it will be seen, that with the exception of a

FIG. 140.



Oil globules from the liver of the *Cod*.

few secreting cells, it is composed of a mass of oil globules of various sizes, as shown in Fig. 140; but in a specimen from the peritoneum of the same Cod, a mass of adipose cells full of a dark brown liquid oil are visible. The adipose cells are grouped in lobules, and surrounded by

areolar tissue, which invests the lobules and supports the capillaries distributed to the cells. This is well exhibited in two preparations, one of the subcutaneous adipose tissue of a young child, the other of the same tissue from an adult, in which the adipose cells far exceed those of the child in diameter.

Hunter's observations on this subject, contained in the third volume of the Physiological Catalogue of the Museum, are so valuable as to merit an attentive perusal; they contain, however, a few trifling errors which arose from his not being aware of the existence of the adipose cell, especially in the Whale, the fat of which animal he supposed to be contained in the interstices of the fibres that constitute a great portion of the blubber. A preparation put up to demonstrate this fact is still in the Museum, but the receptacles it exhibits are for the lodgment of the adipose cells containing the oil, and not for the oil itself. Another preparation, is a section of the skin of a Whale, taken from near the tail, also put up by Hunter, to show that in this part of the animal there are no receptacles for the oil. The chief value of the adipose tissue of the Whale tribe consists in its containing oil which is liquid at ordinary temperatures, the train oil of commerce; the fat of the Bear is also very valuable on the same account, and is largely employed by the perfumer. I once had an opportunity of dissecting a Bear, and, although it was during frosty weather, the quantity of oil that flowed from between the muscles was very great, amounting to many gallons, and hence I concluded, that the value of the oil of the Bear, or grease—as it is usually termed—to the perfumer, depends on its continuing in a fluid state, even at so low a temperature. The nearest approach to the fat of the Bear, is perhaps, the marrow from the extremities of the long bones of ruminants, which is often

used as a substitute for bear's grease. From the whole of this tribe we procure a very firm fat known as tallow; in the Piq it is still firmer, and ealled lard; if a portion of either of these substances be cut transversely, the fat is so dense as to allow of the adipose cell being divided, and such tallow or lard is perfectly solid in all temperatures of this climate, and is eapable of being employed in the manufacture of eandles. In the Turtle, there is under the earapace a quantity of adipose tissue contained within delieate areolæ, formed principally of loose white fibrous tissue, and is well known as the green fat, being much prized by epicures. This fat is more digestible than that from ordinary animals, and I have frequently heard it remarked, that those persons who eannot easily digest the fat of beef or mutton, ean eonsume large quantities of the fat in question without annoyance.

I have stated, that the adipose tissue is supported by a fibrous net-work; this holds good in the majority of eases, but, in the marrow of bones, no supporting tissue is present, and I here take the opportunity of demonstrating a point of some interest, which is of eonstant occurrence throughout the vertebrate kingdom, viz., that in the bones of the arm or leg, the marrow of the humerus and femur is exceedingly firm, whilst that of the lower part of the tibia or metatarsal bone is always more or less liquid, and it is from these bones that the ordinary neat's foot oil is procured. The same thing is evident even in skeletons, in those of birds, it may be

observed that the boncs of the extremities are greasy at their distal ends, because the contents of the adipose cells are always of a liquid nature, and have little or no tissue to support them.


According to chemists, fat consists of a liquid and solid principle, the former being termed elaine, the latter margarine; in all the firmer fats, such as lard, a third principle, termed stearine, is also present. When speaking of oil in vegetables, I noticed the analogy between the fatty substances derived from plants and animals, and I mentioned that a stearine had for some time been procured from palm-oil, either by distillation, or by subjecting it to great pressure. Stearine may be obtained from human fat, and cannot readily be distinguished from that of vegetable origin.

It is generally known that the elaine and stearinc of the fats, arc compounds of the oleic and stearic acids, with a base termed glycerine, a substance now frequently employed in medicine, and also in mounting microscopical objects. This material is extensively obtained in the manufacture of soap. When oils or fats are boiled with caustic alkali, the compounds of stearie, oleic and margaric acids with glycerine, are decomposed, the former combining with the alkali, while the latter is set free, and is found in the soap-lees; from which it is separated by a chemical process. The purest glycerine, which is almost colourless, is obtained in the manufacture of lead plaster. In this process, oil, litharge (oxide of lead) and water arc boiled together, the oxide

of lead combines with the fatty acids, and the glycerine is separated in a nearly pure state.


It sometimes happens, especially in the adipose tissue of old persons, that the solid element, or margarine, separates in the form of acicular crystals from the elaine. In a specimen of adipose tissue from a female, seventy years of age, every cell (Fig. 141) contains a stellate mass of these needle-shaped crystals. This fact is more strikingly exemplified in the adipose tissue of the *Sper*-

FIG. 141.



Adipose cells, each containing needle-shaped crystals of margarine.

FIG. 142.



Adipose tissue of Sperm-whale: A A, cells containing crystals of spermaceti. B B, crystals of spermaceti on the outside of cells.

maceti Whale, the adipose cells, as shown by Fig. 142, are of large size, and in every cell may be seen the crystalline substance known as spermaceti. This substance, however, as represented by AB, frequently occurs in masses of needle-shaped crystals external to the cells, but in some instances tabular crystals are more common in these situations than the acicular.


The subcutaneous adipose tissue is most abundant in animals destitute of hair, such as the Whale, and even man himself; in other animals, especially ruminants, it is stored up around the loins and kidneys, and that obtained from the latter situation has received the name of suet. The Whales, on the contrary, have no adipose tissue within the cavity of the abdomen, and some wild animals, especially the Hare, have seldom a trace of adipose tissue in any part of the body. In fishes, adipose tissue occurs among the muscles; but in the Cod and some of the cartilaginous species it is found in the liver in the form of oil. In reptiles, as the Frogs and Newts, it is stored up against winter in the form of long cœcal appendages situated above the testicles. If a Froq be examined in the autumn, these appendages will be found of large size; but by the spring they will have nearly disappeared. The same thing occurs amongst hybernating quadrupeds, the Dormouse and Hedgehog for example, but in these animals the fat is principally subcutaneous, forming in fact a winter overcoat. In serpents the fat exists in the form of lobules attached to the sides of the mesentery; in turtles it is principally beneath the carapace; in birds it is chiefly subcutaneous, and in the peritoneum; in many of these animals, especially those of the order palmipedes, the adipose tissue contains an abundance of oil.

Having made frequent allusion to a peculiar kind of fat found in the *Sperm Whale*, and known by the name of spermaceti, I will now describe it more parti-

cularly. It is a crystalline substance, of a white or vellowish white colour, which is deposited in the large cranial cells or interspaces of the Spermaceti Whale. During the life of the animal it is liquid, but it readily crystallizes and scparates after death. It was formerly considered useless, and hundreds of tons have been thrown into the Thames, but it is now largely employed in the manufacture of candles, and is even more valuable than the oil itself, from which it separates in flaky crystals. When spermaceti is exposed to a heat of 1.15° Fahrenheit, it melts, and again crystallizes on cooling; in this condition it has the property of polarizing light, and exhibits a beautiful series of colours. The same power of polarizing light is also possessed by a substance somewhat like spermaceti, produced by the decomposition of certain animal tissues, especially the muscular, and known by the name of adipocirc. A fine specimen of this material, formed from the thigh of a man by the action of water, is prescried in the College Museum. Like spermaceti, adipocire is easily melted, and crystallizes on cooling.

I have already stated that the cells are arranged in lobules, and that each lobule is invested by areolar tissue, which passes among the cells, and gives support to the blood-vessels. The blood-vessels rapidly divide into a capillary network, which not only surrounds the individual lobules, but branches in the form of loops encircle each fat-cell. These capillary loops are represented, in Fig. 143, under a magnifying power of two hundred and fifty

FIG. 143.




Portion of an injected lobule of human adipose tissue.

diameters. This vascular network can only be demonstrated by injected specimens, and the arrangement of the vessels may be rendered more distinct by dissolving out the fat by oil of turpentine. When large masses of adipose tissue are treated in this way, the walls

of the cells are readily distinguished when viewed by transmitted light. In birds, the adipose cells are of smaller size than in mammalia, and when injected the capillary net-work is closer. Small lobules of adipose

FIG. 144.



a b, injected lobules of adipose tissue from the skin of a Pigeon; c, portion of the same magnified 200 diameters.


tissue sometimes surround the pulps upon which the feathers are formed, and the capillaries of these lobules very much resemble those of some glandular organs, as is strikingly exemplified in a portion of the skin of a young *Pigeon*, Fig. 144. At a and b, are shown the capillaries of two lobules, magni-

fied forty diameters, and at c, a portion of one of the same, magnified two hundred diameters.

When speaking of the adipose tissue of the Whale, I mentioned the receptacles between the fibres for the lodgment of the cells; a specimen of this tissue would,

when injected, exhibit an extraordinary amount of vaseularity. An injected specimen of the blubber of a young *Porpoise*, is represented by Fig. 145. The upper part

FIG. 145.



Vertical section of the skin of a young *Porpoise:* a, cuticle; b, papillæ; c, vessels of the adipose tissue, or blubber.

of the preparation, α , exhibits the non-vaseular eutielc, into which long papillæ, b, project; below these are seen numerous trunks, c, giving off capillary vessels which belong to the adipose tissue. The amount of vascularity, therefore, of the blubber of a large Whale must be truly astonishing, if we may judge from this specimen from one of the smallest of the eetaeea. A somewhat similar structure to that of the blubber of the Whale occurs in the Piq, which, when salted and dried.

is known as bacon. In this substance the eapillaries, when successfully injected, are as numerous as in other specimens of adipose tissue.

The adipose tissue of young animals as in children, generally consists of a series of small oval, or eircular lobules seattered throughout the areolar tissue; if the injection be perfect, it will be found that the capillaries of the cells are much smaller than those of the adult

human subject, represented in Fig. 143. In fatty tumours, the fat is usually more or less yellow, and contains few if any vessels which bleed on its removal or when cut; but it sometimes happens that the mass is of a red colour, and contains a very large amount of blood. A tumour of this character, removed after death from the mesentery of a child, by Mr. Pittard, formerly one of our students, had at first sight an appearance like that of malignant disease, and when an incision was made into it, a large quantity of blood escaped. Its true nature was only ascertained by microscopic examination, when it turned out to be nothing more than a mass of adipose tissue, and although it has been preserved in spirit and exposed to the light for the last two years, it still retains a portion of its red colour.


LECTURE XVII.

FATTY DEGENERATIONS.

There is a particular morbid condition of many distinct organs, in which fatty matter, or true adipose tissue, takes the place of healthy structures. The first of these abnormal changes is called fatty degeneration, but to that of the second the term fatty substitution might be applied; they occur most frequently in muscular tissue. Before, however, I can present a satisfactory idea of the commencement of this disease, it will be necessary for me to describe very briefly the normal anatomy of muscular tissue. For our knowledge of the minute anatomy of muscle we are principally indebted to the labours of Mr. Bowman, and the account I shall give will differ but little from that published by him in the first part of the "Philosophical Transactions" for 1841. Every voluntary

muscle consists of a series of fibres, each of which is termed a *fasciculus*, and each fasciculus is invested by a sheath of structureless membrane, termed the *sarcolemma* (Fig. 147, B). The muscle itself is surrounded

FIG. 146.




Capillaries of voluntary muscle.

by areolar tissue, which dips in among the fasciculi, and gives support to the bloodvessels and nerves. If a muscle be injected, the eapillaries, as represented in Fig. 146, are readily seen; they run in parallel lines between the fasciculi, and transverse connecting branches are given off

at tolerably regular distances.


The fasciculi exhibit transverse and longitudinal striæ, but, in most cases, the former, as shown in Fig. 147, A, are more plainly exhibited than the latter. In some animals the fasciculi break up transversely, in others longitudinally, so that, in the one ease, we have a series of dises, and in the other numerous filaments termed fibrillæ. The fasciculi of the *Eel* readily break up into dises, whilst those of man and most mammalia commonly separate into fibrillæ. If the flat surfaces of the dises, at F, in Fig. 147, be examined, they present a granular aspect, which is due to their being made up of the ends of the fibrillæ; and, if the fibrillæ be viewed with a power of five hundred diameters each one, as shown at B, will exhibit a beaded structure, the



A, portion of a fasciculus of voluntary muscle of an *Eel* splitting up into discs. BC, fibrillæ. D, fasciculus invested with its sarcolemma. E, masses of myoline detached from the fasciculus.

part forming the bead being a minute portion of muscular substance, termed myoline; but if the power be increased, the masses of myoline (c) will be found to be surrounded by a thin cell-wall. In the muscular substance of the *Eel*, the structureless sarcolemma surrounding a fasciculus (Fig. 147, B) is readily seen. In Fig. 148 is represented a portion of the muscle of a *Pig*, prepared by

Mr. Lealand; on the lower edge of one of the fasciculi are distinctly seen the very delicate fibrillæ aa, and, if the power with which this specimen is examined be sufficiently high, the central mass of myoline b may



Fasciculus of voluntary muscle of a Pig: a a, fibrille; b, enlarged representation of one of the fibrille.

be noticed in each little cell. The fibrilæ vary in size in different portions of the same muscle, in the *Skate* and *Shrimp* I have seen specimens more minute than $\frac{1}{60000}$ th of an inch; in these cases the particles of myoline can be discerned, but not the cell-wall around them. If the fasciculi have been kept

long in spirit or acid, the fibrillæ break up, and the masses of myoline are found either in short lengths or in detached

pieces, in all parts of the field of view. A specimen of this kind from the *Eel* is represented in Fig. 147, E; the same thing is very common in other fishes, especially the *Lamprey* and *Lancelet*.

Before fatty degeneration commences in voluntary muscle, the transverse striæ disappear; and I have long known that the first trace of this disease is marked by a disturbance of the particles of myoline, which appear as so many very minute granules scattered irregularly within the sarcolemma, leading one to suppose that the delicate cell around each particle had given way, thereby allowing the myoline to escape, and destroying all regularity both of the transverse and longitudinal markings. As the disease progresses, the myoline is replaced by minute, highly-refracting globules of oil, until at last the whole sheath is full of them.

In a specimen of this diseased condition of mus-

FIG. 149.

A B

A, fasciculus of voluntary muscle full of oil globules. B, fasciculus of voluntary muscle exhibiting a disturbance of the particles of myoline.

cle from the human subject, Fig. 149, B, the transverse striæ are visible in the upper part, and a partial disturbance of the myoline in the lower; in another preparation (Fig. 149, A) the disease has so far advanced that all trace of striæ is completely lost, and globules of oil, in this case of nearly equal size, but in others of variable diameter, occupy the sarcolemma. The

fibres of the heart are very subject to fatty degeneration, and for our knowledge of this disease we are, in a great measure, indebted to the labours of Dr. Ormerod; but the subject has been lately investigated with great care by Dr. Richard Quain; and in his paper, published in the fifteenth volume of the "Medico-Chirurgical Transactions," you will find all that is at present known respecting it. A very excellent example of fatty degeneration of the muscular fibres of the heart, is one taken from a man a hundred and three years of age, for which I am indebted to the kindness of Dr. Edward Smith, it exhibits the transverse striæ in some parts, but in others, as in Fig. 142, A, these are replaced by highly-refracting globules of oil. I have lately had the opportunity of examining a most interesting case in which the disease was present in the voluntary muscles of the extremitics. In one family of nine children, six of whom were girls and three boys, all the girls were perfectly healthy, but the boys, on arriving at the age of three or four, began to lose the use of their limbs. One of them, the eldest, has lately died, and, on examination of the brain and spinal chord, both were found to be healthy, the muscle, however, had not only undergone fatty degeneration, but the fasciculi themselves were much diminished in size, which would, of course, account for the want of power in the limbs. This disease from the first was supposed to be seated in some part of the nervous system, probably arising from imperfect innervation of the muscle;

but the discovery of its real seat will, it is to be hoped, lead to such a mode of treatment as may be beneficial to the two afflieted survivors.

Fatty degeneration of voluntary musele is very eommon among the lower animals. All are aware, no doubt, of the difference in colour between the flesh of the breast of a Fowl and that of its legs, the former being more or less white, the latter of a reddish hue. If that from the breast be examined, traces of fatty degeneration are very apparent in the faseieuli, arising, no doubt, from the want of use of the musele in the aet of flying. The museular structure of the legs, on the eontrary, which are always in exercise, is perfectly healthy. I have detected the same disease in the museles of the legs of the Ostriches kept for a long time in the Zoologieal Gardens. These from want of use, are perceptibly whiter than those employed merely in the support of the body, and also exhibit a eonsiderable amount of fatty degeneration. The same disease I have also found in their bones; and it is a faet well known to the keepers, that when they onee take to lying down, their doom may be said to be sealed. A knowledge of this eireumstanee may perhaps lead to the more suecessful preservation of these gigantic members of the feathered tribe. In the Sheep many instances are on record in which almost the entire museular substance of the trunk has been converted into adipose tissue. In the Pathological Series is a mutton chop, Prep. 10, nearly all the museular substance of which

is replaced by adipose tissue; in this case the disease no doubt first commenced in early life, and as the fasciculi diminished in size, the adipose tissue was deposited to occupy their place, forming a good instance of fatty substitution. A portion of a fasciculus, surrounded by adipose cells, is represented in Fig. 150, B, and an entire fasciculus, full of nearly equal-sized

FIG. 150.
A
B

A, fatty degeneration of voluntary muscle. B, substitution of adipose tissue for voluntary muscle.

globules of oil, in Fig. 150, A. The leg of a man thirty-five years of age was removed a few months since at King's College Hospital, by Mr. Partridge, in which all the muscles had been replaced by adipose tissue. The limb had been rendered useless by an attack of paralysis, occurring three years after birth. Only a few fasciculi could be

diseovered amongst the adipose cells, and these were of very small size and perfectly healthy. Fatty degeneration appears also to occur in osseous tissues, and indeed the disease termed mollities ossium is of this nature. All bones so affected have thin walls, are always more or less soft, and contain an abundance of oil. I have examined the bones in several cases, and found that the disease first commences in the bone cells, the cell itself becoming larger and larger, its canaliculi disappearing, and several of these cells uniting to form a cavity, in which oil globules soon make their appearance,

all the parts of the bone in the neighbourhood of the cells, becoming at the same time thin and transparent from the removal of the granules of earthy matter. In a vertical section of the lower end of the tibia, from a case of mollities ossium; the walls of the bone are exceedingly thin, and so soft as to be readily cut with a knife, the entire medullary cavity, and all that remains of the cancellated structure, being full of oil and adipose tissue.

There are still one or two more examples of fatty degeneration of tissues, of sufficient importance to merit notice. The first of these is a deposit of fatty matter between the layers of the cornea, giving rise to the arcus senilis. The discovery of the true nature of this change, which is common in the eyes of old people, is due to Mr. Canton, who has published an account of it in the "Lancet" for 1850.

In a vertical section of a cornea, given to me by Mr. Canton, the arcus senilis has formed, and between some of the layers of which the cornea is composed there is a deposit of small, highly-refracting globules of oil. The arcus is not situated in that part of the cornea joining the sclerotic, but a little nearer the centre, there being generally a transparent zone between the sclerotic and the arc. The last example of degeneration of tissues of which I shall speak, is the formation of Adipocire, which takes place after death, when flesh is exposed to a certain amount of moisture, or to a running stream of water. A preparation of the thigh of

a human subject converted into adipocire, was presented to the College some years ago by the late Regius Professor of Medicinc at Oxford, Dr. Kidd. It was a portion of a subject which had been partially dissected, and, not being further required, was placed in a pit near the dissecting-room, into which fresh water was occasionally admitted. When a mass of adipocire is handled, it will be found to have a soapy feel, and to possess little or no odour. On exposure to heat, it readily melts like spermaceti, crystallizcs again on cooling, and also polarizes light. When examined microscopically in the fluid state, or after solution in ether, no trace of muscular substance remains, the residue consisting principally of areolar tissue. A thin slice exhibits no appearance of cellular structure. Adipocire, like spermaceti, is capable of being made into candles; and on the discovery of this substance, many years ago, a patent was taken out for the conversion of the offal of slaughter-houses into adipocire, but the process died with the patentee. A very large quantity of adipocire was found at the commencement of the present century in the burial-ground of Lcs Innocens at Paris, where one thousand five hundred bodies had been buried in one pit, most of which had been converted into this material by the action of water. A considerable quantity of adipocirc was once formed in a pit in this College, into which all the parts removed in dissection are thrown; the adipocire, however, in this case, was derived principally from the muscular substance of a young Elephant.

LECTURE XVIII.


PIGMENT.

I now arrive at the description of the last variety of permanent cellular tissue occurring in the human subject, pigment. This consists either of a solid or fluid matter, contained within a cell-wall. When speaking of pigment in vegetables, I stated that, in some few cases as in Ferns, we had colour belonging to the cell-wall itself, but in all other instances, the various and beautiful hues of flowers were produced by a fluid colouring-matter occurring within the cells. We have likewise a striking example of fluid colouring-matter in animal cells, in the blood discs, but in animal tissues generally the pigmental matter occurs in the form of granules. The most striking examples of pigment occur in the eye, upon the choroid coat of which is a distinct layer of hexagonal cells, most probably cpithelial, termed the

pigmentum nigrum, within these are found innumerable minute granules, which, if examined soon after death, exhibit a molecular movement. The size of these granules does not generally exceed the $\frac{1}{20000}$ th of an inch in diameter, the depth of the colour depending upon the quantity aggregated within a certain space. The granules are said to consist of a peculiar animal principle, the chief constituent of which is carbon, and on this account neither the strongest acids nor chloring is capable of destroying the colour. The most common form of the cells is hexagonal, as in the pigmentum nigrum before described; but between the choroid and sclerotie they are somewhat fusiform, and oecasionally have bifid extremities. In some of the lower animals, as the reptiles and fishes, they have a stellate appearance, and in some cases bear a strong resemblance to bone cells. The principal seat of pigment in animals is the cuticular layer of the skin, in which it also occurs in hexagonal cells. In the negro the black colour was formerly supposed to occur only in a distinct layer of the skin, termed the rete mucosum, but such is now found not to be the case, the rete mucosum being merely the layer of cutieular cells last formed, and which contain pigmental granules. Similar cells are present in Europeans, those of the last deposited layer being always darker than those nearer the free surface; the difference in quantity of the pigment in the cells gives rise to all the varieties of colour which the skin of the human race presents. In Albinocs the cells

are present, but the granules, if not absent, are so few in number as to oeeasion little or no shade of colour. The skin of a *Lamprey*, Fig. 151, exhibits a number of

FIG. 151.



Pigment cells of the skin of a Lamprey.


large cells of stellate figure, somewhat like the lacunæ of bone. The red spots on the skin of the *Plaice* are produced by the presence of minute irregular cells occurring in great abundance between others of large size, which are of a black colour, and stellate figure. Pigment-cells are very frequently found in the

peritoneum of fishes and reptiles.

In the skin of a Newt (Triton palmatus), the pigment is very abundant, giving a marbled appearance to the animal; the cells are of a stellate figure, and are chiefly arranged in circles, the central part of the circle being a folliele, which, in some cases, contains granular matter. Numerous pigment-cells are present in the skin of the web of the frog's foot; these always excite attention when the circulation in the capillaries is seen for the first time. In some of the Reptiles there are pigment-cells containing a white material; in the iris of a Tiger-boa (Python tigris), a collection of white pigment-cells exists, in which the branches of the cells are of great length, and when these are viewed by transmitted light they appear perfectly black. I am still uncertain whether the branches of these cells ever

communicate or anastomose; in the majority of cascs they certainly do not, but in the last named instance this point cannot be easily determined. The pigmentum nigrum from the eye of a Sheep, Fig. 152, A,

FIG. 152.



A, pigmentum nigrum of the eye of a Sheep. B, pigmentum nigrum of the eye of an adult human subject.

FIG. 153.



Pigment-cells between the choroid and sclerotic coats of the eye of a *Sheep*.


consists of cells of hexagonal figure, full of black granules; if, however, the posterior part of the choroid be examined, as in the specimen, Fig. 153, it will be found that some of the cells are fusiform, whilst others are bifurcated at one or both extremities; in the centre of each cell is a large

white spot, which is the nucleus.

In the *pigmentum nigrum* from the eye of an adult human subject (Fig. 152, B), the cells are much larger than those in the *Sheep*, and the granules also more

plainly seen. In Albinoes, that is in the white offspring of black parents, the hexagonal eells are present, but

FIG. 154.




A, pigmentum nigrum of a black *Rabbit*. B, cells from the choroid of a white *Rabbit* in which the pigment is wanting.

the pigment is so small in quantity as to allow the vaseular ehoroid to be seen through the pupil, giving a pink appearance to the eyes of these people. The same may be said of white *Rab*bits and *Cats*. A portion of the pigmentum of the ehoroid of a black *Rabbit* is represented in Fig. 154, A, the

eells are full of granules, and similar eells from a white Rabbit, in which the pigment-granules are either absent or void of eolour, are shown at B; the eyes of these animals are, from this eircumstance, more or less pink.

In the pigmentum nigrum from the eye of a feetus,

FIG. 155.




Pigmentum nigrum from the eye of a human fectus: A, cells magnified 250 diameters. B, portion of the same magnified 600 diameters.

the cells are not only smaller than those of the adult, but the granules also are fewer in number, and the centre of each cell is occupied by a large white spot, the nucleus. A portion of this pigment, as seen under the same magnifying power as that of the adult, in Fig. 152, B, is represented in Fig. 155, A, and

in order to show the nucleus and the granules, the portion B is magnified six hundred diameters.

I shall next describe the pigment developed in the skin, but must first state, that, until the last few years, a coloured layer, distinct from the cutiele, and termed

FIG. 156.




A portion of the skin of a $Negro: a \ a$, eutiele; b, portion of eutiele reflected; c, layer termed $rete \ mucosum; \ d$, true skin.

rete mucosum, was supposed to exist in the skin of the Negro; a similar layer was subsequently found in the skin of the white man, but devoid of colour. In two preparations by Hunter, the one, the skin of a Negro, represented by a a, in Fig. 156, in which, in addition to the eutiele b, the black layer, c,

termed rete mucosum, is shown; while in the other, which is the skin of a European, a similar layer, unstained with pigment, is turned down; these layers,

FIG. 157.




Vertical section of enticle of the Negro, showing that the last formed eells contain most pigment.

however, are nothing more than the last formed portions of the eutiele.

The vertical section of the skin of a Negro, Fig. 157, serves to show that not only the cells, but the pigment also, is most abundant in the deepest layer, and, as you ascend towards the free sur-


face, the pigment decreases in quantity, and the cells become flattened into the form of scales, from which the pigment is absent. A portion of the cuticle of a Negro with its inner or attached surface uppermost, which has been dried before being mounted in Canada-balsam, exhibits cells of a more or less hexagonal figure, containing the pigment. In another specimen from the same skin, Fig. 158, the disposition of the cells is better

FIG. 158.



Pigment cells on the under surface of the eutiele of a Negro.

FIG. 159.



Vertical section of the skin of the nose: A, euticular layer having a band of pigment on its attached surface. BB, hairs and schaceous follicles.

shown. The preparation is mounted in fluid; and it may be noticed that the surface is pitted, and that the greater part of the pigment-cells are arranged around the depressions, into which the papillæ of the true skin are rcceived, these depressions being the white parts of the figure. A vertical section of the skin of the nose having a stratum of black pigment in the decpest portion of the cuticle, while the upper part is quite white, is represented in Fig. 159. The specimen is interesting in another point of view, as several hairs, B B, are visible in the cuticle, cach having sebaceous follicles on either side; many of these follicles are enormously dilated, and within them a quantity of dark granular matter is contained, thus affording a good example of one of the forms of the disease called acne, which essentially consists of an enlargement and suppuration of the sebaceous follicles. The colour of the hair frequently depends upon the pigment developed in the cuticle; and it is a well-known fact, that in those animals, as the Pig, in which there are occasional patches of black hair, the skin from which such hairs grow is generally black; so, in the Albino, the hairs are white, because no pigment is secreted.

Pigment is developed in peculiar situations under certain circumstances, as around the nipple during pregnancy, and in certain spots on the face, called freckles, after exposure to a summer's sun; in both these instances the colouring matter disappears on the removal of the exciting cause. Pigment is sccreted by the Cuttle-fish in a special gland, termed the ink-bag, and this pigment is largely employed by the artist, under the name of sepia. Pigment is also secreted in cells in certain states of disease, as in melanosis a striking specimen of which in both ovaries of a female is preserved in the Museum. In the lung of a Calf, here and there, a lobule is quite black, whilst all the others are perfectly white. A section of one of these black lobules, demonstrates that the pigmental matter is deposited in the form of irregular granules in the parenchyma of the lung, and such lobules

would no doubt receive nearly if not quite as much air as the whiter ones.

There are some rare instances in which pigment is secreted in large quantities on the skin of the human face. A remarkable case of this abnormal secretion of pigment is described and figured by Mr. Teevan in the "Medico-Chirurgical Transactions" for 1844.

I possess some pigment given me by Mr. Squire, the distinguished chemist in Oxford Street, which appeared under the right cye of a young woman, and was capable of being brushed off with a camel's-hair pencil. This is contained within cells like those of cpithelium, and consists of granules. Mr. Squire tried various experiments upon it, and found that he could not remove the colour either by the strongest acids or by chlorine.

Having now arrived at the period when my labours, for the present session, must be brought to a close, it only remains for me to thank you most sincerely for your kind attention, and to express a hope that I may, in some measure, have succeeded in imparting such knowledge as may be useful in the practice of your Profession; and if so, I shall feel that I have accomplished one of the great objects for which this Institution was established, viz.: the advancement of those sciences conducive to the alleviation of human suffering, and the good of mankind.

INDEX.

Α.	Birds, fat of, 185.
A dimension 2007	— adipose tissue in, 190.
Adipocire, 201.	—— capillaries of adipose tissue
Adipose tissue, 180. —————————— of Turtle, 185.	in, 190.
of Reptiles, 188.	Boletus, cells of, 25.
Tessels of 180	Bone, ivory-like deposit in, 160.
vessels of, 189. capillaries of, 189.	— structure of, 168.
in Birds, 190.	fatty degeneration of, 175,
in Porpoise, 190.	199.
in Children, 191.	Burr-reed, cells of, 24.
Albinoes, 203, 206.	C.
Alder, ducts of, 105.	
Algæ, 56.	Cactus, raphides in, 45.
Aloe, cuticle of, 13.	Campanula, fibre of, 13. Camphor, 41.
Anchusa italica, hairs of, 12.	Canada-balsam, 42.
Animal tissues, 115.	Canes, silica of, 52.
classification of, 117.	Canna bicolor, vessels of, 98.
— membrane, 117.	Capillaries, naked, 134.
— cells, 117.	of adipose tissue, 189.
Antheridia of Mosses, 72.	——— in Birds, 190.
Apple-tree, raphides in, 46.	in Porpoise, 190.
Arcus senilis, 200.	——— in Children, 191.
Areolar tissue, 127.	in muscle, 194,
of Elephant, 128. development of, 129.	Capsule of lens, vessels of, 130.
of Sheep, 129.	Cartuage, 135.
	vessels in, 141.
Arrow-root, starch of, 29.	— articular, 143.
Articular cartilage, 143.	vessels in, 149, 151,
in Fishes, 146.	155.
vessels of, 149, 151,	— in Fishes, 140, 171.
155.	—— in Reptiles, 140. —— discased, 154.
Artificial raphides, 47.	ossifying, 168, 172.
В.	of Rana paradoxa, 171.
	Cassava, starch of, 21.
Balsam, vessels of, 8.	Cells, nature of, 17.
leaf of, 23.	— typical forms of, 17.
Bamboo, silica of, 51.	—— stellate, 24.
Bear, fat of, 184.	—— contents of, 26.
Berghmehl, 57.	—— animal, 118.
Birds, articular cartilage in, 147.	Cellular tissues, animal, 135.

Cetacea, fat of, 181, 184.	Elæagnus, raphides in, 46.
fibro-cartilage of, 178.	Elder, pith of, 9.
Chara vulgaris, spores of, 73.	Elephant, arcolar tissue of, 128.
China-grass, woody tissue of, 81.	Elm-tree, raphides in, 47.
Chlorophylle, 34.	Enchondroma, 163.
circulation of, 35.	Equisetum, silica in, 49.
Chorda dorsalis, structure of, 136.	spores of, 73.
of Lamprey, 137.	Euphorbia, starch of, 32.
of Trout, 138.	
Classification of animal tissues,	F.
117.	Fat, 181.
Clematis, ducts of, 107.	eomposition of, 186.
Cocoa-nut, oil of, 38.	of Binda 185
—— shell of, 68.	of Birds, 185.
Codfish, fat of, 183.	of Ced feb 183
Collomia grandiflora, cells in seed	of Typests 183.
of, 14.	of Insects, 182.
Composition of fat, 186.	of Turtle 185
Confervæ, spores of, 70.	— of Turtle, 185.
—— conjugation of, 72.	Fatty degeneration, 193.
Cornea, fatty degeneration of, 200.	of somes 200
Croton-oil, seed of, 67.	of cornea, 200.
Cuticle of Aloc, 13.	of muselc, 196.
of Mallow, 56.	of Fowl, 198.
siliceous, 49.	of Ostrich, 198.
pigment in, 207.	Ferns, vessels in, 101.
Cuttlefish, cartilage of, 139.	Festuca pratensis, silica of, 55.
Cycas revoluta, woody tissue of,	Fibre, 13.
91.	of Campanula, 13.
hairs of, 100.	—— of Digitalis, 13.
210120 02, 2000	Fibro-cartilage, 177. ——————————————————————————————————
D.	of vertebræ, 170.
Desmidicæ, 57.	Fibro-cellular tissue, 75.
Deutzia, silica of, 55.	of Pleurothallis, 75
Development of arcolar tissue, 129.	of Orchidaceæ, 76.
Diatomaccæ, 56.	of Seeds, 77.
Digitalis, fibre of, 13.	Fibrous tissue, 120.
Discased cartilage, 154.	of membrana putaminis,
— potatocs, 33.	123.
Dotted ducts, 105.	vessels of, 132.
— tissue, 78.	Figure election 36
Dragon-palm, 79.	Fishes, articular cartilage of, 146.
Ducts, dotted, 105.	— eartilage of, 171.
—— porous, 105.	Fixed oils, 37.
porous, 2000	Flax, 80.
E.	woody tissue of, 80.
Ton of Mouga 110 127	— New Zealand, 82,
Ear of Mouse, 119, 137.	preparation of, 87.
Bat, 119, 137.	Fowl, latty degeneration of muscle
Eel, muscle of, 194.	of, 198.
Elaters, 73. — of Junjermannia, 13.	Fruit, stones of, 68.
- or ounjoinming, ro.	,,

G.

H.

Hairs of Anchusa Italica, 12. Hard tissues, 61. Hare, absence of fat in, 188. Hiccory, raphides in, 45. Histology, definition of, 1.

I.

Indian rubber, 36.
Insects, fat of, 182.
Ivory like deposit of bone, 160.
Ivory nut, structure of, 63.

J.

Jatropa oil, 39. Jungermannia, elaters of, 13.

\mathbf{K} .

Kieselguhr, 59.

L.

Lacunæ, 24.
Lamprey, cartilage of, 136.
— pigment of, 204.
Leck, cells of, 23.
— flower, stem of, 7.
Lepidosteus, scale of, 175.
Lettuce, vessels of, 98.
Lime-tree, raphides in, 45.

M.

Mallow, cuticle of, 56.

Mammalia, cartilage of, 140.

— articular cartilage of, 147.

Marchantia, root of, 9.

Mclanosis, pigment of, 209.

Membrane, 6.

Membrane, animal, 117. —— pupillaris, vessels of, 130. — vessels of, 130. —— synovial, 148, 151, 161. —— unrolling spirally, 100. - vessels of, 150. Milky juiccs, 35. Mineral oil, 40. Mollities ossium, 175. Mucor, 18, 22. Mummy cloth, 83. Muscle, capillaries of, 194. --- of Eel, 194. fatty degeneration of, 196. of Fowl, 198. —— of Ostrich, 198. —— of Sheep, 198. —— structure of, 193.

N.

— and tendon, vessels of, 150.

Naked capillaries, 134. Negro, pigment of, 203. Nettle, sting of, 69. Newt, pigment of, 204. New Zealand Flax, 82. Nitella, 35.

Mushroom, cells of, 25.

Ō.

Oat, silica of, 54. Oils, fixed, 37. Oil, Cocoa-nut, 38. —— Jatropha, 39. —— mineral, 40. —— Palm, 38. —— volatile, 40. Onion, raphides in, 44. Opuntia vulgaris, cells of, 15. Orchidaceæ, fibro cellular tissue of, 76. woody tissue of, 90. Osscous granules, 173. — in pus, 174. Ossifying cartilage, 168, 172. Ostrich, fatty muscle of, 198. tendon of, 132.

P.

Palm oil, 38. Pear, gritty tissue of, 61.

Pelargonium, 26.	Riec-paper, starch of, 29.
Phones wistota illiant 5	Root of Marchantia, 9.
Pharus, eristatus silica of, 54.	Rush, stellate cells of, 25.
Phytozoa, 70. — in Chara, 70.	S.
in Mosses, 71.	Salvia pratensis, seeds of, 14.
Pig, fat of, 185.	Sarcina ventriculi, 19.
— muscle of, 195.	Scale of Lepidosteus, 175.
Pigment, 202.	Sealariform vessels, 101.
———— in the eye, 202.	Selerogen, 61.
skin, 203.	Seed of eroton-oil, 67.
———— Lamprey, 204.	—— of Gourd, 7.
——— Newt's, 204.	of Star anise, 65.
Plaice, 204.	— of Tamarind, 67.
absence of in Albinoes, 203.	Seeds, fibro-eellular tissue of, 77.
Pigmentum nigrum, 202, 205.	—— of Salvia pratensis, 14. Sepia, 209.
Pine-apple fibre, 82. Pinus Webbiana, 11.	Sheep, fatty degeneration of musele
Pith of Elder, 9.	of, 198.
Pleurothallis, cells of leaf of, 9.	— areolar tissue of, 129.
—— fibro cellular tissue of, 75.	Shell of Cocoa-nut, 68.
Porcellanous deposit on bone,	—— of Walnut, 68.
160.	Silica, 49.
Porous tissue, 78.	—— of Bamboo, 51.
—— duets, 105.	—— of Canes, 52.
Porpoisc, adipose tissue of, 191.	—— of Deutzia, 55.
Potatoes, disease of, 33.	of Oct 54
—— starch of, 29.	of Oat, 54. of Rice, 52.
Pus, osseous granules in, 174.	—— of Wheat, 52.
R.	Siliccous cutieles, 49.
Rana paradoxa, cartilage of, 171.	Skin, pigment of, 207.
Raphides, 42.	Snake-wood, structure of, 65.
in Apple-tree, 46.	Sparganium ramosum, cells of pe-
—— artificial, 47.	tiole of, 24.
—— in Caetus, 45.	Spermaecti, 187, 188.
—— in Elæagnus, 46.	Spermaecti-whale, fat of, 187.
—— in Elm, 46.	Sphagnum, vessels of, 8.
— in Hiccory, 45.	—— leaf of, 8.
in Lime-tree, 45.	Spiral vessels, 93, 95. ———————————————————————————————————
in Onion, 44.	——————————————————————————————————————
— in Squill, 43. Reptiles, adipose tissue of, 188.	Spores of Chara, 73.
eartilage of, 140, 146.	— of Confervæ, 70.
Resin, 41.	— of Equisetum, 73.
Rete mucosum, 203.	Squill, raphides in, 43.
Rind of Gourd, 24.	Star anise, seed of, 65.
Rhubarb, starch of, 28.	Starch, 28.
Riband-fish, vertebra of, 175.	of Arrow-root, 29.
Riec-paper, 79.	of Emphorbia 32
—— silica of, 52.	—— of Euphorbia, 32.

Starch of Potato, 29. — of Rhubarb, 28. — of Rice, 29. — of Sago, 29. — tests of, 30. — of Tous les mois, 29. — of Wheat, 29. Stellate cells, 24. — of Rush, 25. — of Sparganium, 24. Stigmariæ, vessels of, 106. Stings of Nettle, 69. Stipules of Ficus elastica, 36. Stones of fruit, 62, 68. Strawberry, cells of, 22. Structure of muscle, 193. Synovial membrane, 148, 151, 161. — vessels of, 150, 156.	V. Vaagmaer, vertebra of, 175. Vallisneria, 35. — circulation in, 35. Vascular tissue, 93. Vegetable tissues, 1. — conclusion of, 109. Vertebræ, fibro cartilage of, 180. Vessels of Balsam, 8. — of cartilage, 141. — of articular cartilage, 149, 151, 155. — of diseased cartilage, 154. — of Ferns, 101. — scalariform, 101. — spiral, 95. Vine, cells of, 9. Volatile oil, 40.
T.	w.
Tabasheer, 50. Tallow plant, 41. Tamarind, seed of, 67. Tendon, structure of, 120. —— vessels of, 132, 157. Tests of starch, 30. Tissue, adipose, 180. —— animal, 115. —— classification of, 117. —— areolar, 127. —— dotted, 78. —— fibro-cellular, 75. —— fibrous, 120. —— porous, 78. —— vascular, 93. —— voody, 80. —— yellow fibrous, 125. Tissues, hard, 61. Torula ccrcvisie, 18. Tous les mois, starch of, 29. Tumours, 163.	Walnut-shell, 68. Wax plant, 41. Wheat, silica of, 52. White animals, absence of pigment in, 206. Woody tissue, 80, 87. ————————————————————————————————————

- of Giraffe, 124.

Just published, 1 vol. 8vo. Price £1. 2s.

PRACTICAL TREATISE

ON THE USE OF THE MICROSCOPE.

BY JOHN QUEKETT,

ASSISTANT CONSERVATOR OF THE MUSEUM OF THE ROYAL COLLEGE OF SURGEONS OF ENGLAND.

SECOND EDITION, REVISED, WITH STEEL AND WOOD ENGRAVINGS.

Just ready, 1 vol. Price 6s.

THE PRESCRIBERS' COMPLETE HANDBOOK,

COMPRISING THE

PRINCIPLES OF THE ART OF PRESCRIBING,

A MATERIA MEDICA CONTAINING ALL THE PRINCIPAL MEDICINES EMPLOYED,

CLASSIFIED ACCORDING TO THEIR NATURAL FAMILIES, WITH THEIR PROPERTIES, PREPARATIONS AND USES, AND

A CONCISE SKETCH OF TOXICOLOGY.

BY M. TROUSSEAU,

PROFESSOR OF THE FACULTY OF MEDICINE, PARIS, AND

M. REVEIL.

EDITED, WITH NOTES,

BY J. BIRKBECK NEVINS, M.D.

LONDON:

HIPPOLYTE BAILLIERE, 219, REGENT STREET, AND 290, BROADWAY, NEW YORK, U.S.

STANDARD SCIENTIFIC WORKS,

PUBLISHED BY

HIPPOLYTE BAILLIERE,

219, REGENT STREET, LONDON;

ANI

290, BROADWAY, NEW YORK, U.S.

J. B. BAILLIERE, LIBRAIRE, RUE HAUTEFEUILLE, PARIS.

BAILLY BAILLIERE, LIBRAIRE, CALLE DEL PRINCIPE, MADRID.

Mr. Bailliere having lately opened a house at 290, Broadway, New York, for the sale of Scientific Works, begs respectfully to state that he shall now be enabled to supply any American Books on Science and General Literature, immediately after Publication, at the rate of Five Shillings the Dollar on the New York Prices.

H. B. continues to receive a weekly parcel from France, containing the newest Works on Science and General Literature, which he supplies at the rate of One Shilling per Franc on the advertised Price in Paris. He begs to acquaint his friends and the Patrons of German Scientific Works, that he is able to furnish German Works and Periodicals every Month, at the rate of Three Shillings and Sixpence the Rix-Dollar.

Chemistry, Physics, Mineralogy, Geology, Astronomy, Rural Economy, &c.

3, 355			
Blakey (R.) History of Logical Science from the Earliest Times to the Present Day, by Robert Blakey, Professor of Logic and Metaphysics, Queen's College, Belfast, Author	£	S	d
of the Mistory of the Philosophia of Mind, in I vol damy sug	0	12	λ Δ
teorology, By J. B. Boussingault Member of the Institute of F.	U		0
with Notes, carefully revised and corrected, I vol. 8vo. cloth boards. London, 1845. Brewster (Sir David). The Natural History of Creation, in I vol. royal 8vo. Illustrated with Engravings and Woodcuts. In preparation.	0	18	0
Campbell. A Practical Text-Book of Inorganic Chemistry Including the Preparations of Substances, and their Qualitative and Quantitative Analyses, with Organic Analyses. By Dugald Campbell, Demonstrator of Practical Chemistry to the University College. 12mo. London, 1849			
Chapman. A Brief Description of the Characters of Minerals; forming a familiar Introduction to the Science of Mineralogy. By Edward J. Chapman. 1 vol. 12mo. with	0	5	6
Practical Mineralogy; or, a Compendium of the distinguishing Characters of Minerals; by which the Name of any Species or Variety in the Mineral Kingdom may be speedily ascertained. By Edward J. Chapman. 8vo. Illustrated with 13 engravings, showing 270 specimens. London, 1843	0	4	0
Chemical Society (Quarterly Journal of the). 4 vols. 8vo. London, 1848—52	0	7	()
Published Quarterly. Each Part	2	12	
Cook. Historical Notes on the Discovery and Progressive Improvements of the Steam Engine; with References and Descriptions to accompany the Plates of the American condensing Steam Engine for River Boats. 18mo. and a large fol. coloured plate on a roller and canvas. New York, 1849	0	3	0
Burnas and Ranssingarite in o	0	18	0
Dumas and Boussingault. The Chemical and Physiological Balance of Organic Nature: an Essay. By J. Dumas and J. B. Boussingault, Members of the Institute of France. 1 vol. 12mo. London, 1844			
M.D., with Additions, 8vo and 28 4to water 18(1) Philadelphis. Edited by R. Knox,	0	4	0
	1	4	
The Text separately with the four additional Plates, for P	2	2	0
the French edition. Plain			
Coloured .		12	
	U.	1.4	11

70	£	8	d
Reverend John R. Morrell, with Critical Annotations, a Biography of Fourier, and a General Introduction, by Hugh Doherty. 2 vols. 8vo. London 1851	1	1	0
History of Drugs; their Effects, Doses, and Adulterations, with the Preparations of the British Pharmacopæias. Svo. London, 1851	0	12	0
Cordon (L.) A Synopsis of Lectures on Civil Englineering and Mechanics. 4to. London, 1849	0	7	6
Gordon and Liddell. Exposition of a Plan for the Metropolitan Water Supply, showing that the Thames at Maple-Durham is the most eligible source from which a supply of pure soft water can be brought for the inhabitants of London and its suburbs. 8vo. London, 1849	0	1	0
Graham. Elements of Chemistry; including the Application of the Science in the Arts. By T. Graham, F.R.S. L. & E., Professor of Chemistry at University College, London. 2nd Edition, entirely revised and greatly enlarged, copiously illustrated with Woodcuts. Vol. 1, 1850	1		0 0
Humboldt. Kosmos: a General Survey of the Physical Phenomena of the Universe. By Baron A. Humboldt. The original English Edition, 2 vols. post 8vo. London, 1848.	0	15	
Kentz. A Complete Course of Meteorology. By L. F. Kæmtz, Professor of Physics at the University of Halle. With Notes by Ch. Martins, and an Appendix by L. Lalanne. Translated, with Additions, by C. V. Walker. 1 vol. post 8vo. pp. 624, with 15 Plates, eloth boards. 1845	0	12	6
KNAPP. Chemical Technology, or Chemistry applied to the Arts and Manufactures. By F. Knapp, Professor at the University of Giessen. Edited, with numerous Additions, by Dr. E. Ronalds, Professor of Chemistry at Queen's College, Galway; and Dr. Thomas Richardson, of Newcastle-on-Tyne. Illustrated with 600 large Woodcuts and 7 Coloured Plates. 3 vols. 8vo. London, 1848—1851	3		
Vol. III. Containing Sugar, Tea, Flour, &c. With 7 Coloured Plates Knipc. Geological Map of the British Isles. Coloured. In a case. London, 1848	4		. 0
Leon (John A.) The Art of Manufacturing and Refining Sugar, including the Manufacture and Revivification of Animal Charcoal. With 14 Plates, illustrative of the Machinery and Building. Large folio. London, 1850	3	3	0
Licbig. Chemistry and Physics, in relation to Physiology and Pathology. By Baron Justus Liehig, Professor of Chemistry at the University of Giessen. 2nd Edition, 8vo. London, 1847	0	3	Q
Mcmoirs of the Literary and Philosophical Society of Man- chester. (Second Series). Vol. 9. 8vo. with Woodcuts and Plates. London, 1851.	0	7	6
Mitchell (J.) Manual of Practical Assaying, intended for the use of Metallurgists, Captains of Mines and Assayers in General. With a copious Table, for the purpose of ascertaining in Assays of Gold and Silver the precise amount, in Ounces, Penoyweights, and Graius, of noble Metal contained in one ton of Ore from a given quantity. 1 vol. post Svo. London, 1846 Treatise on the Adulterations of Food, and the Chemical Means employed to	0	10	G
detect them. Containing Water, Flour, Bread, Milk, Cream, Beer, Cider, Willes, Spirituous Liquors, Coffee, Tea, Chocolate, Sugar, Honey, Lozenges, Cheese, Vinegar, Pickles, Anchovy Sauce and Paste, Catsup, Olive (Salad) Oil, Pepper, Mustard. 1200. London, 1848	0	6	0
Miller. Principles of Physics and Meteorology. By J. Muller, M.D. Illustrated with 530 Woodcuts, and 2 coloured Plates, 8vo. London, 1847	0	18	0
Nichol. Astronomy Historically and Scientifically developed, showing the Rise of the Science from its Growth, and the Character of the illustrions Men who have contributed to it. By J. P. Nichol, Professor of Astronomy in the University of Glasgow. 2 vois.			
The Architecture of the Heavens. Ninth Edition, entirely revised and greatly enlarged. Illustrated with 23 Steel Engravings and numerous Woodcuts. 8vo. London, 1851	0	16	0
Quckett (J.) Lectures on Histology, delivered at the Royal College of Surgeons of	0	10	6
Practical Treatise on the Use of the Microscope. Indistracted With 12 Section Loudon 1859.	1	2	0
Diseased Structure. 8vo. Illustrated with Engraved Plates and Woodcuts. In the			
Recid. Rudiments of Chemistry, with Illustrations of the Chemistry of daily Life, by D. B. Reid, M.D., Lecturer on Chemistry formerly one of Her Majesty's Commissioners for the Health of Towns in England. 4th Edition, with 130 Woodcuts. 12mo. 1850.	0	2	6
Regnant. An Elementary Treatise on Crystallography, Mustrated with 100 wood Engreyings, printed on black ground. Svo. Loudon, 1848.	0	3	0
Reichenbach (Baron Charles). Physico-Physiological Researches on the Dynamics of Magnetism, Electricity, Heat, Light, Crystallization and Chemism, in their Relations to Vital Force. The complete Work from the German Second Edition, with Addltions, a Preface and Critical Notes, by John Ashburnen, M.D. 8vo. With Woodcuts and One Plate. London, 1850.	0	15	0
		_	-

Richardson. Geology for Beginners; comprising a Finmiliar Exposition of the Elements of Geology and its Associate Sciences, Mineralogy, Fossil Conchology, Fossil Botany and Paleontology. By G. F. Richardson, F.G.S. 2nd Edition, post 8vo. With 251	3	e s	d
Woodcuts. 1843. Richardson and Ronalds. Metallurgy; and the Chemistry of the Metals. In	. 0) 8	0
3 vols. 8vo. Illustrated with numerous Wood Engravings. In the press. Stars and the Earth. The Stars and the Earth; or, Thoughts upon Space, Time, and Eternity. 4th Edition, Eighth thousand, 2 Parts in 1, 18mo. London, 1850.	. 0) 2	3 0
Stockhardt. The Principles of Chemistry, illustrated by Simple Experiments. Translated from the German Edition. With 218 Woodcuts. 12mo. Cambridge, 1850.	. 0	7	6
Thomson. Chemistry of Organic Bodies-Vegetables. By Thomas Thomson, M.D. F.R.S. L. & E, Regius Professor of Chemistry in the University of Glasgow. Corresponding Member of the Royal Academy of Paris. 1 large vol. 8vo. pp. 1092, boards.			
London, 1838			0
Chemistry of Animal Bodies. Svo. Edinburgh, 1843 Thomson (R. D.) British Annual and Epitome of the Progress of Science. By R. D. Thomson, Assistant Professor in the University of Glasgow. 3 vols. 1837, 38, 39, 18mo.	0	15	
cloth boards, lettered, each Weisbach (J.) Principles of the Mechanics of Machinery and Engineering. 2 vols. 8vo. Illustrated with 200 Wood Engravings. London 1848.	. 0	19	6
		10	U
			=
Anatomy, Medicine, Surgery, and Natural Histor	у.		
Ashburner. On Dentition and some Coincident Disorders. 18mo. London, 1834 Canton (A.) The Teeth and their Preservation, in Infancy and Manhood to Old Age. 12mo. with Woodcuts			0
Courtenay. Pathology and Rational Treatment of Stricture and Urethra in all its Varieties and Complications, with Observations on the Use and Abuse of Urethral	0	4	0
Instruments The whole illustrated by numerous Cases. By F. B Courtenay, M.R.C.S. &c. 4th Edition, 8vo. London, 1848 A Few Words on Perineal Section, as recommended by Professor Syme, for the	0	5	0
Practical Observations on the Chronic Enlargement of the Prostrate Clandin Old	0	1	0
People: with Mode of Treatment. By Francis B. Courtenay. Svo. with numerous Cases and Plates, boards. London, 1839 A Treatise on the Cure of Stricture of the Urethra; with Practical Observations on	0	7	G
the Treatment of Spermatorrhœa by Cauterization. 8vo. 1851 Crnveilhier and Bonamy. Atlas of the Descriptive Anatomy of the Human Body. By J. Cruveilhier, Professor of Anatomy to the Faculty of Medicine, Paris. With Explanations by C. Bonamy.	0	3	6
Explanations by C. Bonamy. Containing 82 Plates of Osteology, Syndemology and Myology. 4to. London, 1844. Plain Coloured	3	0	0
Fau. The Anatomy of the External Forms of Man (for Artists). Edited by R. Knox, M.D. with Additions. 8vo. Text, and 28 4to, Plates. London, 1849. Plain	5	15	
Gerber and Gulliver. Elements of the General and Microscopical Anatomy of Man and the Mammalia; chiefly after Original Researches. By Professor Gerber. To which Is added an Appendix, comprising Researches on the Anatomy of the Blood, Chyle, Lymph, Thymous Fluid, Tubercle and Additions had Gully	2	2	
Grant. General View of the Distribution of Extinct Animals. By Robert E. Grant, M.D., F.R.S. L. & E., Professor of Comparative Animals.	1	4	0
In the "British Annual," 1839. 18mo. London, 1839 On the Principles of Classification, as applled to the Primary Divisions of the Animal Kingdom. In the "British Annual," 1838. 18mo. Illustrated with 28 Woodcuts. London, 1838	0	3 (6
Outlines of Comparative Anatomy. Svo. Illustrated with 148 Woodcuts, boards.	0	3 (
Part VII. with Title-page Hall (Marshall). On the Diseases and Derangements of the Nervous System, in their Primary Forms, and in their modifications by Aga, Say, Carolina (System), in their	0	8 (
disposition, Excesses, General Disorder and Organic Disease. By Marshall Hall, M.D., F.R.S. L. & E. 8vo. with 8 engraved Plates. London, 1841	0 1	15 ()
The following is an Appendix to the above Work. On the Mutual Relations between Anatomy, Physiology, Pathology, Therapentics and the Practice of Medicine, helps, the Gulstonian Latin			
2 Coloured Plates and 1 Plates London 10 Constollar Bectures for 1842. Svo. with	0	5 (
New Memoir on the Nervous System, true Spinal Marrow, and its Anatomy, Physiology, Pathology, and Therapeutics. 4to. with 5 engraved Plates. London, 1843		0 (
Henriques. Etiological, Pathological and Therapeutical Reflections on the Aslatic Cholera, as observed in Europe, Asia Minor, and Egypt. 8vo. London, 1848		1 6	

Hufeland. Manual of the Practice of Medicine; the Result of Fifty Years' Experience By W. C. Hufeland, Physician to the late King of Prussia, Professor in the University of	£	3	d	
Berlin. Translated from the Sixth German Edition by C. BRUCHHAUSEN and R. NELSON. 8vo. bound. London, 1844	0	15	0	
Jones (W.) An Essay on some of the most Important Diseases of Women, with a Description of a Novel Invention for their Treatment and Relief. Second Edition. 8vo. London, 1850				
Practical Observations on the Diseases of Women, showing the necessity of Physical Examination, and the Use and Application of the Speculum. Illustrated by Cases.	0	1	0	
Woodcuts, and Coloured Plates. 8vo. London, 1850 Lebaudy. The Anatomy of the Regions interested in the Surgical Operations performed	0	7	ŋ	
upon the Human Body; with Occasional Views of the Pathological Condition, which render the interference of the Surgeon necessary. In a Series of 24 plates, the Size of Life. By J. Lebaudy. Folio. London, 1845	1	4	0	
Lec. The Anatomy of the Nerves of the Uterus. By Robert Lee, M.D., F.R.S. Folio, with 2 engraved Plates. London, 1841	0	8	0	
Mattlock. Practical Observations on the Efficacy of Medicated Inhalations in the Treatment of Pulmonary Consumption. By Dr. Maddock. 3rd Edition. Svo. with a coloured Plate. London, 1846	0		6	
Martin. A General Introduction to the Natural History of Mammiferous Animals: with a particular View of the Physical History of Man, and the more closely allied Genera of	V	v	,	
the Order "Quadrumana," or Monkeys. Illustrated with 296 Anatomical, Osteological, and other Engravings on Wood, and 12 full-plate Representations of Animals, drawn by W. Harvey. 1 vol. 8vo. London, 1841	0	16	0	
Moreau (Professor). Icones Obstetricæ; a Series of 60 Plates and Text, Illustrative of the Art and Science of Midwifery in all its Branches. By Moreau, Professor of Midwifery to the Faculty of Medicine, Paris. Edited, with Practical Remarks, by J. S.				
Strecter, M.R.C.S. Folio. Cloth boards. London, 1841. Price Plain	3 6	3 6	0	
Owen. Odontography; or, a Treatise on the Comparative Anatomy of the Teeth, their Physiological Relations, Mode of Development, and Microscopical Structure in the Vertebrate Animals. By Richard Owen, F.R.S., Corresponding Member of the Royal Academy of Sciences, Paris and Berlin; Hunterian Professor to the Royal College of				
Surgeons, London. This splendld Work is now completed. 2 vols. royal 8vo. containing 168 plates, half-bound russia. London, 1840-45.		0		
A few copies of the Plates have been printed on India paper, 2 vols. 4to Phillips. Scrofula: its Nature, Prevalence, Canses, and the Principles of Treatment. By Benjamin Phillips, F.R.S., Surgeon and Lecturer on Surgery to the Westminster	10			
Hospital. 8vo. with an engraved Plate. London, 1846 A Treatise on the Urethra; its Diseases, especially Stricture, and their Cure. 8vo.		12		
boards. London, 1832. Prescriber's (The) Complete Handbook. See Trousseau.	0	0	3 0	
Prichard. The Natural History of Man; comprising Inquiries into the Modifying Influence of Physical and Moral Agencies on the different Tribes of the Human Family.				
By James Cowles Prichard, M.D., F.R.S., M.R.I.A. Corresponding Member of the National Institute of the Royal Academy of Medicine, and of the Statistical Society				
of France; Member of the American Philosophical Society, &c. &c. 3rd Edition, enlarged, with 50 coloured and 5 plain Illustrations, engraved on Steel, and 97 Engray-		10	3 0	
lngs on Wood, royal 8vo. elegantly bound in cloth. London, 1848 Appending to the First and Second Editions of the Natural History of Man, large				
8vo. with 6 coloured Plates. London, 1845 & 1848. Each .	,		3 (
the Researches into the Physical History of Mankind, folio, coloured, and 1 sheet of letter-press, in cloth boards. 2nd Edition. London, 1850. Illustrations to the Researches into the Physical History of Mankind. Atlas of 4-			4 ()
coloured and 5 plain Plates, engraved on Steel, large 8vo. Boards. London, 1841 On the Different Forms of Insanity, in Relation to Jurisprudence. (Dedicated to	·	1	8 ()
the Lord Chancellor of England.) 12mo. London, 1842	, 0) ;	5 ()
M.D., Physician to the Hopital de la Charlet. Talishted by La Charlet and La Char			o (0
400 varieties of Cutaneous Affections. London, 1889 The Text separately, 8vo. In boards.	. I	l :		0
Account of the Diseases of the Genito-Urinary Organs, with the Physiology of Generation in the Vegetable and Animal Kingdoms. By M. Ryan, M.D. 4th Edition, greatly	· ()	6	0
of the British Sp-cles of Burrowing Sand Wasps contained in all the Metropolitan Collections; with their habits, as far as they have been observed. 8vo. with 4 Plates		0 1	0	0
London, 1837			0	0
Elements of British Entomology. Part 1. 1839. 8vo.	. ()	8	U
Street (v. 15.) Oct Monthly			_	

	P	8	d
Tronssean and Reveil. The Prescribers Complete Handbook, comprising the Principles of the Art of Prescribing, with a List of Diseases and their appropriate Remedies, a Materia Medica containing all the Principal Medicines employed, classified according to their Natural Families, with their Properties, Preparations and Uses, and a Concise Sketch of Toxicology. By Troussean, Professor of the Faculty of Med., Paris, and M. Reveil. Edited, with Notes, by J. BIRKBECK NEVINS, M.D. London, 1852. 12mo.	0		0
True and False Spermatorrhoea. 8vo. 1852 Vogel and Day. The Pathological Anatomy of the Human Body. By Julius Vogel, M.D. Translated from the German, with Additions, by George E. Day, M.D., Professor to the University of St. Andrews. Illustrated with upwards of 100 plain and coloured	0	5	0
Engravings, 8vo. cloth. London, 1847	0	18	0
gantly bound in cloth, coloured Plates	1	14	6
Vol. II. containing the Order Rodentia; or, Gnawing Mammalia: with 22 Illustrations, engraved on Steel, and Engravings on Wood, royal 8vo. elegantly bound in cloth, coloured Plates. London, 1848	1 1	14 9	60
The Natural History of Mammalia is intended to embrace an account of the structure and habits of all the known species of Quadrupeds, or Mammals; to which will be added, observations upon their geographical distribution and classification. Since the fossil and recent species illustrate each other, it is also inlended to include notices of the leading characters of the extinct species. The Genera, and many of the species, are illustrated by Engravings on Steel, and by Woodcats. The modifications observable in the structure of the skulls, teeth, feet, and other			
parts, are atmost entirety illustrated by Steel Engravings. Williams. Elements of Medicine: Morbid Poisons. By Robert Williams, M.D., Physician to St. Thomas's Hospital. 2 vols. 8vo. London, 1836—41	1	0	c
Willis. Illustrations of Cutaneous Disease: a Series of Delineations of the Affections of the Skin, in their more interesting and frequent forms; with a Practical Summary of their Symptoms, Diagnosis and Treatment, including appropriate Formulæ. By Robert Willis, M.D., Member of the Royal College of Physicians. The Drawings are after Nature, and Lithographed by Arch. Henning. These Illustrations are comprised in 94 Plates, folio. The Drawings are Originals, carefully coloured. Bound in cloth, lettered.	0	818	
London, 1843. On the Treatment of Stone in the Bladder by Medical and Mechanical Means. London, 1842.	6	5	
Botany.			
Babington. Primitiæ Floræ Sarnicæ; or, an Outline of the Flora of the Channel Islands of Jersey, Guernsey, Alderney, and Sark. 12mo. London, 1839. Fielding and Gardner. Sertum Plantarum; or, Drawings and Descriptions of Rare and undescribed Plants from the Author's Herbarium. By H. B. Fielding;	0	4	0
assisted by G. Gardner, Superintendent of the Royal Botanic Gardens, Ceylou. 8vo. London, 1844 Hooker Icones Plantarum. By Sir W. J. Hooker, Director of the Royal Botanic Gardens, Keylou, New Series, Vols. L. IV. contribution, 100 Press, Press, Press, Vols. L. IV. contribution, 100 Press, Pres	1	1	0
Gardens, Kew. New Series, Vols. I—IV, containing 100 Plates each with Explanations, Svo. cloth. London, 1842—1844. Each vol. Vol. IV. Part 2. London, 1848	1	8	
The Lordon Journal of Part 2. London, 1848	ō :		0
1842-47 Vols. I-VI, with 24 Plates each, boards.	6	0	
Now reduced to 20 Shillings each Vol. Notes on the Botany of the Antarctic Voyage, conducted by Captain James Clark Ross, R.N., F.R.S., in H.M.S. Erchus and Terrors, with Observations with The			
Now reduced to 20 Shillings each Vol. Now reduced to 20 Shillings each Vol. Notes on the Botany of the Antarctic Voyage, conducted by Captain James Clark Ross, R.N., F.R.S., in H.M.S. Erebus and Terror; with Observations on the Tussac Grass of the Falkland Islands. 8vo. with 2 coloured Plates. London, 1843 Niger Flora; or, an Enumeration of the Plants of Western Tropical Africa, Collected by the late Dr. Th. Vogel, Botanist to the Voyage of the Expedition sent by Her Britannic Majesty to the River Niger in 1841, under the Command of Capt. H. D. Trotter, R.N., Including Spicilegia Gorgonea, by P. B. Webb, and Flora Nigritiana, by Dr. J. D. Hooker and George Bentham. With 2 Views a Man, and 50 Plates.		4	0
The London Johrnal of Botany. Vols. I—VI, with 24 Plates each, boards. Now reduced to 20 Shillings each Vol. Notes on the Botany of the Antarctic Voyage, conducted by Captain James Clark Ross, R.N., F.R.S., in H.M.S. Erebus and Terror; with Observations on the Tussac Grass of the Falkland Islands. Svo. with 2 coloured Plates. London, 1843 Niger Flora; or, an Enumeration of the Plants of Western Tropical Africa, Collected by the late Dr. Th. Vogel, Botanist to the Voyage of the Expedition sent by Her Britannic Majesty to the River Niger in 1841, under the Command of Capt. H. D. Trotter, R.N., including Spicilegia Gorgonea, by P. B. Webb, and Flora Nigritiana, by Dr. J. D. Hooker and George Bentham. With 2 Views, a Map, and 50 Plates. Svo. Mather (W.) Outlines of Botany. Part I, with 7 Plates, 12mo, cieth boards. London		0	0
The London Johrnal of Botany. Vols. I—VI, with 24 Plates each, boards. Now reduced to 20 Shillings each Vol. Notes on the Botany of the Antarcic Voyage, conducted by Captain James Clark Ross, R.N., F.R.S., in H.M.S. Erebus and Terror; with Observations on the Tussac Grass of the Falkland Islands. Svo. with 2 coloured Plates. London, 1843 Niger Flora; or, an Enumeration of the Plants of Western Tropical Africa, Collected by the late Dr. Th. Vogel, Botanist to the Voyage of the Expedition sent by Her Britannic Majesty to the River Niger in 1841, under the Command of Capt. H. D. Trotter, R.N., including Spicilegia Gorgonea, by P. B. Webb, and Flora Nigritiana, by Dr. J. D. Hooker and George Bentham. With 2 Views, a Map, and 50 Plates. Svo. London, 1849 Mather (W.) Outlines of Botany. Part I, with 7 Plates, 12mo. cloth boards. London, 1848.		4	0
The London Johrnal of Botany. Vols. I—VI, with 24 Plates each, boards. Now reduced to 20 Shillings each Vol. Notes on the Botany of the Antarcic Voyage, conducted by Captain James Clark Ross, R.N., F.R.S., in H.M.S. Erebus and Terror; with Observations on the Tussac Grass of the Falkland Islands. Svo. with 2 coloured Plates. London, 1843 Niger Flora; or, an Enumeration of the Plants of Western Tropical Africa, Collected by the late Dr. Th. Vogel, Botanist to the Voyage of the Expedition sent by Her Britannic Majesty to the River Niger in 1841, under the Command of Capt. H. D. Trotter, R.N., including Spicilegia Gorgonea, by P. B. Webb, and Flora Nigritiana, by Dr. J. D. Hooker and George Bentham. With 2 Views, a Map, and 50 Plates. Svo. London, 1849 Mather (W.) Outlines of Botany. Part I, with 7 Plates, 12mo. cloth boards. London, 1848. Miers (J.) Illustrations of South American Plants, Vol. I. 4to. With 34 Plates. London, 1847—50 Schieden. The Plant; a Biography, in a Series of Popular Lectures on Botany. Edited and Translated by A. Henfrey, F.L.S. 8vo. with 5 coloured Plates, and 13 Woodcuts. London, 1848	6 0 1 0	1 (0

Wight. Illustrations of Indian Botany; or, Figures Illustrative of each of the Natural Orders of Indian Plants, described in the Author's Prodromus Flore Peninsulæ Indiæ Orlentalis; but not confined to them. By Dr. R. Wight, F.L.S., Surgeon to the Madras Establishment. Vol. I, published in 13 Parts, containing 95 coloured Plates.	£	S	d
Matras, 1888—40	4	17	G
Vol. 11. 3 Parts, containing 200 coloured Plates. Madras, 1841-50 Odd Parts may be obtained to complete Sets.		12	
	A	0	0
- Vol. II, consisting of 4 Parts, containing together 418 Plates. Madras,	-1 ,	·	U
1840-42 Vol. III, Parts 1 to 4, with 426 Plates. Madras, 1843-47	5	5	
Vol. IV, Parts I to 4, with 458 Plates. Madras, 1848—50 Vol. V, Part I, with 140 Plates. Madras, 1851 Odd Parts may be obtained to complete Sets.		0	
Contributions to the Botany of India. By Dr. Robert Wight, F.L.S., Surgeon to		~	^
the Madras Establishment. 8vo. London, 1834 ———————————————————————————————————	0	7	łj
Notices of their Economical Properties and Uses. By Dr. Robert Wight, F.L.S., Surgeon to the Madras Establishment. 3 Parts, 4to. with 150 coloured Plates. Madras, 1846—48	1	10	0
Prodromus Floræ Peninsulæ Indiæ Orientalis; containing abridged Descriptions of the Plants found in the Peninsula of British India, arranged according to the Natural System, By Drs. Robert Wight, F.L.S., and Walker Arnott. Vol. I, 8vo.	-1	10	
London, 1834	0	16	0
	-	_	=
- Hom copathic.	-		
Bellnomini (J., M.D.) Scarlatina; its Treatment Homoopathically. 8vo. London,			
1843	0	1	0
Beenninghausen. Manual of Homeopathic Therapeutics, intended as a Guide to the Study of Materia Medica Pura. Translated, with Additions, by J. Laurie, M.D. 8vo. 1848	0	12	0
Essay ou the Homoopathic Treatment of Intermittent Fevers. 8vo. New York,	0	2	6
Black. A Treatise on the Principles of Homocopathy. Svo. London, 1842	0	9	0
Caspari. Homosopathic Domestic Physician, Edited and enlarged by F. Hartmann, with Additions and Preface by C. Hering, M.D., enriched by a Treatise on Anatomy and Physiology. Illustrated with Woodcuts, by W. Esrey, M.D. 1 vol. 12mo. Phlla-	0	8	. 0
delphia, 1852 Carie (P. F., M. D.) Practice of Homeopathy. 1 vol. 8vo. London, 1838		6	
Principles of Homeonathy, 1 vol. 8vo. London, 1837 · · ·		$\frac{5}{12}$	
Jahr's Homœopathy. New Edition, 2 vols. 12mo. London, 1847. See Jahr. Domestic Practice of Homœopathy. 3rd Editiou, 1850	0		0
Education The Pathogenetic Cyclopedia, a Systematic Arrangement and Analysis of the Hamconathic Materia Medica. Vol. I. Svo. Loudon, 1850	0	18	3 0
Bunsford (Harris). The Pathogenetic Effects of some of the Principal Homeopathic	0	(0 6
Remedies. 8vo. London, 1838. The Practical Advantages of Homeopathy, illustrated by numerous Cases. Dedi-	0		3 0
cated, by Permission, to Her Majesty Queen Adelaide. 1 vol. 8vo. boards. 1841 Epps. Domestic Homœopathy; or, Rules for the Domestic Treatment of the Maladies of Infants, Children, and Adults, and for the Conduct and Treatment during Pregnancy,	0		
Confinement and Suchling Ath Edition, 12mo, 20muon, 1044	0	1 4	1 6
Everest (T. H.) A Popular View of Homocopathy; exhibiting the Present State of		(0
- A Letter addressed to the Medical Practitioners of Great British on the Subject of	()) ;	1 6
Homeopathy. 8vo. London, 1834 Gunther. New Manual of Homeopathic Veterinary Medicine; or the Homeopathic Treatment of the Horse, the Ox, the Dog, and other Domestic Animals. Translated			
from the 3rd German Edition, with considerable Additions and Improvements	0	10	0 6
Hahnemann. Lesser Writings. Collected and Translated by R. E. Didgeon, M. D.			1 0
8vo. 1851 — Muterla Medlea Pura. Translated and Edited by Charles J. Hempel, M.D. 4 vols.	1	1:	2 0
Svo. New York, 1846 Materia Medica Pura. Purt 1. 4to. By Drysdule, Dudgeon and Black. London,	0		7 0
1852. Fublished by the Hahnemann Soviety Organon of Homespathic Medicine. Edited by Dr. Dudgeon. 8vo. London,			
1849 Organon of Homospathic Medicine. Edited by Dr. Brageon. Trans-	C) 10	0 0
The Chronic Diseases, their Specific Nature and Homocopathic Treatment. Translated and Edited by Charles J. Hempel, M.D. 5 vols. 12mo. New York, 1846	2	? (0 0

	£	S	d
Hamilton. A Guide to the Practice of Homoopathy. Translated and compiled in Alphabetical Order, from the German of Ruoff, Haas, and Rückert, with Additions.	0	5	0
12mo. 1844 The Flora Homœoputhica; or Illustrations and Descriptions of the Medicinal Plants used as Homœopathic Remedics. 8vo. with 3 Coloured Plates. 1851. Parts 1 to 8.	_		
Each. (A Part appears Monthly) Market 1 (F. Richard Day). Popular Outlines of Homocopathy. 24mo. London, 1840.	0		0
HRPEMANN (F.) Theory of Acute Diseases and their Homogopathic Treatment. Translated by C. Hempel. 2 vols. 12mo. New York, 1848 Theory of Chronic Diseases and their Homogpathic Treatment. 2 vols. 12mo. New	0	18	0
York, 1849	0	18	
Hayle. An Address on the Homocopathic System of Medicine. 8vo. 1843	0	1	0
Mempiel. Homocopathic Domestic Physician. Svo. New York, 1846 Henriques. The Homocopathic Medical Dictionary, and Home Guide. 12mo. 1851.	0	5 4	0
Hering (of Philadelphia). The Homeopathist; or, Domestic Physician. 3rd Edition, much enlarged. 12mo. London, 1852	0	10	
Momocopathic Examiner (The). By Drs. Gray and Hempel. New Series. Vols. I. and Il. New York, 1846-1847. Each	1	0	0
Homocopathic Handbook, and Clinical Guide for the Treatment of Diseases. Translated from the German by D. Spillan, M.D. 12mo. 1851	0	12	0
Homocopathic Pharmacopoeia and Posology (New); or the Mode of Preparing Homocopathic Medicines, and the Administration of Doses. From the Work of Buchner and Gruner. 12mo. 1850.	0	8	0
Jahr. Manual of Homocopathic Medicine. In 2 Parts.—Part I. Materia Medica. —Part II. Therapeutical and Symptomatological Repository. Translated from the 4th Edition, and Edited with Additions, by P. F. Curie, M.D. 2 vols. 8vo. London,			
1847	1	12	0
The American Edition. Translated with extensive additions from various sources,			
by C. J. Hempel, M.D., and J. M. Quin, M.D., the Materia Medica only. 2 large vols. 8vo. New York, 1848	3	3	0
This Work is intended to facilitate a comparison of the parallel symptoms of the various Homæopathic agents, thereby enabling the Practitioner to discover the characteristic symptoms of each drug, and to determine with ease and correctness the remedy.			
——————————————————————————————————————	0	2	G-
and Repertory for the Treatment of Acute and Chronic Diseases. Translated by C. J. Hempel. Revised and Edited, by J. Laurie. 12mo. London, 1850	0	12	0
Lanrie (J., M.D.) An Epitome of Homocopathic Domestic Medicine. 12mo. 5th Edition. London, 1851	^	5	^
Elements of Homœopathic Practice of Physic. 8vo. 1847		5 16	0
Homœopathic Domestic Medicine. 6th Edition. 8vo. London, 1851. The Parent's Guide, a Treatise on the Method of Rearing Children from their Infancy; Comprising the essential Branches of Moral and Physical Education. 12mo.	0	16	0
North American Homeopathic Journal (The). Conducted by Drs. Hering,	0	6	0
Marcy, Metcalf. New York, 1851 ———————————————————————————————————		18	
Malan. Pocket Book of Homcopathy. 18mo. Second Editlon, 1852.	0	18	0
Marcy. The Homeopathic Theory and Practice of Medicine. Svo. New York, 1850 Marsden. Notes on Homeopathy. Svo. London, 1849		12	
Newman (George). Homeopathic Family Assistant. 2nd Edition 18mo 1847	0	3 5	6
A Concise Exposition of Homeopathy; its Principles and Practice. With an Appendix	0		6
Ran. Organon of the Specific Healing Art. Translated by C. Hempel. Svo. New York, 1847	0	8	
Resenstein (J. G.) The Comparative Merits of Allopathy, the old Medical Practice; and Homoopathy, the Reformed Medical Practice; particularly illustrated. 8vo. Montreal. 1846	0	8	
Rueckert. Therapeutics; or, Successful Homocopathic Cures, collected from the best Homocopathic Periodicals. 8vo. New York			
Russell. A Treatise on Epidemic Cholern, with n Map showing the Course of the Cholera from India to Britain. Svo. London, 1849		15	
Simpson (M.D.) Practical View of Homogopathy. Svo. London, 1836		10 10	
Transactions of the American Institute of Homocopathy. Svo. New York, 1846 Whitefield. Homocopathy. The True Art of Healing. 18mo. Brighton, 1845	0	6	()
Yeldam (S.) Homoopathy in Acute Diseases. Svo. 2nd Edition London 1850	0	0	6

Ollendorff.

Mesmerism. £ 8 d Ashburner (J.) Facts in Clairvoyance, with Observations on Mesmerism, and its Application to the Philosophy of Medicine, and to the Cure of Diseasc. 8vo. London, 1.0 Barth. The Principle of Health Transferable. 18mo. 2nd Edition. London, 1850 0 0 6 - A Manual of Mesmeric Practice, intended for the Instruction of Beginners. 2nd Edition, 12mo, London, 1850 3 0 Bannann. Curative Results of Medical Somnambulism, consisting of several authenticated Cases, including the Somnambule's own Case and Cure. 8vo. London, 1849 1 6 The Handbook of Mesmerism. 3rd Edition. 12mo. London, 1851 1 0 Capern (Thos.) The Fifty Cases. 12mo. The Mighty Curative Powers of Mesmerism proved in a Hundred and Deleuze. Practical Instruction in Animal Magnetism. Translated by T. C. HARTSHORN. 4th Edition. With Notes, and a Life, by Dr. Foissac. 12mo. London, 1850. 46 Early Magnetism, in its Higher Relations to Humanity; as veiled in the Poets and the Prophets. By $\Theta Y O \Sigma$ MAOO Σ . 8vo. cloth. London, 1846. 5 0 Elliotson. Numerous Cases of Surgical Operations without Pain in the Mesmeric State; with Remarks upon the Opposition of many Members of the Royal Medical and Chirurgical Society, and others, to the Reception of the inestimable hlessings of Mesmerism. By John Elliotson, M.D. Cantah, F.R.S. 8vo. London, 1843 A Fine Portrait of, Engraved on Stone. London, 1844 2 6 46 0 Somnolism and Psycheism, otherwise Vital Magnetism or Mesmerism. 3rd Edition. 12mo. London, 1851 46 Jones. The Curative Power of Vital Magnetism; Verified by Actual Application to numerous Cases of Diseases. 12mo. London, 1845 1 0 Kistc. Mesmerism; or, Facts against Fallacies. In a Letter to the Rev. George Sandby. 18mo. London, 1845. Reichenbach. Physico-Physiological Researches on the Dynamics of Magnetism, Elec-1 0 tricity, Heat, Light, Crystallization, and Chemism, in their Relations to Vital Force, by Baron Charles Reichenbach. The Complete Work from the German second Edition, with Additions, Preface, and Critical Notes, by John Ashburner, M.D. 8vo. with Woodcuts, and I Plate. London, 1850 0 15 0 Sandby. Mesmerism and its Opponents. 2nd Edition, 12mo. 1848 5 0 A Practical Manual of Animal Magnetism; containing an Exposition of the Methods employed in producing the Magnetic Phenomena, with its Application to the Treatment and Cure of Diseases. By A. Teste, M.D. Translated from the 2nd Edition, by C. Spillan, M.D. Dedicated to John Elliotson, M.D. Cantab. 12mo. London, 1843. 6 0 Topham and Ward. Account of a Case of successful Amputation of the Thigh during the Mesmeric State, without the knowledge of the Patient. Read to the Royal Medical and Chirurgical Society on the 22nd of November, 1842. 8vo. 1.0 Facts in Mesmerism, with Reasons for a Dispassionate Inquiry into it. the Rev. Ch. H. Townsend. Second edition, with a New Preface, and enlarged. 8vo. 0 9 0 London, 1844 . The most Philosophical Work on the subject. ist. A Journal of Cerebral Physiology and Mesmerism, and their Application to Human Welfare. Published Quarterly, each Number *** This Journal contains papers by Drs. Elliotson, Engeldue, Ashhurner, &c. Zoist. 2 6 Thirty-six Numbers have already appeared. Nos. 1 to 36, forming 9 vols. 8vo. cloth hoards. Each vol. Vol. I., Commenced April 1, 1843, being nearly out of Print, is £1 10s. 0 11 0 Wilson. Practice of the Water Cure, with authenticated Evidence of its Efficacy and Safety. Part 1, containing 70 authenticated Cases, the Opinions of English Medical Practitioners, a Sketch of the History and Progress of the Water Cure, and an Account of the Processes used in the Treatment. Svo. London, 1844 The Water Cure, Stomach Complaints and Drugs, Diseases, their Causes, Consequences and Cure by Water, Air, Exercise and Diet. 8vo. Third edition, 1843 4 6 Bernstein (L.) Selections from the best German Authors in Prose and Poetry; also 6 0 some Commercial Letters, 12mo. London, 1842 niface. Modern English and French Conversation; containing Elementary Phrases and new Easy Dialogues, in French and English, on the most familiar Subjects: for the Use of the Traveller and Student. By M. Boniface. Sixteenth edition, 18mo. London, 3 0

9 0

0 4 6

lendorff. A new Method of Learning to Read, Write and Speak the German Language in Six Months. By H. G. Ollendorff. Translated from the Flfth French Edition. By G. J. Bertlinchamp, A.B. Fourth edition, revised and considerably improved. 12mo.

HIPPOLYTE BAILLIERE.

HAVING LATELY OPENED A HOUSE

290. BROADWAY, NEW YORK, U.S. AT

IS NOW ENABLED TO SUPPEY ALL AMERICAN BOOKS

AT THE RATE OF 5s. THE DOLLAR.

ALL ORDERS ENTRUSTED TO HIM WILL MEET WITH PROMPT ATTENTION.

H. B. makes up a case for New York every fortnight, and having Correspondence with most of the Scientific Men in the United States, would be glad to receive any communications for them at a small charge.

Allan. Philosophy of the Mechanics of Nature, and the Source and Modes of Action of Natural Motive-Power. 8vo. Illustrated with numerous Woodcuts. New York, 1852.

American Flora (The); or, History of Plants and Wild Flowers, over Six Thousand Plants.
By A. B. Strong. 4 vols. 4to. Coloured Plates. New York, 1848.

American Mechanics' Magazine and Engineers' Journal. Edited by W. Adams. January to December, 1851. Illustrated with Woodcuts. New York, 1852.

Astronomical Observations made during the years 1845—46 at the National Observatory, Washington. Published by authority. 2 vols. 4to. Washington, 1846 and 1851.

Bailey. Microscopical Examination of Soundings. 4to. With a Plate. Washington, 1851.

Microscopical Observations made in South Carolina, Georgia, and Florida. 4to. With 3 Plates. Washington, 1851.

Barry. The Fruit Garden; a Treatise. With 150 Woodcuts. 12mo. New York, 1851.

Bartol. A Treatise on the Marine Boilers of the United States. Svo. With numerous Woodcuts. Philadelphia, 1851.

Bell. Practical Methods of Ventilating Buildings. 8vo. With Plates. Boston, 1848.

Berzelius. The Use of the Blowpipe in Chemistry and Mineralogy. 8vo. With Plates. Boston, 1845.

Bowditch. The Young Stethoscopist; or, the Student's aid to Auscultation. 2nd Edition. 12mo. New York, 1848.

Campbell. Negro-Mania; being an Examination of the falsely assumed equality of the various Races of Men. 12mo. Philadelphia, 1851.

Coucs. Outlines of a System of Mechanical Philosophy; being a Research into the Laws of Force. 8vo. Boston, 1851.

Baguerreian Jourual (The). Devoted to the Daguerreian and Photogenic Arts. Vols. 1 and 2. 8vo. New York, 1850-51.
 Third Volume. (Now in course of Monthly Publication). 1852.
 Bana. A System of Mineralogy, comprising the most recent discoveries. 3rd Edition. 8vo. Illustrated with Diagrams. New York, 1850-00 that Devote Management Photogenic Arts. Vols. 1

Divis. The Great Harmonia; being a Philosophical Revelation of the Natural, Spiritual, and Celestial Universe. 2 vols. 12mo. Boston, 1851.

The Philosophy of Spiritual Intercourse, being an Explanation of Modern Mysterics. 8vo.

New York.

Dods. The Philosophy of Electrical Psychology. 12mo. New York. 1851.

Eastman. Dahcotah; or, Life and Legends of the Sionx around Fort Snelling. 12mo. With Plaics. New York, 1849.

Eberle. A Treatise of the Materia Medica and Therapeutics. 6th Edition, 2 vols. in 1. Svo.

Eberle A Treatise Philadelphia, 287.

Ellet. Physical Geography of the Misslssippl Valley. 4to. Washington, 1849.

Foster and Whitney. Report on the Geology and Topography of a Portion of Lake Superior Land District. 8vo. With 12 Plates and Map. Washington, 1850.

Fremont (Captain). Report of the Exploring Expedition to the Rocky Mountains in 1842, and to Oregon and North California in 1843-44. 8vo. With Maps and Plates. Washington, 1845—Oregon and California. The Exploring Expedition to the Rocky Mountains; to which is added a Description of the Physical Geography of California. With Notices of Gold Region. With Plates. 8vo. Buffalo, 1851.

French. Historical Callections of Louislana, embracing Translations of many rare and valuable Documents relating to the Natural, Civil and Political History of that State. 2 vols. 8vo. With Maps. New York and Philadelphin, 1846—50.

Gilson. Institutes and Practice of Surgery, being outlines of a Course of Lectures. Edition. With Plates. 2 vols. 8vo. Philadelphin, 1850.

Gray. The Botanical Text-Book, an Introduction to the Scientific Botany. Svo. With Woodcuts. 3rd Edition. New York. 1850.

Green. Observations on the Pathology of Cronp; with Remarks on its Treatment. 12mo. New York, 1849.

Grimes. Phreno-Geology: the Progressive Creation of Man, indicated by Natural History. 12mo. New York, 1850.

- Etherology and the Phreno-Philosophy of Mesmerism and Magic Eloquence. 12mo. New York, 1850.

Hare. Memoir on the Explosiveness of Nitre; with a view to Elucidate its Agency. 4to. Wash-

ington, 1849.

Harris. The Principles and Practice of Dental Surgery. 4th Edition. 8vo. With 200 Woodcuts. Hastings. The Practice of Surgery; embracing Minor Surgery, and the Application of Dressings. With numerous Illustrations. 12mo. Philadelphia, 1850.

Handbook of Young Artists and Amateurs in Oil Painting. By an American Artist. 12mo. New York, 1849.

Mannt. General Theory of Bridge Construction; with Practical Illustrations. 15 Plates. Svo. New York, 1851.

Hines. Oregon; its History, Condition, and Prospects; with Personal Adventures among the Indians. 8vo. Buffalo, 1852.
 Holgate. American Genealogy, being a History of some of the early Settlers of North America and their Descendants, from their first Emigration to the Present Time. 4to. New York, 1851.

Jay (J. C.) A Catalogue of the Shells, arranged according to the Lamarkian System, with their Authorities, Synonyms and References to Works where figured or described. 4th Editlon, 4to. New York, 1850.
 Johnson. The Coal Trade of British America. 8vo. Washington, 1850.
 Knapen. Mechanic's Assistant: a thorough Practical Treatise on Mensuration and the Sliding Rule. 12mo. New York, 1850.

Leconte. Descriptions of New Species of Coleoptera from California. 8vo.

Synopsis of the Cleridae of the United States. 8vo. New York, 1849.
On the Platycomy Compressus: a New Fossil, Pachyderm. 4to. New York, 1851.

With 4 Plates. New York, 1848.

Loomis. The Recent Progress of Astronomy. 12mo. New York, 1850.

LOPA. The Epoch of Creation. The Scripture Doctrine contrasted with the Geological Theory. 12mo. New York, 1851.

Macfarlanc. History of Propellers and Steam Navigation. 12mo. With Woodcuts. New York, 1851.

ury. Explanations and Sailing Directions to accompany the Wind and Current Charts, approved by Commodore Lewis Warrington, and published by Authority. 4to. With several large folio Maps. Washington, 1851. Maury.

Meigs (C.) Obstetrics: the Science and the Art. 8vo. With 121 Woodcuts. Philade A Practical Treatise on the Diseases of Children. 8vo. Philadelphia, 1848.

Women; her Diseases and Remedies. 2nd Edition. 8vo. Philadelphia, 1851.

Minific. Text Book of Geometrical Drawing for the Use of Mechanics and Schools. Philadelphia, 1849.

With Illustrations for Drawing Plans, Sections, and Elevations of Buildings and Machinery. Second Edition. With 56 Steel Plates. Royal 8vo. Baltimore, 1850. tratinns for

Mitchell. Materia Medica and Therapeutics: with ample Illustrations of Practice, and very copious Notices of Toxicology. 8vo. Philadelphia, 1850.

Morfit. Chemical and Pharmaceutical Manipulations: a Manual of the Mechanical and Chemico-Mechanical Operations of the Liboratory. With 423 Illustrations. 8vo. Philadelphia, 1849.

— Chemistry Applied to the Manufacture of Soap and Candles. 8vo. Philadelphia, 1847.

Neill and Smith. An Analytical Compendium of the various branches of Medical Sciences, Anatomy, Physiology, Surgery, Materia Medica, Chemistry and Practice of Medicinc. 5 vols. 8vo. with Woodcuts, bound in 1. Philadelphia, 1848.

Observations at the Magnetic and Meteorological Observatory at Girard College, Philadelphia, in 1840-45. 3 vols. 8vo. and Atlas of 200 Plates. Washington.

Overman. The Miniufacture of Iron in all its various Branches. 8vo. Philadelphin, 1850. Owen. Report on the Mineral Lands of the United States of America. Svo. with numerous Plates.

Washington, 1840.

- Report of a Geological Exploration of Part of Java, Wisconsin, and Illinois, with Charts

and Illustrations. Svo. Washington, 1848.

Report of a Geological Recomnaissance of the Chippewa land District of Wisconsin, with

Charts and Illustrations. Svo. Washington, 1848.

Reports to the Secretary of War with reconnaissance in New Mexico and Texas. Svo. with Charts and Plates. Washington, 1850.

Pharmaconcia of the United States of America (The). By Authority. 8vo. Philadelphia, 1851.

Putnam's Home Cyclopedia. Handbook of Useful Arts. 12mo. with Woodcuts. 670 pages. New York, 1852.

Handbook of Universal Biography. 12mo. 821 pages. New York, 1852. Handbook of Literature and the Fine Arts. 12mo. With Woodcuts. 648 pages. New York, 1852.

Robinson. Catalogue of American Minerals, with their Localities; including all which are known to exist in the United States and British Provinces. 8vo. Boston, 1825.

Ruchenbergen. Lexicon of Terms used in Natural History. 12mo. Philadelphia, 1850.

Saint John. Elements of Geology, intended for the Use of Students. 12mo. 180 Woodcuts. New York, 1851.

Schoolcraft. Personal Memoirs of a Residence of Thirty Years with the Indian Tribes on the American frontiers. 8vo. with a Portrait. Philadelphia, 1851.

Historical and Statistical Information respecting the History, Condition and Prospects of the Indian Tribes of the United States. Vol. I, 4to. with 76 coloured Plates. Published by authority of Congress. Philadelphia, 1851.

- The American Indians, their History, Condition and Prospects. New Edition. With Plates. Buffalo, 1851.

Smithsonian Contributions to Knowledge. 2 vols. 4to. With Maps and Plates. Washington, 1848-51.

Squier. Nicaragua, its People, Scenery, Monuments and the Proposed Interoceanic Canal. With numerous original Maps and Illustrations. 2 vols. Svo. New York, 1852.

Antiquities of the State of New York; being the result of extensive original Surveys and Explorations. 8vo. Buffalo. 1851.

Stille. Elements of General Pathology. 8vo. Philadelphia, 1848.

Taylor. Statistics of Coal, the Geographical and Geological Distribution of Mineral Combustibles or Fossil Fuel, including also Notices and Localities of the various Mineral Bitumiuous Substances employed in the Arts and Manufactures. Illustrated by Maps and Diagrams. Royal Svo. Philadelphia, 1848.

Trautwine. New Method of calculating the Cubic contents of Excavations and Embankments, by the aid of Diagrams. 8vo. Philadelphia, 1851.

The Field Practice of laying out circular Curves for Railroads. 12mo. Philadelphia, 1851.

Tucker. Elements of the Principles of Midwifery. 8vo. Illustrated with Woodcuts. Philadelphia, 1848.

Vestiges of Civilization; or, the Ætiology of History, Religious, Æsthetical, Political and Philosophical. 8vo. New York, 1851.

Von Amringe. An Investigation of the Theories of the Natural History of Man. By Lawrence Prichard and others, founded on Animal Analogics. 8vo. New York, 1848.

Walker. Researches relative to the Planet Neptune. 4to. Washington,

Wilkes. Narrative of the United States Exploring Expedition during the years 1838 to 1842.
With Illustrations and Maps. 5 vols. 8vo. Half bound. Philadelphia, 1845.

Wisc. System of Aeronautics, comprising its Earliest Investigations and Modern Practice and Art. 8vo. With Plates. Philadelphia, 1850.

Wheeler. Historical Sketches of North Carolina, from 1584 to 1851. Illustrated with Engravings. 2 vols. in 1. 8vo. Philadelphia. 1851.

Rural Homes; or, Sketches of Houses suited to American Country Life. 8vo. With Plates.

Wood. Treatise on the Practice of Medicine. 2nd Edition. 2 vols. 8vo. Philadelphia, 1849. Wood and Bache. The Dispensatory of the United States of America. 9th Edition. 8vo. Philadelphia, 1851.

Introduction to the Science of Government. With a Brief Treatise on Political Economy. 20th Edition. 8vo. Buffalo, 1851.

Troppaueger. German Grammar; with Reading Lessons, systematically arranged to show the affinity existing between the English and German Language and Progressive ક હૈ Exercises. Fourth edition, 12mo. London, 1849 0 6 6

Revue des Denx Mondes. Publié tous les 15 Jours en un Cahler de 10 à 12 feullles d'impression: - Souscription pour l'Année . 6 Mois 6 0 3 Mois 0 15 0 Un des meilleurs Journaux de Littérature publié à Paris.

Wellesley (W.) De la France Contemporaine et de ses divisions Hiérarchiques. Réponse à l'ouvrage de M. Guizot, de la "Démocratie en France," 8vo. Londres, 1849 0 6 0

VOL. X. OF THE

LIBRARY OF ILLUSTRATED

STANDARD SCIENTIFIC WORKS.

Already Published.

Τ.

Professor Muller's Principles of Physics and Meteorology. with 530 woodcuts and two coloured engravings. 8vo. 18s.

II. AND V.

Professor Weisbach's Mcchanies of Machinery and Engineering. 2 vols. with 900 woodcuts. £1 19s.

III. IV. AND X.

Professor Knapp's Technology; or,
Chemistry Applied to the Arts and Manufactures.

EDITED BY DR. RONALDS, AND DR. T. RICHARDSON,
3 vols. splendidly illustrated, price £3 4s.

VI.

Quekett's (John) Practical Treatise on the Use of the Microscope. second edition, with 12 steel and numerous wood engravings. 8vo. £1 2s.

VII.

Professor Fau's Anatomy of the External Forms of Man. FOR ARTISTS.

EDITED BY R. KNOX, M.D.

8vo. and an atlas of 28 plates 4to. Plain £1 4s. coloured £2 2s.

VIII.

Professor Graham's Elements of Chemistry, with Its Application in the Arts.

SECOND EDITION, WITH NUMEROUS WOODCUTS. VOL. I. £1 1s.

TX.

Professor Nichol's Architecture of the Heavens.

NINTH EDITION, WITH 23 STEEL PLATES AND MANY WOODCUTS. LONDON, 1851. 168.

In the Press.

A Complete Treatise on Metallurgy, and the Chemistry of the Metals.

BY DRS. RONALDS AND RICHARDSON.

WITH ILLUSTRATIONS.

CATALOGUE

OF

SCIENTIFIC BOOKS.

AMERICAN, FRENCH AND GERMAN.

HIPPOLYTE BAILLIÈRE,

219, REGENT STREET, LONDON;

AND

290, BROADWAY, NEW YORK, U.S.

Mr. Bailliere having lately opened a house at 290, Broadway, New York, for the sale of Scientific Works, begs respectfully to state that he shall now be enabled to supply any American Books on Science and General Literature, immediately after Publication, at the rate of Five Shillings the Dollar on the New York Prices.

H. B. makes up a case for New York every fortnight, and having Correspondence with most of the Scientific Men in the United States, would be glad to receive any communications for them at a small charge.

H. B. continues to receive a weekly parcel from France, containing the newest Works on Science and General Literature, which he supplies at the rate of One Shilling per Franc on the advertised Price in Paris. He begs to acquaint his friends and the Patrons of German Scientific Works, that he is able to furnish German Works and Periodicals every Month, at the rate of Three Shillings and Sixpence the

J. B. BAILLIÈRE,

LIBRAIRE, RUE HAUTEFEUILLE, PARIS.

C. BAILLY-BAILLIÈRE,

LIBRAIRE, CALLE DEL PRINCIPE, MADRID.

AMERICAN WORKS.

the contract of the contract o	0	8	.7
Actiology and Schill's Outlines of Pathological Semeiology. 8vo. Philadelphia, 1841.			
Agassiz. Lake Superior; its Physical Character, Vegetation, and Animals, compared with those of other and similar Regions. With a Narrative of the Tour. 8vo. Plates.	0	8	0
Allan. Philosophy of the Mechanics of Nature, and the Source and Modes of Action of	0	18	0
Natural Motive-Power. 8vo. Illustrated with numerous Woodcuts. New York, 1852. Allen. A History of the American Revolution, comprehending all the Principal Events	0	18	0
of the Continental Congress, and many of the most important Resolutions of the Continental Congress, and many of the most important Letters of General Washington. 8vo. 2 vols. Baltimore, 1822, (scarce).			
American Agriculturist; designed to Improve the Planter, the Farmer, the Stock Breeder, the Horticulturist. Royal 8vo. With Woodcuts. Vols. I. and II. New			
Almanac and Repository of Useful Knowledge for the Year 1852-53. 12mo.	8	12	
Association for the Advancement of Science (Proceedings of). Fourth Meeting.	0	5	0
8vo. Washington, 1851 Flora (The); or, History of Plants and Wild Flowers, with the Chemical	0	14	0
and Medical Properties of, over Six Thousand Plants. By A. B. Strong. 4 vols. 4to. Coloured Plates. New York, 1848 Institute of Instruction at Providence, Lectures Delivered before	5	10	0
the; including the Journal of Proceedings, and a List of the Offices. 12mo. New Series. 11 vols. Boston, 1841-50	2	5	0
Journal of Science and Arts. See Silliman.	_		
Annals of the Lycenm of Natural History of New York, Vols.	0	12	0
I. to IV. New York, 1852. Vol. V. Parts 1 to 8 Appleton's Mechanics' Magazine and Engineers' Journal. 8vo.	0	16	0
January to December, 1851. With 600 Woodcuts. New York, 1852.	0	17	6
Astronomical Journal. 4to. Published Fortnightly. Vol. I. 1851 Observations made during the years 1845-46, at the National Observatory,	1	1	v
Washington. Published by authority. 2 vols. and Supplement, 4to. Washington, 1846 and 1851	2	2	0
Andubon. Illustrated Notes of an Expedition through Mexico and California. Part I. folio. With 4 Coloured Plates. New York, 1852	0	15	0
Baldwin. Pronouncing Gazetteer: containing Topographical, Statistical, and other Information of all the more Important Places of the Known World, from the most Recent and Authentic Sources; with an Appendix, containing more than 10,000 Additional Names, chiefly of the Small Towns and Villages; and a Supplement, giving			
the Pronunciation of near 2,000 Names. 12mo. Philadelphia, 1851		10	
Bailey. Microscopical Examination of Soundings. 4to. With a Plate. Washington, 1851 Microscopical Observations made in South Carolina, Georgia, and Florida. 4to. With 3 Plates. Washington, 1851	0		6
Bancroft. History of the Colonization of the United States. 14th Edition. 4 vols. 8vo.			
Plates. Boston, 1850 . Barry. The Fruit Garden; a Treatise. With 150 Woodcuts. 12mo. New York, 1851 .	0		0 6
Bartol. A Treatise on the Marine Boilers of the United States. 8vo. With numerous Woodcuts. Philadelphia, 1851	0	7	. 6
Beck. Lectures on Materia Medica and Therapeutics, delivered in the College of Physicians and Surgeons of the State of New York. 8vo. New York, 1851	1	0	0 (
Berzelins. The Use of the Blowpipe in Chemistry and Mineralogy. 8vo. With Plates. Boston, 1845	0		6
Blackwell. The Laws of Life, with Special Reference to the Physical Education of Girls. 12mo. New York, 1852	0	3	3 0
Bonyinge. The Future Wealth of America; being a glance at the Resources of the United States and the Commercial and Agricultural of Cultivating Tea, Coffee, Indigo.			
Booth and Morfit. On Recent Improvements in the Chemical Arts. (Smithsonian			i ()
Report.) 8vo. Washington, 1851 Roydon (Simon) A System of Useful Formulæ Adapted to the Practical Operations	0	J	5 0
of Locating and Constructing Railroads. A Paper Read before the Boston Society of Civil Engineers, Dec. 1849. Svo. Boston, 1851 Bowditch. A History of the Massachusetts General Hospital. Svo. 2 Plates. Boston,	0	14	0
1851	0	15	0
The Young Stethoscopist; or, the Student's ald to Auscultation. 2nd Edition. 12mo. New York, 1848	0	6	0

Bradford. Notes on the North-West or Valley of the Upper Mississippi. 12mo. New	£ s d 0 4 0
Yurk, 1846 Brocklesby. Elements of Metcorulogy, with Questions for Examinations, designed for Schools and Academies. 12mn. 44 Engravings. New York, 1849	0 4 0
Views of the Microscopic World; designed for General Reading, and as a Handbook for Classes in Natural Science. Svo. Illustrated with 247 Wuodcuts and Drawings. New York, 1851	0 6 0
Buchanan. The Culture of the Grape and Wine Making; with an Appendix containing Directions for the Cultivation of the Strawberry. By N. Longworth. 12mo. Cincinnati,	
Burleigh. The Legislative Guide, containing all the Conducting Business in Congress; Jefferson's Manual and the Citizen's Manual, &c. 12mo. Philadelphia, 1852.	0 4 0
Bushe (Gco., M.D.) A Treatise on the Malformations, Injuries, and Diseases of the Rectum and Anus. 8vo. and Atlas 4to. New York, 1837	1 2 6
Byrne. The Practical Metal-Workers' Assistant: containing all Metals and Alloys Forging of Iron and Steel, with the Application of the Art of Electro-Metallurg, to Manufacturing Processes. 8vo. With 537 Engravings. Philadelphia, 1851	1 0 0
The Practical Model Calculator for the Engineer, Mechanic, Manufacturer of Engine Work, Naval Architect, Miner, Millwright, &c. 8vo. With Cuts. Philadelphla	0 18 0
Calhoun (John C.) Works. Vol. I. A Disquisition on Government and a Discourse on the Constitution and Guvernment of the United States. Edited by R. K. CRALLE. 8vo. Charleston, S. C., 1851	0 16 6
Canals. Report of the Select Committee of the Assembly of 1846, upon the Investigation of Frauds in the Expenditures of the Public Moneys upon the Canals of the State of	
New York. 8vo. Albany, 1847. Carson (Joseph, M.D.) Illustrations of Medical Botany, consisting of Coloured Figures of the Plants affording the important articles of the Materia Medica, and	0 10 0
descriptive Letter-press. 2 vols. 4to. Half-bound. Philadelphia, 1847 Channing. A Treatise on Etherization in Childbirth. Illustrated by 581 cases. Syo. Boston, 1848	0 12 0
Collection of Drawings in Detail of the most Approved Construction of American Machinery, with Descriptions, Calculations and Specifications, by the American	0 12 0
Engineering Society of New York. Folio. With 18 Plates. New York, 1852. Comfort. Thompsonian Practice of Midwifery, and Treatment of Complaints peculiar to	2 5 0
Women and Children. Svo. Philadelphia, 1845 Coolcy (B. F.) An Exposition and Explanation of the Modern Phenomena, called	0 5 0
Spirit Manifestations; comprising the Rappings, Movements, Writing Mediums, and various other Phenomena connected therewith; with Extracts from different Writers, of Remarkable Instances of Presentiment, Prophesying, Visions, Clairvoyance, &c. 12mo. Springfield, Mass., 1852	0 16
Copway. The Traditional History and Characteristic Sketches of the Ohjibway Nation. 8vo. With Plates. Boston, 1851	0 10 6
Cones. Outlines of a System of Mechanical Philosophy; being a Research into the Laws of Force. Svo. Bostou, 1851	0 5 6
Curtis. The Relations of Homeopathy to Chemistry; an Inaugural Address delivered before the Hahnemann Academy of Medicine. Svo. New York, 1852	0 1 0
Daguerreian Journal (The). Devoted to the Daguerreian and Photogenic Arts. Vols. I. and H. 8vo. New York, 1850-51 Third Volume. (Now in course of Monthly Publication.) 1852.	1 10 0
Dana. Manual of Mineralogy, including Observatious on Mines, Rocks, Reduction of Ores, and the Applications of the Science to the Arts. 12mo. With 260 Woodcuts.	
Newhaven, 1851 A System of Mineralogy, comprising the most Recent Discoveries. 3rd Edition. 8vo. Illustrated with Dlagrams. New York, 1850	0 8 0
Davis. The Great Harmonia; being a Philosophical Revelation of the Natural, Spiritual, and Celestial Universe. 3 vols. 12mo. Boston, 1851	1 7 0
The Philosophy of Spiritual Intercourse; being an Explanation of Modern Mysteries. 8vo. New York. The Principles of Nature, her Divine Revelations and a Voice to Maukind. 2 vols.	0 4 6
The Law of Deposit of the Flood Tide: its Dynamical Action and Office. 4to.	0 18 0
Washington, 1852 The Approaching Crisis; being a Review of Dr. Bushnell's Recent Lectures on	0 1 6
Supernaturalism. 8va. New York, 1852. The Philosophy of Special Providences, a Vision. 5th Edition. 8vo. Boston, 1851 Dehaas. History of the Early Settlement and Indian Wars of Western Virginia; em-	$\begin{array}{cccc} 0 & 4 & 0 \\ 0 & 2 & 6 \end{array}$
De Tocqueville. American Institutions and their Influences: with Notes by Hon	0 13 0
J. C. Spencer. Post 8vo. New York, 1851 10018. Immortality Trlumphant. The Existence of a God and Human Immortality practically 1008 dered, and the Truth of Divine Revelation Substantiated. 12mo. New York, 1959.	0 8 0
Lectures of the Philosophy of Mesmerism. 12mg New York 1952	0 3 6 0 2 0
The Philosophy of Electrical Psychology. 12mo. New York, 1851	0 4 6

Dunglison. Human Physiology. With upwards of 300 Illustrations. 5th Edition.	£	s d
The Practice of Medicine. A Treatise on Succial Pathology and (Ch.	1 1	0 0
a vois, ovo, a madelphia, 1848	1 1	2 0
Dinlop (William). History of the New Netherlands, Province of New York, and State of New York, to the Adoption of the Federal Constitution. 2 vols. 8vo. With Portrait. New York, 1839	0.1	5 0
Dwight. History of the Hartford Convention; with a Review of the Pollcy of the United States Government which led to the War of 1812. 8vo. New York, 1833		
Eastman. Dahcotah; or, Life and Legends of the Sioux around Fort Snelling. 12mo. With Plates. New York, 1849		6 6
Eberle. A Treatise on the Practice of Medicine, with Notes and Additions by George	0	3 0
M'Lellan, M.D. Two vols. in one. 7th Edition. 8vo. Philadelphia, 1849. Eckfeldt and Dubois. New varieties of Gold and Silver Coins, Counterfeit Coins and Bullion, with Mint Values. With a brief Account of the Collection of Coins belonging to the Mint Counterfeit Counte	1	1 0
Elements of Spiritual Philosophy: being an Exposition of Autorior Principles	0	5 0
Elliott, Cottages and Cottage Life: containing Plans for Country Houses adopted to the	0	2 6
Means and Wants of the People of the United States; with directions for Buildiog and Improving. With some Sketches of Life in this Country. 8vo. With 16 Plates. Cincinnati, 1848		
Emory. Notes of a Military Reconnaissance, from Fort Leavenworth in Missouri to	0 1	0 0
8vo. With Maps and Plates. Washington, 1848	0 1	0 0
Espy. Second Report on Metcorology made to the Navy Department of the United States. Ohlong Charts. 1849	0 1	8 0
Everett. Orations and Speeches on various Occasions. Svo. 2ud Edition. 2 vols. With a Portrait. Boston, 1820	4)	5 0
Finance. Report of the Secretary of the Treasury on the State of the Finances. 8vo.	0	8 0
Flagg (J. F. B.) Ether and Chloroform, their Employment in Surgery, Dentistry, Midwifery, Therapeutics, &c Post 8vo. Philadelphia, 1851	0	4 6
Flint. Clinical Reports on Continued Yellow Fever, based on Analyses of 164 Cases, with its Management, Identity, &c. to which is added a Memoir on the Transportation and		
Diffusion by Contagion of Typhoid Fever. 8vo. Buffalo, 1852 Foster and Whitney. Report on the Geology and Topography of a Portion of	0 1	0 0
Lake Superior Land District. 8vo. With 12 Plates and Map. Washington, 1850.	0	7 0
Fowler. Marriage: its History and Ceremonies: with a Phrenological and Physiological Exposition of the Functions and Qualifications for Happy Marriages. 12mo. With Cuts. New York, 1852	0	4 0
Fremont (Captain.) Report of the Exploring Expedition to the Rocky Mountains in 1842, and to Oregon and North California in 1842-44. 8vo. With Maps and Plates.		
Washington, 1845	0 1	5 0
Oregon and California. The Exploring Expedition to the Rocky Mountains; to which is added a Description of the Physical Geography of California. With Notices of Gold Region. With Plates. Post 8vo. Buffalo, 1851	0 1	0 0
Gayarre (Chas.) Louisiana; its Colonial History and Romance. Svo. New York, 1851 Gerhard. On the Diagnosis of Diseases of the Chest; based upon the Comparison of	0 1	2 6
their Physical and General Signs. Svo. Philadelphia, 1836.	0	5 0
Ciliepsie. A Manual of the Priuciples and Practice of Road-Making, comprising the Location, Construction, and Improvement of Roads, Common, Macadam, &c. and Rail-Roads. 8vo. With 155 Engravings. New York, 1849	0 1	2 0
Girard. Contributions to the Natural History of the Fresh Water Fishes of North America 1. A Monograph of the Cottoids. 4to. With 3 Plates. Washington, 1851.	0	7 0
Girard College. Observations at the Magnetical and Meteorological Observatory at the Girard College, Philadelphia, made under the direction of A. D. BUCKE, in 1840 to		
1845. 3 vols. 8vo. and folio Atlas. Washington, 1847	3	0 0
Gray. Plantæ Wrightianæ Texano Neo-Mexicanæ: an Account of a Collection of Plants made by Charles Wright, in an Expedition from Texas to New Mexico. 4to.		
art I. With 10 Plates. Washington, 1852 The Genera of the Plants of the United States. Illustrated by Figures and Analyses	0 1	8 0
from Nature, by Isaac Spraoue. 2 vols. 8vo. With 186 Plates. New York, 1849. The Botanical Text-Book, an Introduction to the Scientific Botany. 8vo. With	2 1	5 0
Woodcuts. 3rd Edition. New York, 1850	0 1	0 0
Green. Observations on the Pathology of Croup; with Remarks on its Treatment. 12mo. New York, 1849	0	4 0
On the Surgical Treatment of Polyni of the Larynx and Œdema of the Glottis. 8vo. With 4 Plates. New York, 1852	0	6 6
Greeg. Commerce of the Prairies; or, the Journal of a Santa Fé Trader, during Eight Expeditions across the Grent Western Prairies, and n Residence of nearly Nine Years in Northern Mexico. 12mo. 2 vols. With Mans and Engravings. Philadelphia, 1850.	0 1	2 0
Griffith. A Universal Formulary: containing the Methods of Preparing and Administering Officinal and other Medicines. 8vo. Philadelphia, 1850	0 1	

Grines. Phreno-Geology: the Progressive Creation of Man, indicated by Natural History.	£ s d 0 8 0
12mo. New York, 1850 Etherology and the Phreno-Philosophy of Mesmerlsm and Magic Eloquence, 12mo.	0 8 0
New York, 1850 Grinike. Considerations upon the Nature and Tendency of Free Institutions. 8vo. Cincinnati, 1848	0 18 0
Gunnison. The Mormons; or, Latter-Day Saints, in the Valley of the Great Salt Lake. A History of their Rise and Progress, peculiar Doctrines, present Condition, and	0 10 0
Prospects derived from personal observation during a Residence among them. 12mo. With a Plate. Philadelphia, 1852	0 3 0
Mahnemann. Materia Medica Pura. Translated and Edited by Charles J. Hempel, M.D. 4 vols. 8vo. New York, 1846	1 12 0
The Chronic Diseases, their Specific Nature and Homœopathic Treatment. Translated and Edited by Charles J. Hempel, M.D. 5 vols. 12mo. New York, 1846	2 0 0
Hall. The West: its Commerce and Navigation. 12mo. Cincinnati, 1848 Handbook of Young Artists and Amateurs in Oil Painting. By an	0 5 0
American Artist. 12mo. New York, 1849	0 7 0
Harc. Memoir on the Explosiveness of Nitre, with a view to Elucidate its Agency in the Tremendous Explosion of July, 1845, in New York. 4to. Washington, 1849.	0 1 0
Harris. The Principles and Practice of Dental Surgery. 4th Edition. 8vo. With 200 Woodcuts	1 0 0
Harris and Blandy, The American Journal of Dental Science. New Series. 8vo. Vol. I. and Vol. II. Parts 1, 2, 3. Philadelphia, 1850-52. Per Annum. Published Quarterly.	1 10 0
Hartmann (F.) Theory of Acute Diseases and their Homocopathic Treatment. Translated by C. Hempel. 2 vols. 12mo. New York, 1848	0.10.0
Theory of Chronic Diseases and their Homœopathic Treatment. 2 vols. 12mo. New York, 1849	0 18 0
Hastings. The Practice of Surgery; embracing Minor Surgery, and the Application of Dressings. With numerous Illustrations. 12mo. Philadelphia, 1850	0 18 0
Hinckley. Theory of Effect. Embracing the Contrast of Light and Shade of Colour and Harmony. 8vo. With 15 Illustrations. Philadelphia, 1851	0 12 0
Holgate. American Genealogy: being a History of some of the Early Settlers of North	0 4 0
America and their Descendants, from their First Emigration to the Present Time, with their Intermarriages and Collateral Branches, including Notices of Prominent Families and Distinguished Individuals; with Anecdotes, Sketches of the Founding of Cities, &c.	
Home-Book of the Picturesque; or, American Scenery, Art, and Literature.	1 10 0
4to. With 13 Engravings on Steel. New York Homeopathic. The North American Homeopathic Journal. 1851 and 1852. Each.	1 16 0 0 18 0
Examiner. Edited by Drs. Gray and Hempel. 8vo. New Series. Vols. I. and II. New York, 1846-47	1 10 0
Quarterly Homœopathic Journal of Boston. 1852 Hornor. The Medical Students' Guide in Extracting Teeth; with Numerous Cases in the	0 7 0
Surgical Branch of Dentistry. 8vo. With 2 Plates. Philadelphia, 1851 Howard. The Anatomy, Physiology, and Pathology of the Eye. 8vo. Montreal and	0 3 6
Jay (J. C.) A Catalogue of the Shells, arranged according to the Lamarkian System,	0 7 6
with their Authorities, Synonyms, and references to Works where figured or described. 4th Edition. 4to. With Supplement. New York, 1852	1 10
Johnson. The Coal Trade of British America. 8vo. Washington 1850	1 1 0 0 9 0
Jones. Historical Sketch of the Electric Telegraph; including its Rise and Progress in the United States. 8vo. With Woodcuts. New York, 1852	0 6 0
Martin. The Book of Home Beauty. With 12 Portraits of American Ladies. By Martin. 4to. New York, 1852	2 2 0
Knapen. The Mechanics' Assistant; a thorough Practical Treatise on Mensuration and the Sliding Rule, &c. 12mo. With Woodcuts. New York, 1850	0 5 6
Lca. Observations on the Genus Unio, together with Descriptions of new Genera and Species In the Families Najades, Conchie, Collinger, Lymphana Melaplace and Barrier	,,,
I.cc. The Poetical Remains of the late Mary Lee. With a Biographical Memoir 12mo	1 4 0
Charleston, 1851 Leconte. Descriptions of New Species of Colcoptera from California. 8vo. New York, 1851	0 7 6
Synopsis of the Cleridae of the United States Syn New York 1940	$\begin{array}{cccc} 0 & 2 & 6 \\ 0 & 2 & 6 \end{array}$
On the Platygonus Compressus: a New Fossil, Pachyderm. 410 With 4 Plates.	0 5 0
Adminals, with human and general Animal Physiology, Botany, Vegetable Physiology and Geology. Translated from the French with additions, and arranged as a Text Book. By Tuos. Wyatt, A.M. With 49 Plates. 8vo. Philadelphia, 1849	
Traces, evo. Prinadelphia, 1849	0 15 0

Lippincott's Cabinet History of the States, from the Earliest Settlement to the Present Time:—	£	8	d
HISTORY OF GEORGIA. By ARTHUR and CARPENTER. 12mo. Philadelphia, 1852.			
HISTORY OF KENTUCKY. By ARTHUR and CARPENTER. 12mo. Philadelphia,	0	4	6
Locke. Observations on Terrestrial Magnetism. 4to. Washington, 1851	0	4	6
Longienow. Kayanah. A Tale. 12mo. Boston, 1849	0		0
Loomis. The Elements of Geolngy. 12mo. With 82 Cuts. Boston, 1852 The Recent Progress of Astronomy. 12mo. New York, 1850	0	6	0
Lord. The Epoch of Creation. The Scripture Doctrine contrasted with the Geological Theory. 12mo. New York, 1851			
Lynch. Narrative of the United States' Expedition to the River Jordan and the Dead Sea. 7th Edition. 8vo. With Plates. Philadelphia, 1850.	0		0
Macfarlane. History of Propellers and Steam Navigation. 12mo. With Woodcuts. New York, 1851	0		0
McKenny (Thos. L.) Memoirs, Official and Personal; with Sketches of Travels among the Northern and Southern Indians; embracing a War Excursion, and Descriptions of Scenes along the Western Border. Two vols in one. 8vo. With Plates. New York, 1846			
Mansfield. American Education, its Principles and Elements. 8vo. New York, 1851 .	0	8	0
Marcy. The Homopathic Theory and Practice of Medicine. 8vo. New York, 1850. Mather and Brockett. Geography of the State of New York: embracing its Physical Features, Climate, Geology, Mineralogy, Botany, Zoology, History, Pursuits	0	12	0
With Maps. Hartford, 1847	0	5	0
Meigs (C.) Obstetrics: the Science and the Art. 8vo. With 121 Woodcuts. Philadelphia, 1849	0	18	0
A Practical Treatise on the Diseases of Children. 8vo. Philadelphia, 1848 . Woman; her Diseases and Remedies. 2ad Edition. 8vo. Philadelphia, 1851 .	0	18	
Message from the President of the United States to the two Houses of Congress at the Commencement of the First Session of the 31st Congress. 8vo. 3 vols. Wash-		0	
Minific. Text Book of Geometrical Drawing for the Use of Mechanics and Schools. With Illustrations for Drawing Plans, Sections, and Elevations of Buildings and Machinery.			
2nd Edition. With 56 Steel Plates. Royal 8vo. Baltimore, 1850	1	18 0	0
Mitchell. On the Cryptogamus Origin of Malarious and Epidemic Fevers. 12mo. Philadelphia, 1849	0	2	6
Mone. An Outline of Mechanical Engineering, with Drawings of Finished Machines, Tools and Apparatus of every Description which are used in the different branches of Industry, of the Newest and most Approved Construction. With an Appendix, coa-			
taining the necessary Descriptions and Calculations. Folio. New York, 1851-52 . Montgomery, Illustrations of the Law of Kindness. 2nd Edition. With a Sup-	2	5	0
plementary Chapter on Aimsgiving. 12mo. Londou and New York, 1845	0	5	0
and 1844. 8vo. Washington, 1845	0	8	0
Morfit. Chemistry Applied to the Manufacture of Soap and Candles, a thorough Exposition of the Principles, &c. of the Trade, &c. 8vo. With 170 Woodcuts. Philadelphia, 1847	1	8	0
Natural History, New York State Cahinet of. Third Annual Report of the Regents of the University. 8vo. With Cuts and Coloured Plates. Alhany, 1850.	0	10	6
Newman. Fascination, or the Philosophy of Charming; illustrating the Principles of Life in connection with Spirit and Matter. 12mo. With Cuts. New York, 1851.	0	4	6
Nicollet. Report intended to illustrate a Map of the Hydrographical Basin of the Upper Mississippi River. 8vo. With Map. Washington, 1845	0	4	0
Norris. Handbook for Locomotive Engineers and Machinists: comprising the Proportions and Calculations for Constructing Locomotives, Manner of Setting Valves, Tables of Squares, Cubes, Areas, &c. 8vo. With Plate and Cuts. Philadelphia, 1852.	0	7	6
Overman. A Treatise on Metallurgy, comprising Mining, and general and particular Metallurgical Operations, with descriptions of Furnaces, &c. With 377 Wood Engravings. 8vo. New York, 1852	1	5	0
The Manufacture of Steel; containing the Practice and Principles of Working and Making Steel. 12mo. With Woodcuts. Philadelphia, 1851	0	6	0
The Manufacture of Iron in all its various Branches; including a Description of Wood-Cutting, Coal-Digging, &c &c., to which is added an Essay on the Manufacture of Steel. 8vo. With 150 Woodcuts. Philadelphia, 1850	1	10	0
Owen. Report on the Mineral Lands of the United States of America. Svo. With numerons Plates. Washington, 1840	0		0
Palfrey. Statistics of the Condition and Products of certain Branches of Industry In Massachusetts, for the Year endling April 1, 1845. 8vo. Boston, 1846	0	5	0
Patents. Annual Report of the Commissioners of Putents. 1847-51. 8vo.	2	0	0
		-	-

	£sd
Peirce. Examinations of Drugs, Medicines, Chemicals, &c., as to their Purity and	0 5 0
Adulterations. 12mo. Cambridge, 1852. Pharmacopæia of the United States of America (The); By Authority.	0 16 0
Pilgrimage of Thomas Paine and Others to the Seventh Circle in the Spirit	0 4 6
World. Rev. C. Hammond. Medium 12mo. Rochester, 1852 Pocy (F.) Memorias sobre la Historia Natural de la Isla de Cuba. 8vo. Parts 1 and 2. With 16 coloured Plates. Habana, 1851	1 5 0
Westen Organon der Weltgeschichte. 8vo. Cincinnati, 1846	0 4 0
Homœopathic Domestic Physician; containing the Treatment of Diseases, with Popular Explanations of Anatomy, Physiology, Hygiène, and Hydropathy; also an abridged Materia Medica 2nd Edition. 8vo. Cincinnati, 1851	0 10 6
Putnam's Home Cyclopedia. Handbook of Useful Arts. 12mo. With Woodcuts. 670 pages. New York, 1852	0 14 0
Handbook of Universal Biography. 12mo. 821 pages. New York, 1852. Handbook of Literature and the Fine Arts. 12mo. With Woodcuts. 648	0 14 0
pages. New York, 1852 Rau. Organon of the Specific Healing Art. Translated by C. Hempel. 8vo. New York,	0 14 0
1847. Reesc. A Plain and Practical Treatise on the Epidemic Cholera, as it prevailed in the City of New York in the Summer of 1832. With a brief Essay on the use of Ardent Spirits. 8vo. With a Plate. New York, 1833	0 4 0
Reports of the Secretary of War, with Reconnaissances of Routes from San Antonio to El Paso; also the Report of Capt. R. B. Marcy's Route from Fort Smith to	
Santa Fé; and the Report of Lieut. J. N. Simpson, of an Expedition into the Navajo Country; and the Report of Lieut. W. H. C. Whitting's Reconnaissances of the Western	0 12 0
Frontiers of Texas. 8vo. With many Plates and Maps. Washington, 1850. Republic of the United States of America, its Duties to itself and its	0 12 0
Responsible Relations to other Countries; embracing also a Review of the late War between the United States and Mexico, its Causes and Results. 12mo. New York, 1848	0 4 6
Robinson. Catalogue of American Minerals, with their Localities; including all which are known to exist in the United States and British Provinces. 8vo. Boston, 1825.	0 8 0
Mathematical Recreations, containing Solutions of many very Difficult and Important Equations, and of several useful Problems in Geometry, Surveying, &c. 8vo.	0 6 0
Sanderson. Biography of the Signers to the Declaration of Independence. 8vo. With	
Portraits. Philadelphia, 1852 Say. The Organic Laws; or, the Laws which Govern the Human Organism. 12mo. New	2 0 0
Scenes and Thoughts in Europe. By an American. 12mo. New York, 1846.	$\begin{array}{cccc} 0 & 4 & 6 \\ 0 & 2 & 0 \end{array}$
Seconi. Researches on Electrical Rheometry. 4to. With 3 Plates. Washington, 1851. Shepard. A Treatise on Mineralogy. 3rd Edition. With 488 Woodcuts. Vol. I. 8vo.	0 5 0 0 6 0
Shew (Joel, M.D.) Consumption; its Prevention and Cure by the Water Treatment. With Advice concerning Hæmorrhage from the Lungs, Coughs, Colds, Asthma, Brou-	
chitis, and Sore Throats. 12mo. New York, 1850. Silliman, Dana, and Gibbs. The American Journal of Science and Arts. 1852.	$\begin{array}{cccc} 0 & 6 & 0 \\ 1 & 7 & 6 \end{array}$
Published every other month.	1 70
Silliman. Remarks made on a Short Tour between Hartford and Quebec, in the Autumn of 1819. 12mo. 2nd Edition. With Plates. New Haven, 1824. (Very scarce.)	0 5 0
Simpson. Journal of a Military Reconnaissance from Santa Fé, New Mexico to the Navajo Country. 8vo. With Colonred Plates, Maps, &c. Philadelphia, 1852	0 16 6
Smith (Jerome V. C., M.D.) Natural History of the Fishes of Massachussets; embracing a Practical Essay on Angling. 12mo. Boston, 1833	0 4 6
(Mrs. E. Oakes). Woman and her Needs; and Shadow Land or the Seer. 1 vol. 12mo. New York, 1851-52.	0 4 0
A System of Operative Surgery; based upon the Practice of Surgeons in the United States: and comprising a Bibliographical Index and Historical Record of many of their Operations for a Period of 200 Years. Svo. Parts 1 and 2. With 28 Plates.	
Phlladelphia, 1851	1 0 0 0 0 6 0
Smithsonian Contributions to Knowledge. Vols. 1, to IV. 4to. With Maps and Plates. Washington, 1848—52.	
Snell. The Managers' Assistant: being a Condensed Treatise on the Cotton Manufacture.	8 0 0
with Suitable Explanations, &c. 12mo. Roan tuck. Hartford, 1850. Squier (E. G., M.A.) Antiquities of the State of New York; being the Results of Extensive Original Surveys and Explorations, with New York; and Activation of	0 6 6
Extensive Original Surveys and Explorations; with a Supplement on the Antiquities of the West. With 14 Plates and 80 Woodcuts. Post 8vo. Buffalo, 1851 State Papers and Public Documents of the United States, from 1789 to 1815. 8 vols.	0 12 6
8v0. Boston, 1815	1 15 0
Stockhardt. Principles of Chemistry, illustrated by Simple Experiments. From the German. By Principles of With 210 Woodcuts. Cambridge, U.S., 1850. Reduced from 10s. to	0 7 0

Stoddard (Richard Henry). Poems. Boston, 1852	£		d
Massachussets, With 4 Plates, 800 Hels hand 1 1000 Herpetology of	·	J	U
Story. Life and Letters of Joseph Story, Associate Justice of the Supreme Court of the	0	8	0
Story. Life and Letters of Joseph Story, Associate Justice of the Supreme Court of the United States, &c. Edited by his son, WILLIAM W. STORY. 8vo. 2 vols. Boston, 1851 gravings on Steel. 4to. New York 1852	1	16	0
	1	16	0
Stuart (Moses.) A Commentary on Ecclesiastes. 12mo. New York, 1851 Swett. A Treatise on the Diseases of the Chest; being a Course of Lectures Delivered at the New York Hospital. 8vo. New York 1859	0	7	0
	0	15	0
Adylor (Bayard). A Book of Romances, Lyrics and Songs 12mg Poster 1059	0	5	
tibles or Fossil Fuel, including also Notices and Jestiflution of Mineral Combus-			
	1	10	0
Transactions of the American Institute of Homocopathy. Vol. I. 12mo. Philadelphia, 1846	a	8	0
Trantwine. New Method of calculating the Cubit Contents of Excavations and Emhank-	U	0	U
ments, by the aid of Diagrams. 8vo. Philadelphia, 1851 The Field Practice of laying out Circular Curves for Railroads. 12mo. Philadelphia,	0	5	0
1001	0	6	6
Tucker. Elements of the Principles of Midwifery. 8vo. Illustrated with Woodcuts. Philadelphia, 1848	ŭ	Ů	
United States and Canada (Map of the). Folded in Case. New York, 1850		10	
Very Cattle A Complete Treatise on Midwifery or the Theory and Practice of Toyology	U	10	U
including the Diseases of Pregnancy, Labour, and the Puerperal State. Translated and Edited by C. Meros, M.D. 8vo. With Plates. Philadelphia, 1852			
Vestiges of Civilization: or the Etiology of History Religious Esthetland	0	18	0
Tomoreal and Thilosophical, 8vo. New York, 1891	0	10	6
Voices from the Spirit World; being Communications from Many Spirits. By the hand of Isaac Post. Medium 12mo. Rochester, 1852.	0	5	0
Walker. Transactions of the Agricultural Societies in the Commonwealth of Massachusetts, for the Year 1850. 8vo. Boston, 1851	0		Ť
Warren. Supernal Theology, and Life in the Spheres: deduced from Alleged Spiritual	U	8	
Manifestations. Svo. New York, 1852	0	2	6
tutes for Laws and Governments, for the Harmonious Adjustment and Regulation of the			
Pecuniary, Intellectual, and Moral Intercourse of Mankind, proposed as Elements of New Society. 12mo. New York	0	2	e
Wheeler. Historical Sketches of North Carolina, from 1584 to 1851. Illustrated with	U	2	
Engravings. Two vols, in one. 8vo. Philadelphia, 1851 Rural Homes; or, Sketches of Houses suited to American Country Life. 8vo. With	0	14	0
Plates. New York, 1891	0	7	0
Whittlesey. Descriptions of Ancient Works in Ohio. 4to. With 7 Plates. Washington, 1850	^		~
Wilkes. Narrative of the United States' Exploring Expedition during the years 1838 to	O	4	5
1042. With illustrations and Maps. 5 vols. Svo. Half-bound. Philadelphia, 1845.	4	0	0
Wise. System of Aeronautics, comprising its Earliest Investigations and Modern Practice and Art. Svo. With Plates. Philadelphia, 1850	0	8	0
Wood. Treatise on the Practice of Medicine. 3rd Edition. 2 vols. 8vo. Philadelphla,	U	0	0
1852	1	16	0
Wood and Bache. The Dispensatory of the United States of America. 9th Editlon. 8vo. Philadelphia, 1851	1	12	0
Wythes (Jos. H., M.D.) The Microscopist; or, a Complete Manual on the Use of the	*	1.2	0
Microscope; for Physicians, Students, and all Lovers of Natural Science. With Illus-	0	C	0
trations. 12mo. Philadelphia, 1851 Young (Alex.) Chronicles of the Pilgrim Fathers of the Colony of Plymouth, from 160	U	6	(P
to 1625. Now first Collected from Original Records and Contemporaneous Printed	0		^
Documents, and Illustrated with Notes. 2nd Edition. 8vo. Boston, 1844 .	0	14 (13

FRENCH WORKS.

Abcille. Traité des Hydropisies et des Kystes, on des collections séreuses et mixtes	a	9 1	s d
dans les cavites closes naturelles et accidentelles. 8vo. Paris, 1852	Λ	7	6
Agassiz. Système Glaciaire, ou recherches sur les Glaciers, leur Mécanisme, leur ancienne extension et le Rôle qu'ils ont joué dans l'histoire de la Terre. lère partie. Nouvelles			
études et expériences sur les Glaciers actuels, &c. Roy. 8vo. 3 Maps and 9 Plates, folio. Paris, 1847			
Alibert. Des Eaux Minérales dans leurs rapports avec l'Economie publique la Médocine	2	10	0
et la Degislation. 8vo. Paris, 1892	0	2	6
Audre-Michaux. Mémoire sur les causes de la Fièvre Jaune. 8vo. Paris, 1852 Annuaire de l'Economie Politique et de la Statistique pour 1852. 12mo. Paris	0		0
(The previous nine years can be had.)	0	4	0
Ausianx. Faits Cliniques de Chirurgie 8vo. Liège 1852	0	9	6
Al'Chives du Milseum d'Histoire Naturelle publiées per les Professiones Ad	Ü		U
teurs de cet Etablissement. 4to. Vols. I. to V. 170 Plain and Coloured Plates. Paris, 1840—1851. Per vol.	2	٥	0
Arthaud. Examen Medico-Legal des faits relatifs au procès criminel de Jobard. 8vo. Paris, 1852	~	U	U
Bassereau. Traité des Affections de la Peau Symptomatiques de la Symbilia Con-	0	3	6
	0	7	6
Beckensteiner. Etudes sur l'Electricité. Nouvelle méthode pour son emploi Médical. 8vo. Vol. I. avec 8 Planches. Paris, 1852	0	10	C
Beylard. Du Rachitis, de la Fragilité des Os, de l'Ostéomalagie. 4to. 8 Planches. Paris,	U	12	0
Bicheno. Observations on the Linnean Come Linnean Come	0	6	0
Descriptions of two new Shells dto I Plate What Britain 4to. Plate.—Marryat,			•
Species of the Genus Terebolla of Lines Committee Descriptions of five British			
Blanchard. Organisation du Rague Animal (170m 170m).	0	5	0
coloriées. Paris, 1852. Each	0	6	0
See Cutalogue No. 1, p. 37.			
Blanchet. La Sourdi-Mutité-traité Philosophique et Médical. 8vo. 2 vols. avec Figures.			
Bonnet. De la Monomanie du Mountre and 11/4	0	16	0
légale. 8vo. Bordeaux, 1852 Boubee. Cours de Géologie Agricole, Théorique et Pratique. 8vo. Paris, 1852 Bouchardat. De l'Alimentation Insufficient de Pratique. 8vo. Paris, 1852	0	1	6
	0	3	
melle, précédé d'un préciseur l'uniformatique des nouveaux Nés, et des Enlans à la Mam-	0	2	D
2nd Edition. Paris 1859	٥	0	0
et 1851, Collegit Sva Paris 1850 et 1851, Collegit Sva Paris 1850	0	9	y
	0	2	6
Bollighy. Base d'une Nouvelle Phanta	1	0	0
l'Etat Sphéroidal. 8vo. Evreux, 1842	0	r.	0
des Visions, des Songes, de l'Extase, du Magnétisme et du Somnambulisme. 2nd Edition. Brierre de Boismont. Des Hallucinations, ou Histoire Raisonnée des Apparltlons, des Visions, des Songes, de l'Extase, du Magnétisme et du Somnambulisme. 2nd Edition.	U	5 (,
ovo. Paris, 1852	0	6 (0
8vo. Paris, 1852. Vol. 11		0 (,
authentiques. 8vo. Paris 1847	0	7 ()
BIIVIERIEF Statistique Cial to an in the control of	0	4 ()
du Département de la Meuse. 8vo. avec un Atlas de 32 Planches fol. Paris, 1852 Cabart. Leçons de Physique et de Chimie. Rédicées de la Paris, 1852	2	8 (
a l'École Polytechnique & Commission	~	0 (
Plantes, les plus connues et la miadles ou études sur les Propriétés Médicinales de 150	0	7 ()
though the Magnation to the state of the Landstillian flag divarges and			
Productions to A Plant of the Island of Tristan de Cunha and of its Notes) :	2 (•
8vo. Paris, 1852	,	4 0	
Charpiguen. Physiologic, Médecinc et Métaphysique du Magnétisme. 8vo. Paris,) :	2	
1040) (; 0	

	£	8	d	
Chinchilla. Annales Historicos—Historia Generale de la Medecina. 2 Vois.—Historia				
de la Medecina Española. 4 Vols.—Historia de las Principales Operaciones de Cirugia.	5	5	0	
In all, 8 vols. Royal 8vo. Valencia, 1843. Clerget. Analyse des Substances Saccharifères au moyen des Propriétés Optiques de	,	0	•	
leurs Dissolutions. Evaluation du rendement Industriel. 8vo. Plate. Paris, 1850	0	2	0	
Colebrooke. Description of Select Indian Plants. 4to. 6 Plates. 1818.—Knight.				
Upon the Different Species of Esculent Strawberries. 4to.—Salisbury. On the Ger-				
mination of Lyconodium Deuticulatum, 4to, Plate, MACBRIDE, Some Account of the				
Lycoperdon Solidium of the Flora Virginica, the Lycoperdon Cervinum of Walter. 4to.				
-Saure. An Account of the Rhynzomorpha Medullaris, a new British Fungus. 4to.	٥	5	٥	
Plate. The above together in one Vol. (From Transactions.)	v	U	U	
Comptes Rendus des Séances et Mémoires de la Société de Biologie. Royal 8vo. Vol.	0	7	0	
III. Paris, 1852		14		
Cruveilhier. Traité d'Anatomie Pathologique Général. 8vo. Vol. II. Paris, 1852	0	9	0	
De Caudolle. Prodromus Systematis Naturalis Regni Vegetabilis, &c., &c. 8vo. Vol.				
VIII Part 1 Paris, 1852	0	16	0	
The Curzon. Etudes sur les Enfants Trouvés au point de vue de la Législation, de la		-	^	
Move et de l'Economie Politique, 8vo. Paris, 184/ · · · · ·	0	7	0	
Essai d'Ichtvologie des Côtes Océaniques et de l'Intérieur de la France, ou	0	2	0	
diametrica des Poissons observés Svo. 2 Plates. Angers, 1001	V	9	U	
Thevereie. Medecine Legale, Theorique et Pratique, revus et annotes par Denaussy de	1	3	0	
Robecourt. 3rd Edition. 3 vols. 8vo. Paris, 1892.	1		0	
D'Orbigny. Prodrome de Paléontologie. 3 vols. 12mo Cours Elémentaire de Paléontologie et de Géologie Stratigraphiques. 3 vols. 12mo.				
area Sympos at Atlas Ato Paris, 1852	0	15	0	
ot Conto Gaologie annliquée aux Arts et à l'Agriculture, comprenant	_	1.0		
11 bla deg Dévolutions du Globe Svo. Fips. Paris, 1001 .		$\frac{10}{12}$		
The manual Maladies de la Matrice. 3rd Edition. 0vo. 2 vois. 1 ans, 1000		3		
Du Potet. Manuel de l'Etudiant Magnétiseur. 2nd Edition. 12mo. Paris, 1851	·	·		
Duval (Vincent). Traité Théorique et Pratique de la Maladie Scrofuleuse. 8vo. Paris,	0	8	0	
1852 . Edwards. Introduction à la Zoologie générale, ou Considerations sur les Tendances de Edwards. Introduction à la Zoologie générale, ou Considerations sur les Tendances de Edwards. Introduction à la Zoologie générale, ou Considerations sur les Tendances de Edwards.				
la Nature dans la Constitution du Règne Animal. 12mo. (Première Partie.) Paris,	_			
		6	3 0	
25 / 114 de la familió de Médecine de Strasbourg. Description de				
deux Fœtus Monstres dont l'un Acephale et l'autre Monopode. Folio. avec 4 planches.	() (0	
Elie de Beaumont. Notice sur les systèmes de Montagnes. 3 vols. 12mo. Paris,		15) ()	1
1852 . Escallier. La Méthode Homeopathique et la Médication Ordinaire comparées dans le	,	, ,		
		,	1 6	1
Fairet. De la Construction et de l'Organisation des Etablissements d'Allénés. 4to.	. () :	3 ()
Paris, 1352 Ferussae (M. de) Histoire Naturelle, Générale et particullère des Mollusques, tant des Espèces qu'on trouve aujourd'hui vivantes que des déponilles Fossiles de celles qui des Espèces qu'on trouve aujourd'hui vivantes que des déponilles Fossiles de celles qui des Espèces qu'on trouve aujourd'hui vivantes que des déponilles Fossiles de celles qui				
n'existent plus, classes d'après les caractères essentits que livraison, par G. P. leurs Coquilles. Par M. de Férussac, continué depuis la 20ème livraison, par G. P. leurs Coquilles. Par M. de Férussac, Coloned Original price, £63, now re-				
	2	4 1	0 ()
duced to		0		
Plain, original price, £15 los. now reduced to the Cephalodes of the Corbigny. Histoire Naturelle, Générale et Particulière des Cephalodes et d'Orbigny. Histoire Naturelle, Générale et Particulière des Cephalodes et d'Orbigny. Histoire Naturelle, Générale et Particulière des Cephalodes et d'Orbigny. Histoire Naturelle, Générale et Particulière des Cephalodes et d'Orbigny. Histoire Naturelle, Générale et Particulière des Cephalodes et d'Orbigny. Histoire Naturelle, Générale et Particulière des Cephalodes et d'Orbigny. Histoire Naturelle, Générale et Particulière des Cephalodes et d'Orbigny. Histoire Naturelle, Générale et Particulière des Cephalodes et d'Orbigny. Histoire Naturelle, Générale et Particulière des Cephalodes et d'Orbigny. Histoire Naturelle, Générale et Particulière des Cephalodes et d'Orbigny. Histoire Naturelle, Générale et Particulière des Cephalodes et d'Orbigny. Histoire Naturelle, Générale et Particulière des Cephalodes et d'Orbigny. Histoire Naturelle, Générale et Particulière des Cephalodes et d'Orbigny.		-1	0 ()
Acétabuliféres vivants et lossites à la sant Mallysques classés en Familles Naturelles				
dans lesquelles on a établi la concordance de tous les systèmes, suivis d'un Prodrome dans lesquelles on a établi la concordance de tous les systèmes, suivis d'un Prodrome dans lesquelles on a établi la concordance de tous les systèmes, suivis d'un Prodrome	3	0 1	0 (0
dans lesquelles on a établi la concordance de tous les systemes, suivis d'activité de général pour tous les mollusques terrestres ou fluviatiles, vivants ou fossiles. Gr. 4to général pour tous les mollusques des Principales découvertes scientifiques modernes	•	0 1	0	0
général pour tous les mollusques terrestres ou nuvatiles, découvertes scientifiques modernes Figuier. Exposition et Histoire des Principales découvertes scientifiques modernes Figuier. Exposition et Histoire des Principales découvertes scientifiques modernes Vaneur à Vaneur Bâteaux à Vaneur Chemins de Fer.) 12mo. Paris, 1850	2	0	3	G
Figurer. Exposition et Histoire des Principales decourres de Fer.) 12mo. Paris, 1850. Vol. 11I. (Machine à Vapeur, Bâteaux à Vapeur, Chemins de Fer.) 12mo. Paris, 1850.				
Vol. III. (Machine à Vapeur, Baccaux à Capeur, Partique et Raisonné d'Hydrothérapie. Recherches Cliniques sur l'appli Fleury. Traité Pratique et Raisonné d'Hydrothérapie. Recherches Cliniques sur l'appli				
cation de cette Medication au tratement des		0	7	6
8vo. Paris, 1892	_			
Forest. De la Question du reboisement et Nouvel Examen des theoristes. Svo. Paris logiques et des faits Economiques qui se rattachent à l'existence des Forêts. Svo. Paris	,	0	2	0
1852	t	O.	-	-
1852 Le Matérialisme et la Phrénologie combattus dans leurs fondements, e Forichon. Le Matérialisme et la Phrénologie combattus dans leurs fondements, e Haller dans son état Normal et ses Aberrations; dans le Délire, les Haller dans son état Normal et ses Aberrations;				
Forichon. Le Matérialisme et la Phrénologie compattus dans le Délire, les Hal l'Intelligence étudiée dans son état Normal et ses Aberrations; dans le Délire, les Hal l'Intelligence étudiée dans son état Normal et ses Aberrations; dans le Délire, les Hal	•		6	
lucinations, 1a Fone, les Songes et chez les		0	3	6
Foliquet. De la Dysentérie. 800. Values, 1002. Physical Rumaines mises à la porté	C			
Galet. Le Corps de l'Homme, ou l'Anatomie et la Physiologie d'anatomies. Paris, 1836 de toutes les classes de la Société, avec 177 Planches coloriées lithographiées. Paris, 1836	. 1	0	0	0
de toutes les classes de la Societe, avec				
4 vols. 4to. Ganot. Traité Elémentaire de Physique Expérimentale et Appliquée. With 420 Wood		0	6	6
cuts. 12mo. Parls, 1852				

			Ξ.
Garnier. Traité des Facultés de l'âme comprenant l'Histoire des principales Théories	£		d
Psycologiques. 8vo. 3 vols. Paris, 1852. Gauthier. Traité Pratique du Magnétisme et du Somnambulisme. 8vo. Paris, 1845.	1 0		0
Gantier (d'Agouty). Explication de tous les Muscles de la tête en huit planches et de tont l'Anatomie de la Tête, en neuf planches. IMPRIMÉES EN COULEUR. Follo. n. d			0
Girardin. Leçons de Chimie Elémentalre, appliquées aux arts industriels et faites le Dimanche, à l'Ecole Municipale de Rouen. 3rd Edition. 8vo. 2 vols. Figs. et Echan-	-	v	U
tillons d'Indiennes, Paris, 1846	0	14	0
et Duhreuil. Cours Elémentaire d'Agriculture. 12mo. 2 vols. With 462 Engravings. Paris, 1852	0	15	0
Grisolle. Traité Elémentaire et Pratique de Pathologie Interne. 5th Edition. 2 vols. 8vo. Paris, 1852	0	18	0
Gnerard. Du Choix et de la Distribution des Eanx, dans une Ville. 8vo. Paris, 1852.	0	2	
Guislain. Leçons Oraies sur les Phrénopathies ou Traité Théorique et Pratique des Maladies Mentales. 8vo. 3 vols. 54 Cuts. Gand, 1852	1	4	0
Hirschweld et Leveillee. La Nevrologie, ou Description et Iconographie du Système Nerveux et des Organs des sens de l'Homme, avec leur Modes de Préparations.			
(Il sera publice en 10 livraisons chacune de 9 planches, avec texte descriptif.) Parts 1 to 9 are published, the rest to appear in two months. Price, each Part, Coloured	0	10	0
Hoffmann. La Rage et le Choléra. 8vo. Paris, 1852		5	0
Jamain. Nouveau traité Elémentaire d'Anatomie descriptive et des préparations Ana	U	1	U
tomiques, suivi d'un Précis d'Embryologie, par A. Verneuil. 12mo. With 146 Cuts. Paris, 1853	0 1	12	0
Manuei de petite Chirurgie, contenant les Pansements, les Bandages, les Ventouses, les Moxas, Saignée, &c. &c. 2nd Edition. 12mo. Paris, 1852	0	6	0
James. De l'Emploi des eaux Minérales dans le traitement des Accidents consécutifs de la Syphilis. 8vo. Paris, 1852	0	1	6
Jarjavay. Traité d'Anatomie Chirurgicale ou de l'Anatomie dans ses Rapports avec la Pathologie externe et la Médecine opératoire. Vol. I. 8vo. Paris, 1852	0	7	
Jobert (De Lamballe.) Traité des Fistules Vesico-Uterines, Vesico-Utero- Vaginales, Entero-Vaginales et Recto-Vaginales. 8vo. 10 Cuts. Paris, 1852.	0		
Jordan. Pugillus Plantarum Novarum Presertim Gallicarum and la Bannart and	0	7	6
l'Essai de Phytostatique appliqué à la Chaine du Jura. Svo. Lyon, 1850 Kenneha. Ethnographical Atlas. 4 sheets. Grand folio, coloured, and 8vo. Text. (In Bussian) St. Potorchurch 1859	0	4	0
Kiener. Spécies général et iconographie des Cognilles Vinentes Bulliés une Re	1 !	2	6
lection de Lamarck, celle de M R Deleggert De L Viere Co-			
L'ouvrage se composera d'environ 160 à 180 livrigans, et de celles du Prince Masséna.			
pages de texte. Les liv. l à 138 sont en vente. Prix de chacune, grand 8vo., papier raisin superfin, satiné, fig. col.	0	0	_
In 4to. fig. col. Kirby. A Century of Insects, including several new Genera described from his Cabinet. 4to. With 3 coloured Plates. (From Typescotium)	0 1	6	
	0 1	0	0
Knight. On Esculent Strawberries. 4to. (From Transactions.) See Colebrooke. Kobelt (De). Appareil du sens Genital des deux Sexes dans l'espèce Humaine et dans quelques Mampifères au point de vue Apatemises als l'espèce Humaine et dans			
l'Allemande par H. Kaula, D.M., avec 5 plauches Que Strackard livil	0	5 (0
de Beigique. 4to. 2 vols, and Supplement With 74 Plates Lin 1010 arbonnifère			
With Woodcuts. Paris 1846	4 1	9 ()
Labourt. Recherches Historiques et Statistiques sur l'Intempérance des Classes	0 :	3 ()
remedier à l'abus des Boissons enjurantes des Boyletts qu'il convient d'employer pour	0	7 (;
de l'Institut sulvi du Rauport de MM Théres d'Il e sang. Lu à l'Académie des Sciences		1 6	
Sciences, Morales et Politiques, 27 Mars et 12 Amil 10:00 Lin à l'Académie des			
les Théologiens, par M. L'Abba I B. I. Sura Reis Corps savants, la Cour de Romc et	0	1 ()
Pétersbourg, pour servir à l'Histoire de Russie et particullèrement à Saint	0 2	7 ()
Bienfaisance dans l'Emplre de Russia 19ma Distribution et de	0 :	2 0	
4to. With 20 Plates. Berlin 1848	0 18		
Machride. On the Lycoperdon Solidum of the Flora Virginica, &c. 4to. (From Transactions.) See Colebrooke.		<i>J</i> 0	
Macquart. Les Arbres et Arbrisseaux d'Europe, et leurs Insectes. 8vo. Lille, 1852. Maillot. Histoire de la Percussion depuis les Temps les plus reculés jusqu'à nos jours. 8vo. Paris, 1852	0 6	3 0	
8vo. Parls, 1852 and deputs les Temps les plus reculés jusqu'à nos jours.		3 0	

	£	s d	
Maisonneuve. Leçons Cliniques sur les Affections Cancéreuses, professée à l'Hôpital			
Cochin, recusillies et publiées, par M. le Dr. Alexis Favnor, avec plauches, lère partie,	_		
comprenant les Affections Cancéreuses en général. 8vo. Paris, 1852.	0	2 0	
Mandl. Manuel d'Anatomle générale appliquée à la Physiologie et à la Pathologic. 8vo. With 5 Plates. Paris, 1843	0	8 0	
Marchal (de Calvi.) Des Epidémies. 8vo. Parls, 1852.	0	3 (
Marryat. New Shells, &c. (From Transactions.) See BICHENO.			
Martin. Photographie, Nouvelle procédé pour obtenir des Epreuves positives directes sur			
Glace. 8vo. Paris, 1852	0	2 6	
Montagn. Genus Terebella, &c. (From Transactions.) See BICHENO.			
Morel. Etudes cliniques, ou traité Théorique et Pratique des Maladies Mentales, con-			
siderées dans leur uature, leur traitement et dans leur rapport avec la Médecine légale		0 1	
des Aliénés. Vol. I. and Plates. 8vo. Paris, 1852	0	8 (1
Mongcot (Dr. A.) Essai d'une Flore du nouveau grès rouge des Vosges, ou description	0	3 (
des Végétaux Silicifés qui s'y rencontrent. 8vo. fig. Epinal, 1852	v	0 (
Plate. Paris, 1852	0	3 ()
Museum d'Histoire Naturelle de Paris. Royal 8vo. Paris, 1850-51:			
Catalogue Méthodique des Reptiles. Parts 1 and 2. Paris, 1851	0	10 ()
De la Collection Entomologique. Parts 1 and 2. Paris, 1851	0	10)
De la Collection des Mammifères. Part 1. Paris, 1850	0	5 ()
Orfila. Traité de Toxicologie. 5th Edition. 2 vols. Svo. Paris, 1852	1	0 ()
Ostzowski. Etudes sur la Mathèse ou Anarchie et Hiérarchie de la Science. 8vo. avee			
Figures. Paris, 1849	0	4	0
Parchappe. Des Principes à suivre dans la Fondation et la Construction des Asiles			
d'Aliénés. Royal 8vo. 3ème Partie, avec Figures. Paris, 1852	0	9	0
Parigot. Thérapeutique Naturelle de la Folie. 8vo. Bruxelles, 1852.	0	2	6
Patissier. Rapport sur le Service Médical des Etablissements Thermaux, pour les Anuées			
1849 et 1850. 4to. Paris, 1852	0	6	0
Payen. Traité de la Fabrication et du Raffinage des Sucres de Cannes, de Betteraves,		_	
Sucre Candi, Sucre d'Orge, de Pommes, &c. 8vo. 12 Plates. Paris, 1832	0	5	()
Pictet et Roux. Description des Mollusques Fossiles qui se trouvent dans les grès		=	^
verts des Environs de Genève. 4to. Parts I to 3. With 40 Plates. Geneve, 1847-32 .	2	5	U
Part 1—Céphalopodes. Part 2—Gasteropodes. Part 3—Acéphales Orthoconiques.			
Plec. Types de chaque Famille et des principaux genres des Plantes croissant spontané-			
ment en France; l'exposition détaillée et complete de leurs Caracteles et de l'Emoly- ologie. 4to. With Coloured Plates. Parts 1 to 65. Paris, 1852. Each Part	0	1	65
Margarias cohre la Historia Natural de la Isla de Cuba, accompanados de Sumarios			
Latinos y Extractos la Frauces. Vol. I. Parts 1, 2. With 16 Coloured and Plain Plates.	1	5	0
Habana, 1851	Ī		
To be completed in 24 Parts, forming 4 Vols.; one Part to appear every Six Weeks.			
Pouson. Traité de l'Exploitation des Mines de Houille ou Exposition Comparative des			
Méthodes employées en Belgique, en France, en Allemagne, et en Angioterie, pour l'Arrachement et l'Extractiou des Mineraux Combustibles. Vol I. Royal 8vo. Atlas and	1	0	6
Méthodes employées en Belgique, en France, en Allemagne, et en Angetetie, pour l'Arrachement et l'Extractiou des Mineraux Combustibles. Vol I. Royal 8vo. Atlas and Figures. Liège, 1852. Pouillet. Notions Générales de Physique et de Météorologie à l'usage de la Jeunesse.	1		
Méthodes employées en Belgique, en France, en Allemagne, et en Angetetre, pour l'Arrachement et l'Extractiou des Mineraux Combustibles. Vol I. Royal 8vo. Atlas and Figures. Liège, 1852 Pouillet. Notions Générales de Physique et de Météorologie à l'usage de la Jeunesse. 12mo. 353 Cuts. Paris, 1850	0	6	0
Méthodes employées en Belgique, en France, en Anemagne, et en Angetet April Parrachement et l'Extractiou des Minéraux Combustibles. Vol I. Royal 8vo. Atlas and Figures. Liège, 1852 Pouillet. Notions Générales de Physique et de Météorologie à l'usage de la Jeunesse. 12mo. 353 Cuts. Paris, 1850 De l'Hygiène de l'Habitation. 8vo. Paris, 1851.	0 0		0
Méthodes employées en Belgique, en France, en Anemagne, et en Angetet April Parrachement et l'Extractiou des Minéraux Combustibles. Vol I. Royal 8vo. Atlas and Figures. Liège, 1852 Pouillet. Notions Générales de Physique et de Météorologie à l'usage de la Jeunesse. 12mo. 353 Cuts. Paris, 1850 De l'Hygiène de l'Habitation. 8vo. Paris, 1851.	0 0	6 1	0 6
Méthodes employées en Belgique, en France, en Antemagne, et en Angetetic, pour l'Arrachement et l'Extractiou des Minéraux Combustibles. Vol I. Royal 8vo. Atlas and Figures. Liège, 1852 Pouillet. Notions Générales de Physique et de Météorologie à l'usage de la Jeunesse. 12mo. 353 Cuts. Paris, 1850 Queriere. De l'Hygiène de l'Habitation. 8vo. Paris, 1851. Rackett. Observations on a Viper found in Cranborne Chace, Dorsetshire. 4to. London, 1877. From Trynspotions.)		6 t	0 6
Méthodes employées en Belgique, en France, en Antemagne, et en Angetetre, pour l'Arrachement et l'Extractiou des Minéraux Combustibles. Vol I. Royal 8vo. Atlas and Figures. Liège, 1852 Pouillet. Notions Générales de Physique et de Météorologie à l'usage de la Jeunesse. 12mo. 353 Cuts. Paris, 1850 Queriere. De l'Hygiène de l'Habitation. 8vo. Paris, 1851. Rackett. Observations on a Viper found in Cranborne Chace, Dorsetshlre. 4to. London, 1817. (From Transactions.)		6 1	0 6
Méthodes employées en Belgique, en France, en Antemagne, et en Angetetre, pour l'Arrachement et l'Extractiou des Minéraux Combustibles. Vol I. Royal 8vo. Atlas and Figures. Liège, 1852 Pouillet. Notions Générales de Physique et de Météorologie à l'usage de la Jeunesse. 12mo. 353 Cuts. Paris, 1850 Queriere. De l'Hygiène de l'Habitation. 8vo. Paris, 1851. Rackett. Observations on a Viper found in Cranborne Chace, Dorsetshlre. 4to. London, 1817. (From Transactions.)		6 1	0 6
Méthodes employées en Belgique, en France, en Allemagne, et angetetit, pour l'Arrachement et l'Extractiou des Minéraux Combustibles. Vol I. Royal 8vo. Atlas and Figures. Liège, 1852 Pouillet. Notions Générales de Physique et de Météorologie à l'usage de la Jeunesse. 12mo. 353 Cuts. Paris, 1850 Querière. De l'Hygiène de l'Habitation. 8vo. Paris, 1851. Rackett. Observations on a Viper fouud in Cranborne Chace, Dorsetshlre. 4to. London, 1817. (From Transactions.) Reveille-Parise. Traité de la Vieillesse Hygiénique, Médical et Philosophique, on Recherches sur l'Etat Physiologique, les Facultés morales, les Maladies de l'Age avancé, et sur les moyens les plus surs, les mieux experimentés de soutenir et de prolonger l'Articlé Vivile à cette époque de l'existence. 8vo. Paris, 1853	6	6 1 0	0 6 6
Méthodes employées en Belgique, en France, en Allemagne, et angetetit, pour l'Arrachement et l'Extractiou des Minéraux Combustibles. Vol I. Royal 8vo. Atlas and Figures. Liège, 1852 Pouillet. Notions Générales de Physique et de Météorologie à l'usage de la Jeunesse. 12mo. 353 Cuts. Paris, 1850 Querière. De l'Hygiène de l'Habitation. 8vo. Paris, 1851. Rackett. Observations on a Viper fouud in Cranborne Chace, Dorsetshlre. 4to. London, 1817. (From Transactions.) Reveille-Parise. Traité de la Vieillesse Hygiénique, Médical et Philosophique, on Recherches sur l'Etat Physiologique, les Facultés morales, les Maladies de l'Age avancé, et sur les moyens les plus surs, les mieux experimentés de soutenir et de prolonger l'Articlé Vivile à cette époque de l'existence. 8vo. Paris, 1853	6	6 1 0	0 6 6
Méthodes employées en Belgique, en France, en Anemagne, et an Angetette, pour l'Arrachement et l'Extractiou des Minéraux Combustibles. Vol I. Royal 8vo. Atlas and Figures. Liège, 1852 Pouillet. Notions Générales de Physique et de Météorologie à l'usage de la Jeunesse. 12mo. 353 Cuts. Paris, 1850 Queriere. De l'Hygiène de l'Habitation. 8vo. Paris, 1851. Rackett. Observations on a Viper fouud in Cranborne Chace, Dorsetshire. 4to. London, 1817. (From Transactions.) Reveille-Parise. Traité de la Vieillesse Hygiénique, Médical et Philosophique, ou Recherches sur l'Etat Physiologique, les Facultés morales, les Maladies de l'Age avancé, et sur les moyens les plus surs, les mieux experimentés de soutenir et de prolonger l'Activité Vitule à cette époque de l'existence. 8vo. Paris, 1853 Ricard. Le Magnétisme Traduit en Cour d'Assisses. Acquittement. 8vo. Paris, 1845.	0	6 1 0	0 6 6
Méthodes employées en Belgique, en France, en Antemagne, et angetet, pour l'Arrachement et l'Extractiou des Minéraux Combustibles. Vol I. Royal 8vo. Atlas and Figures. Liège, 1852 Pouillet. Notions Générales de Physique et de Météorologie à l'usage de la Jeunesse. 12mo. 353 Cuts. Paris, 1850 Querière. De l'Hygiène de l'Habitation. 8vo. Paris, 1851. Rackett. Observations on a Viper found in Cranborne Chace, Dorsetshlre. 4to. London, 1817. (From Transactions.) Reveille-Parise. Traité de la Vieillesse Hygiénique, Médical et Philosophique, on Recherches sur l'Etat Physiologique, les Facultés morales, les Maladies de l'Age avancé, et sur les moyens les plus surs, les mieux experimentés de soutenir et de prolonger l'Activité Vitule à cette époque de l'existence. 8vo. Paris, 1853 Ricard. Le Magnétisme Traduit en Cour d'Assises. Acquittement. 8vo. Paris, 1845. Robin et Verdeil. Traité de Chimie Anatomique et Physiologique ou des Principes	0	6 1 0	0 6 6
Méthodes employées en Belgique, en France, en Alemagne, et angetetre, pour l'Arrachement et l'Extractiou des Minéraux Combustibles. Vol I. Royal 8vo. Atlas and Figures. Liège, 1852 Pouillet. Notions Générales de Physique et de Météorologie à l'usage de la Jeunesse. 12mo. 353 Cuts. Paris, 1850 Querière. De l'Hygiène de l'Habitation. 8vo. Paris, 1851. Rackett. Observations on a Viper fouud in Cranborne Chace, Dorsetshlre. 4to. London, 1817. (From Transactions.) Reveille-Parise. Traité de la Vieillesse Hygiénique, Médical et Philosophique, on Recherches sur l'Etat Physiologique, les Facultés morales, les Maladies de l'Age avancé, et sur les moyens les plus surs, les mieux experimentés de soutenir et de prolonger l'Activité Vitule à cette époque de l'existence. 8vo. Paris, 1853 Ricard. Le Magnétisme Traduit en Cour d'Assises. Acquittement. 8vo. Paris, 1845. Robin et Verdeil. Traité de Chimie Anatomique et Physiologique ou des Principes immédiats du corps de l'Homme et des Mammitères à l'état Normal et à l'état Patholoimmédiats du corps de l'Homme et des Mammitères à l'état Normal et à l'état Patholoimmédiats du corps de l'Homme et des Mammitères à l'état Normal et à l'état Patholoimmédiats du corps de l'Homme et des Mammitères à l'état Normal et à l'état Patholoime de de Microscopie, agrégé à	0	6 1 0	0 6 6
Méthodes employées en Belgique, en France, en Alemagne, et Angetette, pour l'Arrachement et l'Extractiou des Minéraux Combustibles. Vol I. Royal 8vo. Atlas and Figures. Liège, 1852 Pouillet. Notions Générales de Physique et de Météorologie à l'usage de la Jeunesse. 12mo. 353 Cuts. Paris, 1850 Querière. De l'Hygiène de l'Habitation. 8vo. Paris, 1851. Rackett. Observations on a Viper fouud in Cranborne Chace, Dorsetshlre. 4to. London, 1817. (From Transactions.) Reveille-Parise. Traité de la Vieillesse Hygiénique, Médical et Philosophique, on Recherches sur l'Etat Physiologique, les Facultés morales, les Maladies de l'Age avancé, et sur les moyens les plus surs, les mieux experimentés de soutenir et de prolonger l'Activité Vitule à cette époque de l'existence. 8vo. Paris, 1853 Ricard. Le Magnétisme Traduit en Cour d'Assises. Acquittement. 8vo. Paris, 1845. Robin et Verdeil. Traité de Chimie Anatomique et Physiologique ou des Principes immédiats du corps de l'Homme et des Mammifères à l'état Normal et à l'état Pathologique, par MM. Ch. Robin, Professeur d'Anatomie Géuérale et de Microscopie, agrégé à la Esquité de Médecine de Paris, et Verdeil, Professeur de Chimie. 3 vols. 8vo. avec	0	6 1 0	0 6 6
Méthodes employées en Belgique, en France, en Anemagne, et angetetre, pour l'Arrachement et l'Extractiou des Minéraux Combustibles. Vol I. Royal 8vo. Atlas and Figures. Liège, 1852 Pouillet. Notions Générales de Physique et de Météorologie à l'usage de la Jeunesse. 12mo. 353 Cuts. Paris, 1850 Queriere. De l'Hygiène de l'Habitation. 8vo. Paris, 1851. Rackett. Observations on a Viper fouud in Cranborne Chace, Dorsetshire. 4to. London, 1817. (From Transactions.) Reveille-Parise. Traité de la Vieillesse Hygiénique, Médical et Philosophique, on Recherches sur l'Etat Physiologique, les Facultés morales, les Maladies de l'Age avancé, et sur les moyens les plus surs, les mieux experimentés de soutenir et de prolonger l'Activité Vitule à cette époque de l'existence. 8vo. Paris, 1853. Ricard. Le Magnétisme Traduit en Cour d'Assises. Acquittement. 8vo. Paris, 1845. Robin et Verdeil. Traité de Chimie Anatomique et Physiologique ou des Principes immédiats du corps de l'Homme et des Mammifères à l'état Normal et à l'état Pathologique, par MM. Ch. Robin, Professeur d'Anatomie Générale et de Microscopie, agrégé à la Faculté de Médecine de Paris, et Verdeil, Professeur de Chimie. 3 vols. 8vo. avec la Faculté de Médecine de Paris, et Verdeil, Professeur de Chimie. 3 vols. 8vo. avec la Faculté de Médecine de Paris, et Verdeil, Professeur de Chimie. 3 vols. 8vo. avec la faculté de Médecine de Paris, et Verdeil, Professeur de Chimie. 3 vols. 8vo. avec la faculté de Médecine de Paris, et Verdeil, Professeur de Chimie. 3 vols. 8vo. avec la faculté de Médecine de Paris, et Verdeil, Professeur de Chimie. 3 vols. 8vo. avec la faculté de Médecine de Paris, et Verdeil, Professeur de Chimie. 3 vols. 8vo. avec la faculté de Médecine de Paris, et Verdeil, Professeur de Chimie.	0	6 t 0 7 2	0 6 6
Méthodes employées en Belgique, en France, en Alemagne, et Angetette, pour l'Arrachement et l'Extractiou des Minéraux Combustibles. Vol I. Royal 8vo. Atlas and Figures. Liège, 1852 Pouillet. Notions Générales de Physique et de Météorologie à l'usage de la Jeunesse. 12mo. 353 Cuts. Paris, 1850 Querière. De l'Hygiène de l'Habitation. 8vo. Paris, 1851. Rackett. Observations on a Viper fouud in Cranborne Chace, Dorsetshlre. 4to. London, 1817. (From Transactions.) Reveille-Parise. Traité de la Vieillesse Hygiénique, Médical et Philosophique, on Recherches sur l'Etat Physiologique, les Facultés morales, les Maladies de l'Age avancé, et sur les moyens les plus surs, les mieux experimentés de soutenir et de prolonger l'Activité Vitule à cette époque de l'existence. 8vo. Paris, 1853 Ricard. Le Magnétisme Traduit en Cour d'Assises. Acquittement. 8vo. Paris, 1845. Robin et Verdeil. Traité de Chimie Anatomique et Physiologique ou des Principes immédiats du corps de l'Homme et des Mammifères à l'état Normal et à l'état Pathologique, par MM. Ch. Robin, Professeur d'Anatomie Géuérale et de Microscopie, agrégé à la Faculté de Médecine de Paris, et Verdeil, Professeur de Chimie. 3 vols. 8vo. avec Atlas de 45 Planches gravées. Paris, 1853. Rodenberg. Mémoire et Observations sur l'Accouchement Prématuré Artificiel. 8vo.	0	6 t 7 2 16	0 6 6
Méthodes employées en Belgique, en France, en Alemagne, et angetetic, pour l'Arrachement et l'Extractiou des Minéraux Combustibles. Vol I. Royal 8vo. Atlas and Figures. Liège, 1852. Pouillet. Notions Générales de Physique et de Météorologie à l'usage de la Jeunesse. 12mo. 353 Cuts. Paris, 1850. Queriere. De l'Hygiène de l'Habitation. 8vo. Paris, 1851. Rackett. Observations on a Viper found in Cranborne Chace, Dorsetshlre. 4to. London, 1817. (From Transactions.) Reveille-Parise. Traité de la Vieillesse Hygiénique, Médical et Philosophique, on Recherches sur l'Etat Physiologique, les Facultés morales, les Maladies de l'Age avancé, et sur les moyens les plus surs, les mieux experimentés de soutenir et de prolonger l'Activité Vitule à cette époque de l'existence. 8vo. Paris, 1853. Rieard. Le Magnétisme Traduit en Cour d'Assises. Acquittement. 8vo. Paris, 1845. Robin et Verdeil. Traité de Chimie Anatomique et Physiologique ou des Principes immédiats du corps de l'Homme et des Mammifères à l'état Normal et à l'état Pathologique, par MM. Ch. Robin, Professeur d'Anatomie Générale et de Microscopie, agrègé à la Faculté de Médecine de Paris, et Verdeil, Professeur de Chimie. 3 vols. 8vo. avec Atlas de 45 Planches gravées. Paris, 1853. Rodenberg. Memoire et Observations sur l'Accouchement Prématuré Artificiel. 8vo.	1	6 t 7 2 16	0 6 6 0 6
Méthodes employées en Belgique, en France, en Alemagne, et al Algebra, Parrachement et l'Extractiou des Minéraux Combustibles. Vol I. Royal 8vo. Atlas and Figures. Liège, 1852. Pouillet. Notions Générales de Physique et de Météorologie à l'usage de la Jeunesse. 12mo. 353 Cuts. Paris, 1850. Queriere. De l'Hygiène de l'Habitation. 8vo. Paris, 1851. Rackett. Observations on a Viper found in Cranborne Chace, Dorsetshire. 4to. London, 1817. (From Transactions.) Reveille-Parise. Traité de la Vieillesse Hygiénique, Médical et Philosophique, on Recherches sur l'Etat Physiologique, les Facultés morales, les Maladies de l'Age avancé, et sur les moyens les plus surs, les mieux experimentés de soutenir et de prolonger l'Activité Vitule à cette époque de l'existence. 8vo. Paris, 1853. Ricard. Le Magnétisme Traduit en Cour d'Assises. Acquittement. 8vo. Paris, 1845. Robin et Verdeil. Traité de Chimie Anatomique et Physiologique ou des Principes immédiats du corps de l'Homme et des Mammifères à l'état Normal et à l'état Pathologique, par MM. Ch. Robin, Professeur d'Anatomie Générale et de Microscopie, agrégé à la Faculté de Médecine de Paris, et Verdeil, Professeur de Chimie. 3 vols. 8vo. avec Atlas de 45 Planches gravées. Paris, 1853. Rodenberg. Mémoire et Observations sur l'Accouchement Prématuré Artificiel. 8vo. Paris, 1852. Rousseau et Lemonnier. Promenades au Jardin des Plantes. 12mo. With 5 1800.	1	6 t 0 7 2 16 16 1	0 6 6 0 6
Méthodes employées en Belgique, en France, en Alemagne, et angetetic, pour l'Arrachement et l'Extractiou des Minéraux Combustibles. Vol I. Royal 8vo. Atlas and Figures. Liège, 1852 Pouillet. Notions Générales de Physique et de Météorologie à l'usage de la Jeunesse. 12mo. 353 Cuts. Paris, 1850 Queriere. De l'Hygiène de l'Habitation. 8vo. Paris, 1851. Rackett. Observations on a Viper found in Cranborne Chace, Dorsetshire. 4to. London, 1817. (From Transactions.) Reveille-Parise. Traité de la Vieillesse Hygiénique, Médical et Philosophique, on Recherches sur l'Etat Physiologique, les Facultés morales, les Maladies de l'Age avancé, et sur les moyens les plus surs, les mieux experimentés de soutenir et de prolonger l'Activité Vitule à cette époque de l'existence. 8vo. Paris, 1853 Ricard. Le Magnétisme Traduit en Cour d'Assises. Acquittement. 8vo. Paris, 1845. Robin et Verdeil. Traité de Chimie Anatomique et Physiologique ou des Principes immédiats du corps de l'Homme et des Mammifères à l'état Normal et à l'état Pathologique, par MM. Ch. Robin, Professeur d'Anatomie Générale et de Microscopie, agrégé a la Faculté de Médecine de Paris, et Verdeil, Professeur de Chimie. 3 vols. 8vo. avec Atlas de 45 Planches gravées. Paris, 1853 Rodenberg. Mémoire et Observations sur l'Accouchement Prématuré Artificiel. 8vo. Paris, 1852 Rousseau et Lemonnier. Promenades au Jardin des Plantes. 12mo. With Plates. Paris, 1837		6 t 0 7 2 16 16 1	0 6 6 0 6
Méthodes employées en Belgique, en France, en Alemagne, et angetetic, pour l'Arrachement et l'Extractiou des Minéraux Combustibles. Vol I. Royal 8vo. Atlas and Figures. Liège, 1852 Pouillet. Notions Générales de Physique et de Météorologie à l'usage de la Jeunesse. 12mo. 353 Cuts. Paris, 1850 Queriere. De l'Hygiène de l'Habitation. 8vo. Paris, 1851. Rackett. Observations on a Viper found in Cranborne Chace, Dorsetshire. 4to. London, 1817. (From Transactions.) Reveille-Parise. Traité de la Vieillesse Hygiénique, Médical et Philosophique, on Recherches sur l'Etat Physiologique, les Facultés morales, les Maladies de l'Age avancé, et sur les moyens les plus surs, les mieux experimentés de soutenir et de prolonger l'Activité Vitule à cette époque de l'existence. 8vo. Paris, 1853 Ricard. Le Magnétisme Traduit en Cour d'Assises. Acquittement. 8vo. Paris, 1845. Robin et Verdeil. Traité de Chimie Anatomique et Physiologique ou des Principes immédiats du corps de l'Homme et des Mammifères à l'état Normal et à l'état Pathologique, par MM. Ch. Robin, Professeur d'Anatomie Générale et de Microscopie, agrégé a la Faculté de Médecine de Paris, et Verdeil, Professeur de Chimie. 3 vols. 8vo. avec Atlas de 45 Planches gravées. Paris, 1853 Rodenberg. Mémoire et Observations sur l'Accouchement Prématuré Artificiel. 8vo. Paris, 1852 Rousseau et Lemonnier. Promenades au Jardin des Plantes. 12mo. With Plates. Paris, 1837		6 h 0 0 7 2 2 16 16 1 3 3	0 6 6 0 6 0
Méthodes employées en Belgique, en France, en Alemagne, et angetetic, pour l'Arrachement et l'Extractiou des Minéraux Combustibles. Vol I. Royal 8vo. Atlas and Figures. Liège, 1852 Pouillet. Notions Générales de Physique et de Météorologie à l'usage de la Jeunesse. 12mo. 353 Cuts. Paris, 1850 Queriere. De l'Hygiène de l'Habitation. 8vo. Paris, 1851. Rackett. Observations on a Viper fouud in Cranborne Chace, Dorsetshire. 4to. London, 1817. (From Transactions.) Reveille-Parise. Traité de la Vieillesse Hygiénique, Médical et Philosophique, on Recherches sur l'Etat Physiologique, les Facultés morales, les Maladies de l'Age avancé, et sur les moyens les plus surs, les mieux experimentés de soutenir et de prolonger l'Activité Vitule à cette époque de l'existence. 8vo. Paris, 1853. Ricard. Le Magnétisme Traduit en Cour d'Assises. Acquittement. 8vo. Paris, 1845. Robin et Verdeil. Traité de Chimie Anatomique et Physiologique ou des Principes immédiats du corps de l'Homme et des Mammitères à l'état Normal et à l'état Pathologique, par MM. Ch. Robin, Professeur d'Anatomie Générale et de Microscopie, agrègé à la Faculté de Médecine de Paris, et Verdeil, Professeur de Chimie. 3 vols. 8vo. avec Atlas de 45 Planches gravées. Paris, 1853. Rodenberg. Memoire et Observations sur l'Accouchement Prématuré Artificiel. 8vo. Paris, 1852 Roisseau et Lemonnier. Promenades au Jardin des Plantes. 12mo. With Plates. Paris, 1837 Saline. A Memoir on the Birds of Greenland, with Descriptions and Notes on the Species observed in the late Voyage of Discovery in Davis's Straits and Baffin's Bay. 4to With a Plate. London, 1819. (From Transactions.)		6 t 0 7 2 16 1 16 1 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3	0 6 6 0 6 0 6
Méthodes employées en Belgique, en France, en Alemagne, et angetetic, pour l'Arrachement et l'Extractiou des Minéraux Combustibles. Vol I. Royal 8vo. Atlas and Figures. Liège, 1852 Pouillet. Notions Générales de Physique et de Météorologie à l'usage de la Jeunesse. 12mo. 353 Cuts. Paris, 1850 Queriere. De l'Hygiène de l'Habitation. 8vo. Paris, 1851. Rackett. Observations on a Viper found in Cranborne Chace, Dorsetshire. 4to. London, 1817. (From Transactions.) Reveille-Parise. Traité de la Vieillesse Hygiénique, Médical et Philosophique, on Recherches sur l'Etat Physiologique, les Facultés morales, les Maladies de l'Age avancé, et sur les moyens les plus surs, les mieux experimentés de soutenir et de prolonger l'Activité Vitule à cette époque de l'existence. 8vo. Paris, 1853 Ricard. Le Magnétisme Traduit en Cour d'Assises. Acquittement. 8vo. Paris, 1845. Robin et Verdeil. Traité de Chimie Anatomique et Physiologique ou des Principes immédiats du corps de l'Homme et des Mammifères à l'état Normal et à l'état Pathologique, par MM. Ch. Robin, Professeur d'Anatomie Générale et de Microscopie, agrégé a la Faculté de Médecine de Paris, et Verdeil, Professeur de Chimie. 3 vols. 8vo. avec Atlas de 45 Planches gravées. Paris, 1853 Rodenberg. Mémoire et Observations sur l'Accouchement Prématuré Artificiel. 8vo. Paris, 1852 Rousseau et Lemonnier. Promenades au Jardin des Plantes. 12mo. With Plates. Paris, 1837		6 h 0 0 7 2 2 16 16 1 3 3	0 6 6 0 6 0 6

Seringe. Flore des Jardins et des grands cultures; ou Description des Plantes de Jardins,	æ	S	d
d'Orangeries ou de Grands Cultures, leur Multiplication, &c. 3 vols. 8vo. With 20	,	_	
Plates. Paris, 1849. Sestier. Traité de l'Angine Laryngée Oedémateuse. 8vo. Paris, 1852	0	7	
Sheppard. On the Use of the Pedes Scansorii of Birds. 4to. With 1 Plate. London, 1818.			
(From Transactions.) Sieliel. Iconographie Opthalmologique, ou Description avec Figures Coloriées, des	0	1	6
Maladies de l'Organ de la Vue, comprenant l'Anatomie Pathologique, la Pathologie et la			
Thérapeutique Médico-chirurgicales. Large 4to. Parts 1 and 2. 8 Coloured Plates. Paris, 1852. Each	0	_	~
The above Work will be vablished in 20 Livraisons, composed each of 28 pages of	0	7	0
Text and 4 Coloured Plates; a Livraison to annear every 6 weeks			
Smith. Characters of two species of Tordylium. 4to. London, 1817. (From Transactions.) Rhynzomorphia Medullaris, &c. (From Transactions.) See COLEBROOKE.	0	0	6
Socouct. Principes d'Economie Médicale, ou des lois fondamentales de la Médicale.			
déduites de l'observation et de leur application au diagnostic ou prognostic, et au traite- ment des maladies. 8vo. Paris, 1852	^	,	C
Syphilisation (De la) et de la Syphilis Constitutionnelle Communication à l'Académia	0	4	O
Nationale de Médecine, par MM. Auzias, Ricord, Begin, Malgaigne, Velpeau, Dupoul, Larrey, Lagneau, Levy, Gerdy. 8vo. Paris, 1852			
Tardien. Voirées et Cimetières. 8vo. Paris, 1852	0	3	
Dictionnaire d'Hygiène publique et de Salubrité ou Réportaire de toute le			
questions relatives à la Santé Publique. Vol. I. 8vo. Paris, 1852. Tenminck et Langier. Nouveau receuil de planches coloriées d'Oiseaux, pour	0	8 (0
servir de suite et de complément aux planches enluminés de Buffon. Par C. J.			
L'ouvrage se compose de 5 vols., avec 600 planches folio, dessinées d'après poture			
par fretie et fluet, gravees et colorides			
Chaque livraison contient 6 planches, et le texte descriptif correspondant. Ouvrage publié en 102 livraisons, et qui est maintenant complet.			
Prix de la livraison, folio, figures coloriées, au lieu de 15s., réduit à	0 1	0 (0
Grand 4to. au lieu de 10s. 6d., réduit à La dernière livraison contient des tables scientifiques et méthodiques.		7 6	-
Des Oiseaux d'Europe, décrits par C. J. Temminck Atlas de 530 plunches dessinées			
2 vols. Svo. figures coloriées cartonnée	_		
2 Vols. 8vo. figures noires cartonnés	5 1 1	0 (
Il ne reste qu'ini très petit nombre d'exemplaires de cet important ouvrage, dont nous avons baissé le prix de plus des trois quarts.			
AUSTE. Le Magnétisme Animal expliqué ou Lecons Applytiques sur la nature acceptable			
	0	7 ()
Triquet. Abrégé de Pathologie Médico-Chirurgicale ou resumé Analytique de Médecine et de Chirurgie. Vol. I. 8vo. Paris, 1852	۵	G (
Turek. De la Vieillesse, étudiée comme Maladle, et des moyens de la combattre. 8vo. Paris, 1852	v	6 (,
Victor. Description de quelques Colean(Area reconillies	0	4 ()
Voisin. Du Traitement Intelligent de la Folie, et application de quelquesunc de ses principes à la reforme des Criminels. 8vo. Paris, 1847			
raris, 184/	0	2 6	5
CEDATA			
GERMAN WORKS.			
Abhandlungen der Naturhistorischen Gesellschaft zu Nürnberg. 8vo. Part 1. With 3 Plates. Nürnberg, 1852			
3 Plates, Nürnberg, 1852 Apell. Die Reformation der Stevelunde Wie P. t.	0 8	3 6	
8vo. With 5 Plates. Jena 1852			
Bendz, Icones Anatomica Vulgurum Danicoman Ar	0 12	: 6	
	2 2	0	
Willistatt, Jahresbericht fiber die Frankeit ist in der	0 3	0	
Wissenschaften im Jahre 1851. Royal 8vo. Würzburg, 1852	0 5	0	
periode Authorogic und Amerapie. Band V Lief Borrel One 17.1		6	
Bonn, 1852	2 ~	^	
organischen Lebens zu den vulkanischen Masser die Beziehungen des kleinsten) 7	0	
Frev. Die Hartgebilde der Winderston William IV.) 1	6	
Carl Nagell. 8vo. With 2 Plates. Zürich, 1852	0 2	0	
Willi 12 Plates Tene 1960	17		

	£	5	a	
manschen watering watering and ballion of the control of the contr	0	1	0	
	0	10	6	
Huss. Chronische Aikoholskrankheit oder Alcobolismus Chronicus. Ein Beltrag zur Kenntniss der Vergiftungs-Krankheiteu, nach elgener und anderer Erfahrung. 8vo.	0	10	6	
Kissel (Dr. Carl.) Die direkte Kunstheilung der Pneumonien-Monographie. 8vo.	0		0	
Kolliker. Mikroscopische Anatomie oder Gewebelehre des Menschen. Royal 8vo.				
Vol. 11. Parts 1 and 2. With 295 Cuts. Leipzig, 1852 Kreutzer. Grundriss der Veterinärmedezln zum Gebrauche hei Vorlesungen für Studi-	1	3	0	
rende der Medezin, sowie zum Selbstgebrauche für Physikat- und Praktische Aerzte und Thierärzte. Royal 8vo. Parts 1 and 2. Erlangen, 1852	0		6	
Kutzing. Species Algarum. 8vo. Lipsiæ, 1849 Tabulæ Phycologiæ oder Abbildungen der Tange. 8vo. Vol. I. With 100 Plates, Plain. Nordhausen, 1850	2		0	
Lehrbuch der Kriegsbaukunst zum Gebrauche der Kais. Kön. Genie-Akademie. Lief. 1.	1	7	0	
Leygh. Handhuch der Anatomie der Hausthiere. 8vo. With 200 Woodcuts. Stutt-	0	14	9	
Liebig und Kopp. Jahresbericht über die Fortschritte der reinen Pharmaceutischen und Technischen Chemie, Physik, Mineralogie, und Geologie. 1847—51. 3 Parts. 8vo. Lotze. Medecinische Psycologie oder Physiologie der Seele. 8vo. Leipzig, 1852		11		
Lutgert. Tabula Medicina Historica, in ordinem Chronologicum et Philosophicum. Folio. With 8 Plates in a Portfolio. Lugduni Batavorum, 1852	1	0	0	
Mischler. Das Deutsche Eisenhüttengewerhe vom Standpunkte der Staatswirtnschaft.	0	10	0 ()
Muller. Ueber die Larven und die Metamorphose des Echinodermen. 4to. With 9 Plates.	0	7	6	;
Naumannia. Archiv für die Ornithologie, vorzugsweise Europa's. Organ der Deutschen Ornithologen Gesellschaft. Herausgegehen von E. Baldamus. 8vo. Vol. I. 1851. Vol. II. Part 1. 1852	0	10	0 6	
Puggaard. Geologie der Insel Man, eine Untersuchung über die Umwälzungen der Kreide und der Glacialbildung, sowie über die Quaternären Ablagerungen und der Erratischen Blöcke dieser Insel. 8vo. With 13 coloured Plates and numerous Woodcuts. Leipzig, 1852	0		5 6	j
Rammelsberg. Lehrhuch der Krystallkunde oder Anfangsgründe der Krystallographie, Krystallonhygik und Krystallochemie. Royal 8vo. 250 Cuts. Berlin, 1852	0	10	0 ()
Reichel. Die Basalte und Säulenförmigen Sandsteine der Zittauer Gegend in Sächsen	0	1.	4 ()
der Vollständigsten Kupfersammlung der Vögel aller Weltheile. 4to. Parts 1 and 2.	0)	7 (0
Rengott. Uebersicht der Resultate Mineralogischer Forschungen in den Jahren 1844	()	8	6
Ræmer (Dr. F.) Monographie der Fossilen Crinoidienfamme der Blastoiden und der	()	1 (6
Rose. Ausführliches Handbuch der Analytischen Chemie. 8vo. Vol. I.—Die Lehre von den Qualitativen Chemisch-Aualytischen. Vol. II.—Die Lebre von den Quantitativen Chemisch-Analytischen. Braunschweig, 1851.	1	1	8	
Rothamel. Die Wallungen und Stockungen des Bintes. 8vo. Cassell, 1802)	4	
1851. Erstes Halbjahr, Hit. I and 2. Royal 8vo. Göttingen, 1852	(0	7	0
Ruppell. Systematische Uebersicht der Voger Nordost Arma 3.		1 1	16	0
Salisbury. On Lycopodium Denticulatum. 4to. with Plate. See Colebrooke. Schacht. Die Pflanzenzelle, der innere Bnu und das Leben der Gewächse. Mit elgenen Schacht. Die Pflanzenzelle, Chemischen Untersuchungen. Royal 8vo. 20 Plates,				
vergleichenden Mieroscopisch-Chemischen Checkschen		1	4	0
Scheerer. Lehrbuch der Metallurgie, mit besonderer Hillstellt am Oktaber 1117 Cuts. Braunschweig, 1486–48		0	12	9
Schmidt. Beiträge zur Flora der Cap Verdischen Inseln. Mit Berdenstehtigung auch signer jetzt daselbst bekannten wildwachsenden und kultivirten Pflanzen. 8vo. Heidelberg.		0	5	0
Steen-Billes. Bericht über die Relse der Corvette 'Galathea' um die Wettin den Vanschen übersetzt, und thellwelse bearheitet von Rosen. 8vo.		0	12	6
Vol. I. With 7 Plates. Appennagen, 1002		0	8	6
(Dr. F.) Versuch einer Geschichte der Pflanzenwelt. 8vo. Wien, 1852				

Wahar Die Tertierfore der Niedercheinischen Braun Kehlenformetien. 4th Will C	£	s d
Weber. Die Tertiarflora der Niederrheinischen Braun-Kohlenformation. 4to. With 8 Plates. Cassel, 1852	0 1	7 0
Weisenberg. Vollständiges Handwörterbuch der gesammten Arzneimittel von der ältesten bis auf die neueste Zeit für Aerzte und Studirte Wundürzte. 8vo. Part 1.		
Jena, 1852	0	1 6
With 6 Plates. Berlin, 1847	0	6 6
MISCELLANEOUS WORKS.		
Babinet. Elémens de Géométrie descriptive. 8vo. with Atlas. Paris, 1850	0	PT 0
Bertrand. Traité Elémentaire d'Algèbre. 8vo. Paris, 1851.		7 0
Bonnefonx. Manœuvrier Complet ou Traité des Manœuvres de Mer, soit à bord des Batiments à Voiles ou à Vapenr. 8vo. Parls, 1852.	0	7 0
Bonillet (M. N.) Dictionnaire Universel d'Histoire et de Géographie contenant	0 7	7 0
1. L'Histoire Proprement Dite.—2. La Biographie Universel.—3. La Mythologie.—4. La Géographie Ancienne et Moderne. Royal Svo. rel. Paris, 1852.	1 10	0.0
Chasies. Traité de Géométrie Supérieure. 8vo. With 12 Plates. Paris, 1852	0 1	
Cirodde (P. L.) Leçons de Géométrie Analitique, précédées des Eléments de la Trigo- nométrie, Rectilique et Sphérique. 2de Edition. 8vo. Paris, 1848	0 2	7 G
Comte. Système de Politique Positive. 8vo. Vois. I. and II. Paris. 1851—52	0 1	
De Laperrière. Barème ou Tableaux des Comptes faits, d'après le Système Métrique pour le calcul des surfaces et des cubes des Bois carrés, Bois ronds, &c. 8vo. Bordeaux,		
Desmarest. Théorle des nombres. Traité de l'Analyse indéterminée du second des	0 (0 8
à deux inconnues; suivi de l'application de cette auaiyse à la recherche des Racines primitives avec une table de ces Racines, &c. 4to. Paris, 1852		
Flandin. Voyage en Perse. Svo. Relation du Voyage. 2 vols Paris 1852	0 16	0 0
Industriel et Social à l'insage de ceux qui recherchent des rensairements propriété de l'insage de ceux qui recherchent des rensairements propriété de l'insage de ceux qui recherchent des rensairements propriété de l'insage de ceux qui recherchent des rensairements propriété de l'insage de ceux qui recherchent des rensairements propriété de l'insage de ceux qui recherchent des rensairements propriété de l'insage de ceux qui recherchent des rensairements propriété de l'insage de ceux qui recherchent des rensairements propriété de l'insage de ceux qui recherchent des rensairements propriété de l'insage de ceux qui recherchent des rensairements propriété de l'insage de ceux qui recherchent des rensairements propriété de l'insage de ceux qui recherchent des rensairements propriété de l'insage de ceux qui recherchent des rensairements propriété de l'insage de ceux qui recherchent des rensairements propriété de l'insage de ceux qui recherchent des rensairements propriété de l'insage de ceux qui recherchent des rensairements propriété de l'insage de ceux qui recherchent des rensairements propriété de l'insage de ceux qui recherchent des rensairements propriété de l'insage de ceux qui recherchent de l'insage de ceux qui recherchent de l'insage de l'insag		
	0 7	7 6
Guizot. Etudes sur les Beaux. Arts en général. 8vo. Paris, 1852 Corneille et son temps. Etude Littéraire. 8vo. Paris, 1852		0 6
Gnynemer. Dictionnarie d'Astronomie à l'Usage des Cons du Monde d'annèe l'		0
	0 4	10
Lamotte. Cours Méthodique de Dessin Linéaire et de Géométrie, usuelle applicable à tous les Modes d'Enseignement. 2 Parts, 8vo. with Atlas, 4to. Paris, 1843	0.14	
Lepsius. Briefe aus Aegypten, Aethiopien und der Halbinsei des Sinai. 8vo. Plates. Berlin, 1852	0 14	U
Lesnenr. Description of several new Species of Helethure 10-	0 10	6
vents, et autres monuments publics et portivuliere la vents, et autres monuments publics et portivuliere la vents, et autres monuments publics et portivuliere la vents de la	1	
Rome. 2 vols. 4to. and Atlas, folio. de 231 Planches. Relié. Paris, 1850—52.	3 0	0
Nonrrisson. Essai sur la Philosophie de Bossuet, avec des fragments inédits. 8vo.	Δ.	
Nouvelle. Biographie Universelle, depuis les temps les plus reculés jnsqu'à nos jours avec les renseignements Bibliographiques et l'indication des sources à consuiter. Publiée par Firmin Didot, Frères, sous la direction de M. le Dy Hoefen de	0 4	6
par Firmin Didot, Frères, sous la direction de M. le Dy Hessources a consuiter. Publice	0 10	6
Pinaud. Programme d'un Cours Eliments. 2 vois. 8vo. aud Atlas, 4to. Paris, 1847	0 15	
d'Instruction. 8vo. Toulouse, 1851	0 6	6
Edition. Parls 1847.		
Résolution des Problèmes de Géométrie 2000 Edit	0 5	0
Sainte-Benye, Causerles du Lundi 12mg 5 miles. Paris, 184/.	0 - 5 $0 - 17$	0
ment en Europe et des Arts qui es au 1/e Siècle. Histoire du Costume et de l'Ameuble-	0 17	U
This Work will appear every 10 days, and will comprise 180 to 200 Parts, each Part to contain 2 Plates in Gold, Silver or Colours, 14 are out price care to 200 Parts, each Part to	0 1	G
Sounet. Algebre Elementaire avec do nombre cut, price euch, 18. 6d. 4to. Paris, 1852.		
Tarnier (E. A.) Trotté d'Alabam (16.	0 6	0
cercise, et faisant suite à l'Arlebmétique de la compagné d'un grand nombre d'Ex-	0 ~	C
Tresca. Traité élémentaire de Cémario de la Russie. 2 vols. 8vo. Paris, 1852	0 - 7 $0 - 16$	6
programme d'admission à l'école Polytagiant de redige conformément au dernicr		
S S S S S S S S S S S S S S S S S S S	0 7	6

STANDARD SCIENTIFIC WORKS.

Already Published.

Professor Muller's Principles of Physics and Meteorology. WITH 530 WOODCUTS AND 2 COLOURED ENGRAVINGS. 8vo. 18s.

Professor Weisbach's Mechanics of Machinery and Engineering. 2 vols. WITH 900 WOODCUTS. £1 198.

III. IV. AND X.

Professor Knapp's Technology;

Chemistry Applied to the Arts and Manufactures. EDITED BY DR. RONALDS, AND DR. T. RICHARDSON. 3 VOLS. SPLENDIDLY ILLUSTRATED, PRICE £3 4s.

Quekett's (John) Practical Treatise on the Use of the Microscope.

SECOND EDITION, WITH 12 STEEL AND NUMEROUS WOOD ENGRAVINGS. 8vo. £1 2s.

Professor Fau's Anatomy of the External Forms of Man. FOR ARTISTS.

EDITED BY R. KNOX, M.D.

8vo. and an atlas of 28 plates 4to. Plain £1 4s. coloured £2 2s.

Professor Graham's Elements of Chemistry, WITH

Its Application in the Arts.

SECOND EDITION, WITH NUMEROUS WOODCUTS. VOL. I. £1 18.

Professor Nichol's Architecture of the Heavens. NINTH EDITION, WITH 23 STEEL PLATES AND MANY WOODCUTS. LONDON, 1851. 168.

THE PRESCRIBERS' COMPLETE HANDBOOK,

Comprising the Principles of the Art of Prescribing, with a List of Discases and their appropriate Remedies: a Materia Medica, containing all the principal Medicines employed, Classified according to their Natural Families, with their Properties, Preparations, and Uses; and a Concise Sketch of Toxicology.

BY M. TROUSSEAU, PROFESSOR OF THE FACULTY OF MEDICINE, PARIS; AND M. REVEIL. EDITED, WITH NOTES, BY J. BIRKBECK NEVINS, M.D. 12mo. London, 1852. 6s. 6d.

THE PLANT; A BIOGRAPHY,

IN A SERIES OF THIRTEEN POPULAR LECTURES ON BOTANY. BY M. J. SCHLEIDEN, M.D.

EDITED AND TRANSLATED BY A. HENFREY, F.L.S. 8vo. Second Edition, much Enlarged. With 7 Coloured Plates and 16 Woodcuts. London, 1852. 15s.

LECTURES ON HISTOLOGY,

DELIVERED AT THE ROYAL COLLEGE OF SURGEONS OF ENGLAND. BY J. QUEKETT.

Illustrated by 159 Woodcuts. Svo. London, 1852. 10s. 6d.

NATURAL AND MESMERIC CLAIRVOYANCE,

WITH THE PRACTICAL APPLICATION OF MESMERISM IN SURGERY AND MEDICINE. BY JAMES ESDAILE, M.D., LATE PRESIDENCY SURGEON, CALCUTTA; AUTHOR OF "MESMERISM IN INDIA," &c. 12mo. London, 1852. 4s. 6d.

London: Printed by Schulze and Co., 13, Poland Street.





(Walker Ami) R6 1852 - Q 52

1.0

